



Københavns Universitet

Reconstructing the early evolution of Fungi using a six-gene phylogeny.

James, Timothy Y; Kauff , F ; Schoch , C; Matheny, PB; Hoffstetter, V; Cox, C; Celio, G; Gueidan, C; Fraker, E; Miadlikowska, J; Lumbsch, TH; Rauhut, A; Reeb, V; Arnold, AE; Wynns, Anja Amtoft; Stajich, J; Hosaka , K; Sung , G; Johnson, D; O'Rourke, B; Crockett, M; Binder, M; Curtis, JM; Slot , J; Wang, Z; Wilson , A; Schüßler , A; Longcore, JE; O'Donnell , K; Mozley-Standridge , S; Porter , D; Letcher, PM; Powell, MJ; Taylor , JW; White, MM; Griffith , GW; Davies, DR; Humber, RA; Morton , JB; Sugiyama, J; Rossman , A; Rogers , JD; Pfister , DH; Hewitt, D; Hansen, K; Hambleton , S; Shoemaker, RA; Kohlmeyer, J; Volkmann-Kohlmeyer , B; Spotts, RA; Serdani , M; Crous, PW; Hughes, KW; Matsuura, K; Langer, E; Langer, G; Untereiner , W; Lücking, R; Büdel , B; Geiser , DM; Aptroot , A; Diederich, P; Schmitt, I; Schultz , M; Yahr , R; Hibbett, D; Lutzoni , F; McLaughlin, DJ; Spatafora, JW; Vilgalys, R

Published in:
Nature

Publication date:
2006

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
James, T. Y., Kauff , F., Schoch , C., Matheny, PB., Hoffstetter, V., Cox, C., ... Vilgalys, R. (2006). Reconstructing the early evolution of Fungi using a six-gene phylogeny. *Nature*, 443, 818-822.

ARTICLES

Reconstructing the early evolution of Fungi using a six-gene phylogeny

Timothy Y. James¹, Frank Kauff¹, Conrad L. Schoch^{2*}, P. Brandon Matheny^{3*}, Valérie Hofstetter^{1*}, Cymon J. Cox^{1†}, Gail Celio⁴, Cécile Gueidan¹, Emily Fraker¹, Jolanta Miadlikowska¹, H. Thorsten Lumbsch⁵, Alexandra Rauhut⁶, Valérie Reeb¹, A. Elizabeth Arnold^{1†}, Anja Amtoft⁷, Jason E. Stajich⁸, Kentaro Hosaka^{2†}, Gi-Ho Sung², Desiree Johnson², Ben O'Rourke², Michael Crockett², Manfred Binder³, Judd M. Curtis³, Jason C. Slot³, Zheng Wang^{3†}, Andrew W. Wilson³, Arthur Schüßler⁹, Joyce E. Longcore¹⁰, Kerry O'Donnell¹¹, Sharon Mozley-Standridge¹², David Porter¹², Peter M. Letcher¹³, Martha J. Powell¹³, John W. Taylor¹⁴, Merlin M. White¹⁵, Gareth W. Griffith¹⁶, David R. Davies¹⁷, Richard A. Humber¹⁸, Joseph B. Morton¹⁹, Junta Sugiyama²⁰, Amy Y. Rossman²¹, Jack D. Rogers²², Don H. Pfister²³, David Hewitt²³, Karen Hansen²³, Sarah Hambleton²⁴, Robert A. Shoemaker²⁴, Jan Kohlmeyer²⁵, Brigitte Volkmann-Kohlmeyer²⁵, Robert A. Spotts²⁶, Maryna Serdani²⁶, Pedro W. Crous²⁷, Karen W. Hughes²⁸, Kenji Matsuura²⁹, Ewald Langer³⁰, Gitta Langer³⁰, Wendy A. Untereiner³¹, Robert Lücking⁵, Burkhard Büdel⁶, David M. Geiser³², André Aptroot³³, Paul Diederich³⁴, Imke Schmitt^{5†}, Matthias Schultz³⁵, Rebecca Yahr^{1†}, David S. Hibbett³, François Lutzoni¹, David J. McLaughlin⁴, Joseph W. Spatafora² & Rytas Vilgalys¹

The ancestors of fungi are believed to be simple aquatic forms with flagellated spores, similar to members of the extant phylum Chytridiomycota (chytrids). Current classifications assume that chytrids form an early-diverging clade within the kingdom Fungi and imply a single loss of the spore flagellum, leading to the diversification of terrestrial fungi. Here we develop phylogenetic hypotheses for Fungi using data from six gene regions and nearly 200 species. Our results indicate that there may have been at least four independent losses of the flagellum in the kingdom Fungi. These losses of swimming spores coincided with the evolution of new mechanisms of spore dispersal, such as aerial dispersal in mycelial groups and polar tube eversion in the microsporidia (unicellular forms that lack mitochondria). The enigmatic microsporidia seem to be derived from an endoparasitic chytrid ancestor similar to *Rozella allomycis*, on the earliest diverging branch of the fungal phylogenetic tree.

Fungi, Viridiplantae and Animalia are all large clades descended from unicellular, flagellated, aquatic forms that radiated extensively on land. For both plants and animals, biologists have developed unified hypotheses regarding the evolution of morphology and ecology from ancestral to highly derived traits. For example, among green plants, morphologically simple photosynthetic forms, such as unicellular

green algae, gave rise to multicellular forms such as bryophytes, and were followed by a radiation of complex flowering forms with highly derived sexual mechanisms at the tips of the plant phylogeny^{1,2}. Similarly, animals seem to have evolved increasingly complex tissue systems and development from a simple, flagellated, protist-like ancestor similar to extant Choanoflagellida³.

¹Department of Biology, Duke University, Durham, North Carolina 27708-0338, USA. ²Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon 97331-2902, USA. ³Department of Biology, Clark University, Worcester, Massachusetts 01610, USA. ⁴Department of Plant Biology, University of Minnesota, Saint Paul, Minnesota 55108, USA. ⁵Field Museum of Natural History, Chicago, Illinois 60605-2496, USA. ⁶Fachbereich Biologie, Abteilung Pflanzenökologie und Systematik, 67653 Kaiserslautern, Germany. ⁷Institute of Systematic Botany, New York Botanical Garden, Bronx, New York 10458-6126, USA. ⁸University Program in Genetics and Genomics, Duke University, Durham, North Carolina 27708-0338, USA. ⁹Institute of Botany, Darmstadt University of Technology, D-64287 Darmstadt, Germany. ¹⁰Department of Biological Sciences, University of Maine, Orono, Maine 04469, USA. ¹¹National Center for Agricultural Utilization Research, USDA Agricultural Research Service, Peoria, Illinois 61604, USA. ¹²Department of Plant Biology, University of Georgia, Athens, Georgia 30605, USA. ¹³Department of Biological Sciences, University of Alabama, Tuscaloosa, Alabama 35487, USA. ¹⁴Department of Plant and Microbial Biology, University of California, Berkeley, California 94720, USA. ¹⁵Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, Kansas 66045-7534, USA. ¹⁶Institute of Biological Sciences, University of Wales, Aberystwyth, Ceredigion SY23 3DA, UK. ¹⁷Institute of Grassland and Environmental Research, Plas Gogerddan, Aberystwyth, Ceredigion SY23 3EB, UK. ¹⁸United States Plant, Soil and Nutrition Laboratory, USDA-ARS Plant Protection Research Unit, Ithaca, New York 14853-2901, USA. ¹⁹Division of Plant and Soil Sciences, West Virginia University, Morgantown, West Virginia 26506-6057, USA. ²⁰TechnoSuruga, Chiyoda-ku, Tokyo 101-0052, Japan. ²¹Systematic Botany and Mycology Laboratory, USDA Agricultural Research Service, Beltsville, Maryland 20705, USA. ²²Department of Plant Pathology, Washington State University, Pullman, Washington 99164, USA. ²³Harvard University Herbaria, Cambridge, Massachusetts 02138, USA. ²⁴Biodiversity (Mycology and Botany), Agriculture and Agri-Food Canada, Ottawa, Ontario K1A 0C6, Canada. ²⁵Institute of Marine Sciences, University of North Carolina at Chapel Hill, Morehead City, North Carolina 28557, USA. ²⁶Mid-Columbia Agricultural Research and Extension Center, Oregon State University, Hood River, Oregon 97031, USA. ²⁷Centraalbureau voor Schimmelcultures, Fungal Biodiversity Centre, 3508 AD Utrecht, The Netherlands. ²⁸Botany Department, University of Tennessee, Knoxville, Tennessee 37996, USA. ²⁹Faculty of Agriculture, Okayama University, Okayama 700-8530, Japan. ³⁰Institut für Biologie, Universität Kassel, D-34132 Kassel, Germany. ³¹Department of Botany, Brandon University, Brandon, Manitoba R7A 6A9, Canada. ³²Department of Plant Pathology, Penn State University, University Park, Pennsylvania 16802, USA. ³³Adviesbureau voor Bryologie en Lichenologie, NL-3762 XK Soest, The Netherlands. ³⁴Musée national d'histoire naturelle, L-2160 Luxembourg. ³⁵Biozentrum Klein Flottbek und Botanischer Garten, Universität Hamburg, Systematik der Pflanzen, D-22609 Hamburg, Germany. †Present addresses: Biometry and Molecular Research, Department of Zoology, Natural History Museum, London SW7 5BD, UK (C.J.C.); Department of Plant Sciences, University of Arizona, Tucson, Arizona 85721, USA (A.E.A.); Department of Botany, The Field Museum, Chicago, Illinois 60605-2496, USA (K.H.); Department of Biological Sciences, Roy J. Carver Center for Comparative Genomics, University of Iowa, Iowa City, Iowa 52242, USA (Z.W.); Leibniz Institute for Natural Product Research and Infection Biology, Hans-Knöll-Institute, D-07745 Jena, Germany (I.S.); Royal Botanic Garden Edinburgh, Edinburgh EH3 5LA, UK (R.Y.).

*These authors contributed equally to this work.

Currently, no accepted phylogenetic hypothesis exists for the evolution of form and nutritional mode for the earliest fungi. Traditional views of fungal phylogeny indicate that fungi with flagellated cells (Chytridiomycota) are the sister group of the remaining phyla of non-flagellated fungi (Zygomycota, Glomeromycota, Ascomycota and Basidiomycota), implying a single loss of the flagellum coincident with a shift to land. Key adaptations to the terrestrial habit in the fungi include the evolution of a filamentous growth form and the development of aerielly dispersed spores. However, recent phylogenetic studies question the monophyly of the basal phyla Chytridiomycota and Zygomycota^{4,5}. Resolving the phylogeny of the basal groups of the Fungi and their relationships to Ascomycota and Basidiomycota is necessary to understand the sequence of events leading to the colonization of land and the evolution of terrestrial ecosystems. Here we present a multilocus phylogeny of the kingdom Fungi, including representatives of all currently recognized phyla. This analysis provides a robust kingdom-level phylogeny and suggests that there were at least four independent losses of flagella during the early evolution of the Fungi.

We estimated the phylogeny of the Fungi using data from six gene regions: 18S rRNA, 28S rRNA, 5.8S rRNA, elongation factor 1- α (*EF1 α*), and two RNA polymerase II subunits (*RPB1* and *RPB2*). Incongruence among gene regions was tested by maximum likelihood bootstrap (MLBS) analyses of each data partition. This strategy allowed us to identify potential contaminant sequences in addition to conflicting phylogenetic signal. Very little conflicting signal among genes was detected, allowing construction of one super-matrix combining the data for all six gene regions for 199 fungal taxa, 29 of which used data from genome sequencing projects (Supplementary Notes 1). Only 6% of the cells in the super-matrix were missing data, and the number of aligned nucleotides was 6,436. The data were analysed by bayesian methods using a heterogeneous amino-acid and nucleotide model (see Supplementary Notes 2 for a nucleotide-only analysis). Support was estimated at nodes by bayesian posterior probabilities (BPP), MLBS and analysis of individual gene partitions (Supplementary Notes 3).

Chytridiomycota is not monophyletic

The combined gene phylogeny of the Fungi supported monophyly of the Ascomycota, Basidiomycota and Glomeromycota (Fig. 1). The Ascomycota and Basidiomycota formed a clade of 'dikarya' (that is, fungi characterized by having a portion of their life cycle with paired nuclei). Phylogenetic analyses also supported, by BPP, a clade uniting the dikarya and Glomeromycota, in agreement with previously published 18S rRNA phylogenies^{6,7}. The opisthokont clade (Fungi, Metazoa and Choanoflagellida) was also recovered, as has been reported in other studies^{3,8,9}. Two unexpected results were the placements of the endoparasitic, spizellomycetalean chytrids *Olpidium brassicae* and *R. allomycis*. *Olpidium brassicae* grouped with the Zygomycota as sister taxon to *Basidiobolus ranarum*, and *R. allomycis* grouped with the microsporidia as the earliest diverging branch of the Fungi.

The phylum Chytridiomycota consists of true fungi that produce flagellated spores (zoospores). On the basis of ultrastructural studies, the chytrid zoospore is homologous to that of non-fungal opisthokonts¹⁰. The ultrastructural complexity of the opisthokont zoospore suggests that it has evolved only once. Because the zoospore is an ancestral trait, Chytridiomycota is solely defined on a shared ancestral trait (symplesiomorphy) rather than a shared derived trait (synapomorphy). Our phylogeny indicates that the Chytridiomycota is polyphyletic (Fig. 1), consisting of early diverging lineages that have retained the zoospore. However, one large clade of Chytridiomycota uniting the orders Chytriales, Monoblepharidales, Neocallimastigales and some Spizellomycetales (which we call the 'euchytrids') is recovered with high support values in the combined analysis as well as in multiple, single-gene-based analyses (Fig. 1 and Supplementary Notes 3).

In the present phylogeny (Fig. 1), six losses of the flagellum are inferred to have occurred during the evolution of the Fungi. Ancestral state reconstruction of the presence or absence of the flagellum along the phylogeny for each of the 58,611 credible trees demonstrated 4–6 losses (mean 5.86) of the flagellum within the Fungi. One well-supported loss took place along the branch leading to *Hyaloraphidium curvatum*, a unique fungus that grows superficially like a unicellular planktonic alga¹¹. A second loss occurred in the lineage leading to the microsporidia and 2–4 losses occurred among Zygomycota. Variation in the number of losses of the flagellum is attributable, in part, to the uncertain placement of *O. brassicae* and members of the microsporidia. Rearrangement of the phylogenetic position of *O. brassicae* and microsporidia can create phylogenies requiring only two or three losses of the flagellum; however, each of these alternative phylogenies is rejected as statistically worse (in likelihood; $P < 0.05$) than that shown in Fig. 1.

Most molecular phylogenies of the Fungi based on 18S rDNA have placed the zygomycete *Basidiobolus* among Chytridiomycota^{4,12}. This placement indicated that *Basidiobolus* might have made the transition recently from a zoospore state, and that an independent loss of a flagellum occurred in this lineage¹². This argument was strengthened by the presence in two *Basidiobolus* species of a ring-shaped spindle pole body that contains 11–12 singlet microtubules similar to a centriole, but lacks centriolar ninefold symmetry¹³. Our phylogeny is the first to place *Basidiobolus* close to Entomophthorales, the order within which it has been classified traditionally and to which it is ecologically and morphologically allied¹⁴ (for additional phylogenetic support from a paralogous copy of *EF1 α* , see Supplementary Notes 4). Unexpectedly, the phylogeny also suggests a relationship between *B. ranarum* and the chytrid *O. brassicae* (Fig. 1). A functional link between the two taxa is unclear: *O. brassicae* is an endoparasite of plant roots, whereas *Basidiobolus* is associated with insects, soil and amphibians.

Phylogenetic position of the microsporidia

Microsporidia are obligately endoparasitic, protist-like organisms with highly reduced morphology and genomes¹⁵. A defining characteristic of these parasites is the elaborate mechanism by which the spore contents are rapidly injected into the host's cytoplasm through a thin polar tube. Placement of microsporidia in the tree of life has been problematic owing to their extremely accelerated rate of sequence evolution. The earliest phylogenetic analyses of 18S rRNA placed the microsporidia among the earliest diverging lineages of eukaryotes¹⁵; however, these analyses now seem to have been an artefact of 'long branch attraction' of microsporidia to the base of the phylogeny¹⁵. More recent results using *RPB1*, α - and β -tubulin, and other genes, have suggested a fungal origin of the microsporidia^{16–18}, a placement consistent with their having the shared traits of closed mitosis and spores that contain chitin and trehalose¹⁹. Only one study has placed the microsporidia with a specific fungal lineage, in which a relationship was demonstrated between members of the Zygomycota and microsporidia by using tubulin proteins¹⁸. However, tubulin proteins seem to have evolved at different rates in flagellated and non-flagellated fungi^{18,20}.

The microsporidia and *R. allomycis* are intracellular parasites of primarily animals and fungi, respectively. A similarity between microsporidia and *R. allomycis* is the absence of a cell wall when invading host cells, such that the plasma membrane of the parasite makes direct contact with the cytoplasm of the host cell^{19,21}. Although *R. allomycis* does not seem to occupy a long phylogenetic branch, we tested whether the placement of microsporidia with *R. allomycis* was due to long branch attraction. Two different methods suggested that the relationship between microsporidia and *R. allomycis* is not due to long branch attraction (see Supplementary Notes 5). We also tested whether alternative placements for the microsporidia could be statistically rejected from the maximum likelihood phylogeny shown in Fig. 1 using the approximately unbiased test²². Alternative placements of microsporidia with Fungi that have been suggested




Figure 1 | Phylogeny of the kingdom Fungi using bayesian analysis of the combined, six-gene data set. Each fungal species begins with a unique 'Assembling the Fungal Tree of Life' identifier, followed by genus and species. Indicated for each terminal taxon are: nutritional mode, whether they produce flagellated cells and if there is a genome sequence for the taxon completed or underway. Thickened branches indicate those that are supported both by heterogeneous bayesian analysis (BPP $\geq 95\%$) and by MLBS ($\geq 70\%$). Almost every branch was supported by BPP and thus values are not shown. Where indicated, support values (percentage of trees in

agreement out of 58,611 trees) indicate BPP followed by MLBS. Branches are shaded according to reconstruction of nutritional mode. Microsporidia branches have been shortened three times (double black break) to increase readability. Red vertical ticks on branches indicate alternative placements of microsporidia that might be significantly rejected ($P < 0.05$) and green ticks indicate placements that cannot be rejected. Quotation marks indicate non-monophyly of the taxon. The name 'Mucormycotina' will be validated in a manuscript that is in preparation.

include: a sister relationship to the dikarya²³; sister to the zygomycete order Entomophthorales¹⁸; and among the harpellid Trichomycetes¹⁹, represented here by *Smittium culisetae*. We were able to reject ($P < 0.05$) nine alternative placements of the microsporidia (red vertical ticks in Fig. 1), including early divergences among eukaryotes. However, we were unable to reject a placement of microsporidia as sister to Entomophthorales, as sister to the blastocladalean chytrids, as sister to the zygomycete *Dimargaris*, as sister to dikarya and as sister to the Fungi (green vertical ticks in Fig. 1).

Taken together, our results suggest that the relationship between the microsporidia and *R. allomycis* is a result of true phylogenetic signal. The present phylogeny provides an alternative hypothesis for the placement of microsporidia, specifically on the earliest diverging fungal branch with the chytrid *R. allomycis*. However, support for this relationship is derived only from the *RPB1* and *RPB2* gene partitions and is not supported by rDNA (see Supplementary Notes 3); alternative hypotheses in which the microsporidia diverge among early fungi cannot be rejected. The ultimate resolution of the placement of microsporidia will require sampling of additional genes from basal fungal taxa.

Dikarya

The majority (~98%) of described fungal species are members of the dikarya clade, which includes the two phyla Ascomycota and Basidiomycota. Ascomycota is the largest phylum within the Fungi and is characterized by the production of meiospores (ascospores) in specialized sac-shaped meiosporangia (asci), which may or may not be produced within a sporocarp (ascoma). Ascomycota is divided into three monophyletic subphyla: Taphrinomycotina, Saccharomycotina and Pezizomycotina (each of which is well supported as monophyletic in the phylogeny; Fig. 1). Taphrinomycotina is resolved as the earliest diverging clade; it includes a diverse group of species that exhibit yeast-like (for example, *Pneumocystis*) and dimorphic—that is, yeast-like and filamentous (for example, *Taphrina*)—growth forms. The subphylum Saccharomycotina consists of the ‘true yeasts’, including bakers’ yeast (*Saccharomyces cerevisiae*) and *Candida albicans*, the most frequently encountered fungal pathogen of humans. Pezizomycotina is the largest subphylum of Ascomycota and includes the vast majority of filamentous, fruit-body-producing species. Data presented here resolved the Orbiliomycetes and Pezizomycetes as the early-diverging lineages of the Pezizomycotina, with the remaining seven classes sampled forming a well-supported crown clade. Reduced ascumal morphologies, whereby asci are contained within fruit bodies that are enclosed partially (Dothideomycetes, Eurotiomycetes and some Sordariomycetes) or completely (Eurotiomycetes, Leotiomycetes and some Sordariomycetes), are restricted to the crown clade of Pezizomycotina.

The Basidiomycota includes about 30,000 species of rusts, smuts, yeasts, and mushroom fungi²⁴. Most are characterized by meiospores (basidiospores) on the exterior of typically club-shaped meiosporangia (basidia). Phylogenetic relationships among the three subphyla of Basidiomycota are uncertain. The subphylum Pucciniomycotina is primarily distinguished by containing the rust fungi (7,000 species), which are primarily pathogens of land plants. Cytological and biochemical data²⁵ are consistent with a sister group relationship between the subphyla Ustilaginomycotina and Agaricomycotina, as shown in Fig. 1. The Ustilaginomycotina includes 1,500 species of true smut fungi and yeasts, most of which cause systemic infections of angiosperm hosts. The Agaricomycotina includes almost two-thirds of known basidiomycetes, including the vast majority of mushroom-forming fungi. Much of the morphological diversity exemplified in mushroom fruiting bodies is the result of radiations of certain lineages within the Agaricomycotina, and recovering their relationships with confidence has proven difficult^{26,27}. Early-diverging lineages in the Agaricomycotina, which are strongly supported in Fig. 1, also include parasitic and/or saprotrophic fungi capable of dimorphism or yeast-like phases. The mycorrhizal basidiomycetes

seem to have multiple, independent evolutionary origins from saprotrophic ancestors as previously suggested²⁸.

Characteristics of early fungi

We reconstructed ancestral states for major nutritional modes in the Fungi using maximum likelihood (Fig. 1). Most of the ancestral character states of deep nodes are equivocal, with the exception of the common ancestor of members of the Basidiomycota, for which a parasitic ancestor is suggested. The phylogeny suggests that numerous transitions from a pathogenic to a saprophytic nutritional mode have occurred, as well as the reverse (Fig. 1). Although the nutritional mode of the common ancestor of Fungi is ambiguous, the earliest diverging branch in the Fungi contains parasitic species (*R. allomycis* and microsporidia). Recent studies^{9,29} showed that the closest known relative to Fungi is the amoeboid protist *Nuclearia*, which grows phagotrophically on algae and bacteria. Amoeboid phases are also observed in basal fungi: *Rozella* seems to phagocytose the organelles of its host³⁰ and many chytrid zoospores undergo an amoeboid, motile phase before encysting. After the divergence of the *Rozella* and microsporidia lineage, the remaining fungi evolved filamentous growth (for example, hyphae and rhizoids), which aids in substrate attachment and absorptive nutrition involving extracellular digestion. Within the Basidiomycota and Ascomycota, a reversion to a unicellular, yeast-like growth form is observed among the earliest diverging lineages, perhaps implicating a prior advantage for this growth form in the early history of the Fungi.

It is unclear whether the common ancestor of Fungi was marine. Most zoosporic true fungi, including all of the chytrids sampled in this study, grow in freshwater or soil habitats. Therefore, the diversification of the major lineages (phyla) within the kingdom Fungi probably occurred in a terrestrial environment but before the emergence of land plants^{31,32}. Mycorrhiza-like symbioses of the phylum Glomeromycota are suggested to have been crucial in the colonization of land by plants³³. Extant members of the Glomeromycota live exclusively as obligate symbionts of photoautotrophs, including not only vascular plants and bryophytes, but also cyanobacteria. This raises the hypothesis that terrestrial members of the Glomeromycota living symbiotically with cyanobacteria or algae, in semi-aquatic and humid habitats later became the symbiotic partners of early land plants³⁴.

The present multilocus phylogeny explains the possible morphology and ecology of early fungi. The early-diverging lineages consist of a grade of zoosporic fungi, suggesting that the earliest fungi were primarily aquatic and lacked aerial spore dispersal. The loss of flagellated spores is inferred to have occurred at least four times. Each loss seems to have coincided with novel innovations in spore production and dispersal: microscopic wind-dispersed spores in terrestrial fungi; forcibly discharged conidia in the Entomophthorales; non-flagellated, mitotically produced spores in the planktonic *Hyaloraphidium curvatum*; and a complex polar tube apparatus in microsporidia. The sister kingdom to the Fungi (Animalia) evolved diverse body plans capable of feeding by ingestion, whereas the fungal branch developed a myriad of unicellular and filamentous forms optimized for absorptive nutrition. With a well-resolved phylogeny, fungal biologists can now study the evolution of complexity and multicellularity, and compare the evolution of these traits in fungi with their evolution in plants and animals.

METHODS

Molecular techniques. Sequence data were generated from 170 fungal species, primarily using pure cultures and herbarium material (Supplementary Notes 1). We used standard polymerase chain reaction (PCR) protocols²⁵ for amplification and sequencing of six gene regions: the 18S ribosomal RNA gene (nearly full length), the 28S ribosomal RNA gene (primers LR0R and LR7), the internal transcribed spacer (ITS) RNA gene region (full length), *EF1 α* (mostly primers EF1-983F and EF1-2218R), RNA polymerase II largest subunit (*RPB1*, mostly primers RPB1-Af and RPB1-G2R) and RNA polymerase II second largest subunit (*RPB2*, primers RPB2-5F and RPB2-11bR). Information on the PCR primers can be found at <http://www.aftol.org/primers.php>. In a number of basal

fungal taxa, the *EF1 α* gene was not detected, but a paralogous copy of the gene was recovered (*EFL*, or the *EF1 α* -like gene³⁵; see Supplementary Notes 4). We also obtained sequences from fungal and eukaryotic genomes by retrieving sequences from GenBank and genome servers. Although our data set contains both partial sequences and missing data points, in the case of only one taxon (the choanoflagellate *Monosiga brevicollis*) were fewer than four genes sampled.

Phylogenetic reconstruction. The data set consisted of 214 taxa, 199 of which were fungi. Sequences were aligned and ambiguous regions excluded in MacClade³⁶. Conflict among the six genes was assessed by separate MLBS of each data partition using 250 bootstrap replicates in PHYML³⁷. We ignored two conflicts, one including microsporidian 18S sequences (known to be subject to long branch attraction) and the other involving marginally conflicting signal of the Pyrenulales (Ascomycota). Data were combined into one matrix with *EF1 α* , *RPB1* and *RPB2* translated into amino acids and 18S, 28S and 5.8S as nucleotides. We applied a heterogeneous maximum likelihood model to the data set with six unlinked partitions, one for each gene. The 18S and 28S genes were fitted to a general-time-reversible model with a proportion of invariant sites and gamma distributed rates (GTR+I+ Γ), the 5.8S data used GTR+ Γ and proteins used the JTT+I+ Γ fixed rate model. The gamma distribution was approximated using four rate classes. We used MrBayes 3.1.1 (ref. 38) for phylogenetic estimation. Five independent runs were conducted (each with four chains) for 9.5×10^6 generations, sampling every 500 generations. Runs were discarded if they failed to reach the same likelihood plateau observed in other independent runs. We computed the consensus of the sampled trees, the posterior probabilities of clades, and average branch lengths from runs that converged to the same likelihood plateau (58,611 trees). For the analysis of the combined super-matrix we also tested for convergence of runs by analysing frequencies of splits using the software AWTY³⁹ and found that the consensus topology constructed using this criterion trivially differed from that based on log likelihood scores. We also assessed support for nodes on the nucleotide data (third codon positions excluded) by MLBS (500 replicates) using PHYML with a GTR+I+ Γ model. Tests for statistical differences in likelihoods of alternative topologies were assessed using the approximately unbiased test²² on the nucleotide data with site-wise, log-likelihood values calculated using TREE-PUZZLE v5.2 (ref. 40).

Ancestral character state reconstruction of nutritional mode was conducted using the maximum likelihood model Mk1 in Mesquite 1.0 (ref. 41). Taxa were assigned to ecological character states on the basis of published literature, resolving ambiguous assignments when possible. Reconstructions are reported for only those branches significantly assigned an unequivocal character state in a majority of 1,000 trees randomly drawn from the sample of credible trees. The number of losses of the flagellum within the Fungi was also estimated for all 58,611 credible trees using Dollo parsimony as implemented in MacClade.

Received 4 May; accepted 25 July 2006.

- Karol, K. G., McCourt, R. M., Cimino, M. T. & Delwiche, C. F. The closest living relatives of land plants. *Science* **294**, 2351–2353 (2001).
- Groth-Malonek, M., Pruchner, D., Grewe, F. & Knoop, V. Ancestors of trans-splicing mitochondrial introns support serial sister group relationships of hornworts and mosses with vascular plants. *Mol. Biol. Evol.* **22**, 117–125 (2005).
- Lang, B. F., O'Kelly, C., Nerad, T., Gray, M. W. & Burger, G. The closest unicellular relatives of animals. *Curr. Biol.* **12**, 1773–1778 (2002).
- James, T. Y., Porter, D., Leander, C. A., Vilgaly, R. & Longcore, J. E. Molecular phylogenetics of the Chytridiomycota supports the utility of ultrastructural data in chytrid systematics. *Can. J. Bot.* **78**, 336–350 (2000).
- Tanabe, Y., Saikawa, M., Watanabe, M. M. & Sugiyama, J. Molecular phylogeny of Zygomycota based on EF-1 α and RPB1 sequences: limitations and utility of alternative markers to rDNA. *Mol. Phylogenet. Evol.* **30**, 438–449 (2004).
- Schüßler, A., Schwarzott, D. & Walker, C. A new fungal phylum, the Glomeromycota: phylogeny and evolution. *Mycol. Res.* **105**, 1413–1421 (2001).
- Tehler, A., Little, D. P. & Farris, J. S. The full-length phylogenetic tree from 1551 ribosomal sequences of chitinous fungi. *Fungi. Mycol. Res.* **107**, 901–916 (2003).
- Cavalier-Smith, T. & Chao, E. E. The opalozoan *Apusomonas* is related to the common ancestor of animals, fungi, and choanoflagellates. *Proc. R. Soc. Lond. B* **261**, 1–9 (1995).
- Steenkamp, E. T., Wright, J. & Baldauf, S. L. The protistan origins of animals and fungi. *Mol. Biol. Evol.* **23**, 93–106 (2005).
- Barr, D. J. S. Evolution and kingdoms of organisms from the perspective of a mycologist. *Mycologia* **84**, 1–11 (1992).
- Ustinova, I., Krienitz, L. & Huss, V. A. R. *Hyaloraphidium curvatum* is not a green alga, but a lower fungus; *Amoebidium parasiticum* is not a fungus, but a member of the DRIPs. *Protist* **151**, 253–262 (2000).
- Nagahama, T., Sato, H., Shimazu, M. & Sugiyama, J. Phylogenetic divergence of the entomophthorean fungi: evidence from nuclear 18S ribosomal RNA gene sequences. *Mycologia* **87**, 203–209 (1995).
- McKerracher, L. J. & Heath, I. B. The structure and cycle of the nucleus-associated organelle in two species of *Basidiobolus*. *Mycologia* **77**, 412–417 (1985).

- Blackwell, M. & Malloch, D. Similarity of *Amphoromorpha* and secondary capilliconidia of *Basidiobolus*. *Mycologia* **81**, 735–741 (1989).
- Keeling, P. J. & Fast, N. M. In *Insect–Fungal Associations: Ecology and Evolution* (eds Vega, F. E. & Blackwell, M.) 97–118 (Oxford Univ. Press, Oxford, 2005).
- Hirt, R. P. et al. Microsporidia are related to fungi: evidence from the largest subunit of RNA polymerase II and other proteins. *Proc. Natl Acad. Sci. USA* **96**, 580–585 (1999).
- Katinka, M. D. et al. Genome sequence and gene compaction of the eukaryote parasite *Encephalitozoon cuniculi*. *Nature* **414**, 450–453 (2001).
- Keeling, P. J. Congruent evidence from α -tubulin and β -tubulin gene phylogenies for a zygomycete origin of microsporidia. *Fungal Genet. Biol.* **38**, 298–309 (2003).
- Cavalier-Smith, T. In *The Mycota* (eds McLaughlin, D. J., McLaughlin, E. G. & Lemke, P. A.) 3–37 (Springer, New York, 2001).
- Corradi, N., Hijri, M., Fumagalli, L. & Sanders, I. R. Arbuscular mycorrhizal fungi (Glomeromycota) harbour ancient fungal tubulin genes that resemble those of the chytrids (Chytridiomycota). *Fungal Genet. Biol.* **41**, 1037–1045 (2004).
- Held, A. A. *Rozella* and *Rozellopsis*: Naked endoparasitic fungi which dress up as their hosts. *Bot. Rev.* **47**, 451–515 (1981).
- Shimodaira, H. An approximately unbiased test of phylogenetic tree selection. *Syst. Biol.* **51**, 492–508 (2002).
- Gill, E. E. & Fast, N. M. Assessing the microsporidia–fungi relationship: combined phylogenetic analysis of eight genes. *Gene* **375**, 103–109 (2006).
- Kirk, P. M., Cannon, P. F., David, J. C. & Stalpers, J. A. (eds) *Ainsworth & Bisby's; Dictionary of the Fungi* 60 (CAB International, Wallingford, UK, 2001).
- Lutzoni, F. et al. Assembling the fungal tree of life: progress, classification, and evolution of subcellular traits. *Am. J. Bot.* **91**, 1446–1480 (2004).
- Binder, M. et al. The phylogenetic distribution of resupinate forms across the major clades of mushroom-forming fungi (Homobasidiomycetes). *Systematics and Biodiversity* **3**, 113–157 (2005).
- Moncalvo, J.-M. et al. One hundred and seventeen clades of euagarics. *Mol. Phylogenet. Evol.* **23**, 357–400 (2002).
- Hibbett, D. S., Gilbert, L.-B. & Donoghue, M. J. Evolutionary instability of ectomycorrhizal symbioses in basidiomycetes. *Nature* **407**, 506–508 (2000).
- Medina, M. et al. Phylogeny of Opisthokonta and the evolution of multicellularity and complexity in Fungi and Metazoa. *Int. J. Astrobiol.* **2**, 203–211 (2003).
- Powell, M. J. Fine structure of the unwallid thallus of *Rozella polyphagi* in its host *Polyphagus euglenae*. *Mycologia* **76**, 1039–1048 (1984).
- Berbee, M. L. & Taylor, J. W. In *The Mycota* (eds McLaughlin, D. J., McLaughlin, E. G. & Lemke, P. A.) 229–245 (Springer, New York, 2001).
- Heckman, D. S. et al. Molecular evidence for the early colonization of land by fungi and plants. *Science* **293**, 1129–1133 (2001).
- Pirozynski, K. A. & Malloch, D. W. The origin of land plants: a matter of mycotrophism. *Biosystems* **5**, 153–164 (1975).
- Schüßler, A. Molecular phylogeny, taxonomy, and evolution of arbuscular mycorrhiza fungi and *Geosiphon pyriformis*. *Plant Soil* **244**, 75–83 (2002).
- Keeling, P. J. & Inagaki, Y. A class of eukaryotic GTPase with a punctuate distribution suggesting multiple functional replacements of translation elongation factor 1 α . *Proc. Natl Acad. Sci. USA* **101**, 15380–15384 (2004).
- Maddison, D. & Maddison, W. *MacClade Version 4.05: Analysis of Phylogeny and Character Evolution* (Sinauer Associates, Sunderland, Massachusetts, USA, 2002).
- Guindon, S. & Gascuel, O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* **52**, 696–704 (2003).
- Huelsenbeck, J. P. & Ronquist, F. MrBayes: bayesian inference of phylogenetic trees. *Bioinformatics* **17**, 754–755 (2001).
- Wilgenbusch, J. C., Warren, D. L. & Swofford, D. L. AWTY: A system for graphical exploration of MCMC convergence in Bayesian phylogenetic inference. <http://ceb.csit.fsu.edu/awty> (2004).
- Schmidt, H. A., Strimmer, K., Vingron, M. & von Haeseler, A. TREE-PUZZLE: maximum likelihood phylogenetic analysis using quartets and parallel computing. *Bioinformatics* **18**, 502–504 (2002).
- Maddison, W. P. & Maddison, D. R. Mesquite: a modular system for evolutionary analysis. Version 1.0. <http://mesquiteproject.org> (2003).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements Funding for the project was provided by the National Science Foundation's 'Assembling the Tree of Life' and 'Research Coordination Network' programs. For technical assistance we thank L. Bukovnik, C. Roberts and H. Matthews. We also thank the following individuals for sharing research materials: M. C. Aime, W. R. Buck, M. S. Cole, P. Crane, Y. Dalpe, D. M. Hillis, S. L. Joneson, R. Petersen, C. Printzen, E. Vellinga, H. Whisler and A. Zavarzin. We are very thankful to B. Mueller, J. Harer, B. Rankin, J. Pormann and S. Dilda for providing access to the Duke CSEM computer cluster. P. Keeling provided unpublished information used to analyse EFL in Fungi.

Author Information Data for this project have been deposited in GenBank (see Supplementary Notes 1 for accession numbers), and the alignments can be accessed on the Assembling the Fungal Tree of Life website at <http://www.aftol.org/>. Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to T.Y.J. (tyj2@duke.edu) or R.V. (fungi@duke.edu).