



Prewaning mortality in SWAP farrowing pens

Pedersen, Janni Hales; Moustsen, Vivi; Nielsen, Mai Britt Friis; Hansen, Christian Fink

Publication date:
2014

Document version
Early version, also known as pre-print

Citation for published version (APA):
Pedersen, J. H., Moustsen, V., Nielsen, M. B. F., & Hansen, C. F. (2014). *Prewaning mortality in SWAP farrowing pens*. Abstract from The 65th Annual Meeting of the European Federation of Animal Science, Copenhagen, Denmark.
http://www.eaap.org/Previous_Annual_Meetings/2014Copenhagen/Copenhagen_2014_Abstracts.pdf

Preweaning mortality in SWAP farrowing pensJ. Hales¹, V.A. Moustsen², M.B.F. Nielsen² and C.F. Hansen¹¹University of Copenhagen, Department of Large Animal Sciences, Groenegaardsvej 2, 1870 Frederiksberg, Denmark, ²Pig Research Centre, Axeltorv 3, 1609 Copenhagen, Denmark; cfh@sund.ku.dk

Confining sows in farrowing crates increases stress and affects welfare negatively. However, piglets may be at greater risk of dying in loose systems, especially in early lactation. The objective of this study was to investigate if short term confinement around farrowing would decrease piglet mortality and increase number of weaned piglets in a farrowing pen where sows could be confined temporarily (SWAP=Sow Welfare And Piglet protection). The study was conducted in a 1,200 sow piggery where sows were randomly allocated to one of three experimental treatments: loose-loose (LL, n=376), loose-confined (LC, n=369) or confined-confined (CC, n=354). In LL sows were loose from placement in the farrowing unit to weaning, LC sows were loose to birth of the last piglet and thereafter confined to day 4 post farrowing, and CC sows were confined from day 114 of gestation to day 4. All sows were loose housed from day 4 to weaning. Live born and stillborn piglets, equalised litter size and dead piglets were recorded. All data were analysed using linear models in SAS. As expected, the number of total born piglets per litter (17.4 ± 0.10 ; $P=0.55$) did not differ between treatments. However, the number of dead piglets per litter before equalisation was greater in LL (2.0 piglets/litter (1.8-2.1)) and LC (1.8 piglets/litter (1.7-1.9)) than in CC (1.3 piglets/litter (1.2-1.4); $P<0.001$). Piglet mortality from litter equalisation to day 4 was higher in LL (7.6% (7.1-8.7)) than in CC (6.0% (5.3-6.8); $P<0.001$) but did not differ between LC (7.0% (6.2-7.8)) and the other two treatments. From day 4 to weaning the piglet mortality rate was 5.8% in all treatments ($P=0.98$). Consequently there were more weaned piglets in CC (13.7 ± 0.19) than in LL (12.9 ± 0.19 ; $P=0.005$) but there was no difference between LC (13.1 ± 0.19) and LL or CC ($P>0.05$). In conclusion, confinement of sows around farrowing reduced piglet mortality compared to loose housed sows.

Session 43


Theatre 7

Saliva cortisol levels of nurse sows and ordinary sows through lactationC. Amdi¹, V.A. Moustsen², G. Sørensen², L.C. Oxholm² and C.F. Hansen¹¹University of Copenhagen, Grønnegårdsvej 2, Frb C, Denmark, ²Pig Research Centre, Axeltorv 3, Kbh V, Denmark; ca@sund.ku.dk

Nurse sows are used in piggeries with hyper-prolific sows to manage large litters. It is, however, not known if nurse sows experience prolonged stress by having to stay in farrowing crates beyond the normal weaning time. The aim of this study was to quantify the long-term saliva cortisol response as a measurement of the nurse sows stress level compared to ordinary sows (OSOW) weaning their piglets at d 25. In Denmark cascade fostering using two lactating sows are normally performed. The first nurse sow (NURSE1) has her own piglets removed after a week and receives surplus newborn piglets that she fosters until weaning. The second nurse sow (NURSE2) weans her own litter after 21 days and receives the litter from NURSE1 which she rears until weaning. In total 60 sows (n=20) were randomly allocated to become an OSOW, NURSE1 or NURSE2. Saliva cortisol was collected on d 6, 13, 20 and 24 postpartum at 10 h, 13 h and 16 h for all sows and pooled on a daily basis for analyses. Additional samples were taken on d 31 for NURSE1 and NURSE2 and d 38 for NURSE2. Saliva samples were assayed for cortisol levels using a Salivary Cortisol EIA kit (Salimetrics, UK). Cortisol data were log transformed and analysed using proc mixed in SAS. Results showed that there was no effect of group but there was an effect of day ($P<0.001$) with saliva cortisol declining throughout lactation with values of 19.9, 17.1, 12.8 and 10.4 nmol/l (back transformed values) at d 6, 13, 20 at d 24, respectively. NURSE1 tended to have lower values (8.3 nmol/l) on d 31 than on d 24 (11.5 nmol/l; $P=0.08$). NURSE2 had lower cortisol values on d 38 (7.4 nmol/l) and on d 31 (7.5 nmol/l) than on d 24 (11.1 nmol/l; $P<0.05$). Results indicate that saliva cortisol levels decline throughout lactation and that there is no difference in saliva cortisol levels between OSOWS and nurse sows. In conclusion, saliva cortisol levels indicate no additional long-term stress of being selected as a nurse sow.

Arviden

Book of Abstracts of the 65th Annual Meeting of the European Federation of Animal Science



Book of abstracts No. 20 (2014)
Copenhagen, Denmark
25 - 29 August 2014