



Draft whole-genome sequence of *Bacillus sonorensis* strain L12, a source of nonribosomal lipopeptides

Adimpong, David Bichala; Sørensen, Kim I.; Nielsen, Dennis Sandris; Thorsen, Line; Rasmussen, Thomas B.; Derkx, Patrick M.F.; Jespersen, Lene

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.00097-13](https://doi.org/10.1128/genomeA.00097-13)

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Adimpong, D. B., Sørensen, K. I., Nielsen, D. S., Thorsen, L., Rasmussen, T. B., Derkx, P. M. F., & Jespersen, L. (2013). Draft whole-genome sequence of *Bacillus sonorensis* strain L12, a source of nonribosomal lipopeptides. *Genome Announcements*, 1(2). <https://doi.org/10.1128/genomeA.00097-13>

Draft Whole-Genome Sequence of *Bacillus sonorensis* Strain L12, a Source of Nonribosomal Lipopeptides

David B. Adimpong,^{a,b} Kim I. Sørensen,^b Dennis S. Nielsen,^a Line Thorsen,^a Thomas B. Rasmussen,^c Patrick M. F. Derkx,^b Lene Jespersen^a

University of Copenhagen, Faculty of Science, Department of Food Science, Frederiksberg C, Denmark^a; Chr. Hansen A/S, Discovery Department, Hørsholm, Denmark^b; Novo Nordisk, A/S, Department of Molecular Genetics, Måløv, Denmark^c

The *Bacillus sonorensis* L12 draft genome sequence is approximately 4,647,754 bp in size with a G+C content of 45.2%. Over 86% of the genome contains protein-encoding genes, including several gene clusters for *de novo* biosynthesis of the nonribosomal lipopeptides iturin, bacitracin, and fengycin, which could mean that the strain exhibits antifungal effects.

Received 13 February 2013 Accepted 28 February 2013 Published 28 March 2013

Citation Adimpong DB, Sørensen KI, Nielsen DS, Thorsen L, Rasmussen TB, Derkx PMF, Jespersen L. 2013. Draft whole-genome sequence of *Bacillus sonorensis* strain L12, a source of nonribosomal lipopeptides. *Genome Announc.* 1(2):e00097-13. doi:10.1128/genomeA.00097-13.

Copyright © 2013 Adimpong et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to David B. Adimpong, dadimpong@life.ku.dk.

Bacillus sonorensis is a Gram-positive aerobic-endospore-forming bacterium and a member of the *Bacillus subtilis* group of microorganisms. It was first isolated from the Sonoran Desert soil and is phenotypically and genotypically closely related to *Bacillus licheniformis* by sharing traits such as being facultatively anaerobic (1–3). However, the *B. sonorensis* and *B. licheniformis* species can be phenotypically distinguished based on colony pigmentation (1, 4) and sensitivity to different level of clindamycin (5). The *B. licheniformis* sp. is exploited industrially for large-scale production of many enzymes, antibiotics, and biochemicals (3) and works as a host for cloning of several genes encoding α -amylases, etc. Several members of the *B. subtilis* group are also used as probiotics and biocontrol agents in crop farming (6, 7). Thus, given that the industrially relevant organism *B. licheniformis* is closely related to *B. sonorensis*, about which little is known, we sequenced the genome of *B. sonorensis* L12, which was isolated from Gergoush primary starter materials in Sudan (4). It is expected that the information obtained can be used to explore the biotechnological relevance of this organism as well as its genomics and phylogenetic status related to members of the *B. subtilis* group of microorganisms and thereby enhance our limited knowledge on the *B. sonorensis* species.

The genomic DNA of strain L12 was isolated from overnight culture using a commercial DNA isolation kit (GenElute bacterial genomic DNA kit, NA2110; Sigma-Aldrich Co., St. Louis, MO). Sequencing was performed at the Beijing Genomics Institute (BGI, Shenzhen, China) using a combination of randomly sheared libraries with inserts of 0.5 to 2 kb in size. The genomic DNA was sequenced using the Illumina HiSeq platform with a total coverage of 70 \times . The reads were assembled into contigs using CLC Genomics Workbench v. 5 (CLC Bio, Aarhus, Denmark), which was also used to determine the nucleotide sequence statistics. Genome annotation and preparation for submission to GenBank were performed using the NCBI Prokaryotic Genomes Automatic Annotation Pipeline (PGAAP). Putative tRNAs were predicted using the tRNAscan-SE 1.21 Server (8) programs.

The assembled draft genome sequence of strain L12 contains 34 contigs, a genome size of 4,647,754 bp, and a G+C content of 45.2%. The genome was 425,108 bp larger than the *B. licheniformis* DSM13^T genome. Strain L12 has 78 tRNA genes and 4,236 protein-coding genes representing 86.8% of the draft genome sequence.

Preliminary genome analysis revealed that strain L12 carries gene clusters for *de novo* biosyntheses of the nonribosomal lipopeptides fengycin, iturin, and bacitracin, which have potential biotechnological applications (9, 10). It therefore represents a potential strain which can be used as a cloning tool for genetic engineering and production of improved and novel antimicrobial agents and also as a bioprotective agent for controlling plant fungal pathogens in crop farming. Further comparative genomic analyses will also provide valuable insight into the evolutionary and phylogenetic status of this species and contribute to a deeper understanding of its ecology and evolution.

Nucleotide sequence accession numbers. This whole genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [AOFM000000000](https://www.ncbi.nlm.nih.gov/nuccore/AOFM000000000). The version described in this paper is the first version, [AOFM010000000](https://www.ncbi.nlm.nih.gov/nuccore/AOFM010000000).

ACKNOWLEDGMENTS

Financial support for this project was kindly provided by Chr Hansen A/S, Denmark, and the Danish Ministry of Foreign Affairs through Danida. We express our sincere gratitude to them for this assistance.

REFERENCES

1. Palmisano MM, Nkamura LK, Duncan KE, Istock CA, Cohan FM. 2001. *Bacillus sonorensis* sp. nov., a close relative of *Bacillus licheniformis*, isolated from soil in the Sonoran Desert, Arizona. *Int. J. Syst. Evol. Microbiol.* 51:1671–1679.
2. Fritze D. 2004. Taxonomy of the genus *Bacillus* and related genera: the aerobic endospore-forming bacteria. *Phytopathology* 94:1245–1248.
3. Rey MW, Ramaiya P, Nelson BA, Brody-Karpin SD, Zaretsky EJ, Tang M, Lopez de Leon A, Xiang H, Gusti V, Clausen IG, Olsen PB, Rasmussen MD, Andersen JT, Jørgensen PL, Larsen TS, Sorokin A, Bolotin A, Lapidus A, Galleron N, Ehrlich SD, Berka RM. 2001.

- Complete genome sequence of the industrial bacterium *Bacillus licheniformis* and comparisons with closely related *Bacillus* species. *Genome Biol.* 5:R77.
4. Thorsen L, Abdelgadir WS, Rønsbo MH, Abban S, Hamad SH, Nielsen DS, Jakobsen M. 2011. Identification and safety evaluation of *Bacillus* species occurring in high numbers during spontaneous fermentations to produce Gergoush, a traditional Sudanese bread snack. *Int. J. Food Microbiol.* 146:244–252.
 5. Adimpong DB, Sørensen KI, Thorsen L, Stuer-Lauridsen B, Abdelgadir WS, Nielsen DS, Derkx PMF, Jespersen L. 2012. Antimicrobial susceptibility of *Bacillus* strains isolated from primary starters for African traditional bread production and characterization of the bacitracin operon and bacitracin biosynthesis. *Appl. Environ. Microbiol.* 78:7903–7914.
 6. Brannen PM, Kenney DS. 1997. Kodiak®—a successful biological-control product for suppression of soil-borne plant pathogens of cotton. *J. Ind. Microbiol. Biotechnol.* 19:169–171.
 7. Cutting SM. 2011. *Bacillus* probiotics. *Food Microbiol.* 28:214–220.
 8. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25:955–964.
 9. Fickers P. 2012. Antibiotic compounds from *Bacillus*: why are they so amazing? *Am. J. Biochem. Biotechnol.* 8:40–46.
 10. Hoffmann A, Pag U, Wiedemann I, Sahl HG. 2002. Combination of antibiotic mechanisms in lantibiotics. *Farmacol.* 57:685–691.