



## **Gut microbiota composition may relate to weight loss rate in obese pet dogs**

Kieler, Ida Nordang; Vitger, Anne Désiré; Bin Shamzir Kamal, Shamrulazhar; Nielsen, Dennis Sandris; Bjørnvad, Charlotte Reinhard

*Published in:*  
European Society of Veterinary and Comparative Nutrition

*Publication date:*  
2015

*Document version*  
Other version

*Citation for published version (APA):*  
Kieler, I. N., Vitger, A. D., Bin Shamzir Kamal, S., Nielsen, D. S., & Bjørnvad, C. R. (2015). Gut microbiota composition may relate to weight loss rate in obese pet dogs. In *European Society of Veterinary and Comparative Nutrition*

## Gut microbiota composition may relate to weight loss rate in obese pet dogs

Kieler I. N.<sup>1\*</sup>, Vitger A.<sup>1</sup>, Shamzir Kamal S.<sup>2</sup>, Nielsen D. S.<sup>2</sup> Bjornvad C. R.<sup>1</sup>

<sup>1</sup>Dyrlaagevej 16, DK-1870 Frb. C, Denmark, <sup>2</sup> University of Copenhagen, Department of Food Science, Rolighedsvej 26, DK-1958 Frb. C, Denmark  
\*ink@sund.ku.dk

**Introduction:** The gut microbiota (GM) composition of obese individuals differs from the GM of lean individuals. Furthermore, the efficiency of a high-protein diet to induce weight loss in overweight humans depends on the GM (Santacruz et al. 2009). We aimed to study the interaction between exercise, weight-loss rate and the GM during weight-loss in dogs.

**Animals, Material and Methods:** Fifteen privately-owned obese dogs were recruited for a 12-week weight-loss intervention, 8 were exercised and fed restrictively with a high-protein diet, while 7 were only fed restrictively and no changes to their exercise regime was made. Fecal samples were collected and the dogs were weighed at baseline, after 6 and 12 weeks (Vitger et al. 2015). GM was determined using MiSeq-based tag-encoded 16S rRNA gene high-throughput amplicon sequencing, and analyzed by ANOVA, ANOSIM and linear regression analyzes (QIIME 1.7.0).

**Results and Discussion:** Total weight loss, food allowance and GM were not significantly different between groups. Both groups were dominated by genera *Fusobacterium* (~20-50%) and *Bacteroides* (~15-35%). A significant increase in the genera *Suterella*, *Fusobacterium* and a decrease in the *Megamonas* were observed. *Megamonas* abundance negatively correlated to high weight loss (>1% per week). *Ruminococcaceae* were significantly higher at 12 weeks in dogs with a low weight loss (<0.5% per week). Members of *Megamonas* and *Ruminococcaceae* produce acetate, a short chain fatty acid (SCFA), and a substrate for lipogenesis and cholesterol formation (Conterno et al. 2011). Obese rodents also have a GM with a putative increased SCFA producing ability (Turnbaugh et al. 2006), and the GM of some obese individuals may therefore produce more SCFA, which may affect weight loss rate.

**Conclusion:** Following a 12 week weight-loss intervention in dogs a significant GM change, which may relate to weight loss rate, was seen.

### References:

- Conterno, L.; Fava, F.; Viola, R.; Tuohy, K.M., 2011: Obesity and the gut microbiota: does up-regulating colonic fermentation protect against obesity and metabolic disease? *Genes & nutrition* **6**, 241-260.
- Santacruz, A.; Marcos, A.; Warnberg, J.; Marti, A.; Martin-Matillas, M.; Campoy, C.; Moreno, L.A.; Veiga, O.; Redondo-Figuero, C.; Garagorri, J.M.; Azcona, C.; Delgado, M.; Garcia-Fuentes, M.; Collado, M.C.; Sanz, Y.; Group, E.S., 2009: Interplay between weight loss and gut microbiota composition in overweight adolescents. *Obesity* **17**, 1906-1915.
- Turnbaugh, P.J.; Ley, R.E.; Mahowald, M.A.; Magrini, V.; Mardis, E.R.; Gordon, J.I., 2006: An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* **444**, 1027-1031.
- Vitger, A.D.; Stallknecht, B.M.; Nielsen, D.H.; Bjornvad, C.R., 2015: Integration of a physical training program in a weight loss plan for overweight pet dogs. *Journal of American Veterinary Medical Association*, Accepted.