



Non-attending patients in general practice

Ellis, David A; McQueenie, Ross; McConnachie, Alex; Wilson, Philip; Williamson, Andrea

Published in:
The Lancet Public Health

DOI:
[10.1016/S2468-2667\(18\)30020-3](https://doi.org/10.1016/S2468-2667(18)30020-3)

Publication date:
2018

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Ellis, D. A., McQueenie, R., McConnachie, A., Wilson, P., & Williamson, A. (2018). Non-attending patients in general practice. *The Lancet Public Health*, 3(3), [e113]. [https://doi.org/10.1016/S2468-2667\(18\)30020-3](https://doi.org/10.1016/S2468-2667(18)30020-3)

Non-attending patients in general practice

We agree with Frans Smits and Gerben ter Riet when, in their comment,¹ they suggest that it could be valuable to frame questions relating to non-attendance in terms of societal benefits and harms. However, we would like to take this opportunity to provide some additional clarification with regards to our analysis and presentation of results.²

Smits and ter Riet argue that we should have included information on medical diagnoses in our study² rather than provide this analysis in a future publication. Our decision to incorporate this information into a future publication was not taken lightly; the issues were raised during the review process, with one reviewer suggesting that the burden of long-term conditions is likely to be an important factor in the unmet need and behaviours of the patients within this population.³ However, we judged that presenting these data satisfactorily in a single paper would have been overly complex, with other reviewers supporting this decision. In addition to a paper that will focus specifically on patients with multiple long-term conditions, we also plan to publish a future overarching paper focused on unmet need along with health-care utilisation across the health system.

Smits and ter Riet also suggested that our analysis might benefit from a multilevel approach that would involve the use of zero-inflated negative binomial models. Such an approach might be particularly useful, given that 54% of patients did not miss any appointments. We considered that the negative binomial models fitted the data reasonably well. An initial analysis attempted to use a mixed effects regression allowing for random practice effects, but even the simplest of models proved intractable in a dataset of this size, which was

held with limited computational capacity (Safe Haven). To counter this limitation, we adjusted the analysis for available practice-level variables.

In our article,² we focused on describing the data, and the main effects of several patient and practice-level factors. To examine cross-factor interactions would have added another layer of complexity, which would have been very difficult to condense into a single paper. Such analyses would probably best be focused on interactions between a single factor (eg, sex) and factors that predict frequent non-attendance to address a coherent research question.

Finally, we agree that frequent attenders are indeed an interesting subgroup within themselves, which was why all our models were offset for the number of appointments made.² However, the claim that frequent attendance would prompt more changes to clinical work than non-attendance is unfounded. Although frequent attenders and frequent non-attenders are qualitatively different in terms of social and socioeconomic problems, they provide two equally important examples of unmet need.^{2,4,5}

We declare no competing interests.

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

*David A Ellis, Ross McQueenie,
Alex McConnachie, Philip Wilson,
Andrea Williamson
d.a.ellis@lancaster.ac.uk

Department of Psychology, Lancaster University, Lancaster, UK (DAE); General Practice and Primary Care, School of Medicine, Dentistry and Nursing, MVLS, University of Glasgow, Glasgow, UK (RM); Robertson Centre for Biostatistics, Institute of Health and Wellbeing, MVLS, University of Glasgow, Glasgow, UK (AM); Centre for Rural Health, Institute of Applied Health Sciences, University of Aberdeen, Aberdeen, UK (AW).

- 1 Smits FT, ter Riet G. Non-attending patients in general practice. *Lancet Public Health* 2017; 2: e538.
- 2 Ellis DA, McQueenie R, McConnachie A, Wilson P, Williamson AE. Demographic and practice factors predicting repeated non-attendance in primary care: a national retrospective cohort analysis. *Lancet Public Health* 2017; 2: e551–59.

- 3 Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012; 380: 37–43.
- 4 Smits FT, Brouwer HJ, ter Riet G, van Weert HC. Epidemiology of frequent attenders: a 3-year historic cohort study comparing attendance, morbidity and prescriptions of one-year and persistent frequent attenders. *BMC Public Health* 2009; 9: 36.
- 5 Williamson AE, Ellis DA, Wilson P, McQueenie R, McConnachie A. Understanding repeated non-attendance in health services: a pilot analysis of administrative data and full study protocol for a national retrospective cohort. *BMJ Open* 2017; 7: e014120.

