



Acute response of biochemical bone turnover markers and the associated ground reaction forces to high-impact exercise in postmenopausal women

Prawiradilaga, Rizky (Eqi) Suganda; Madsen, Anders Odderup; Jørgensen, Niklas Rye; Helge, Eva Wulff

Published in:
Biology of Sport

DOI:
[10.5114/biolsport.2020.91497](https://doi.org/10.5114/biolsport.2020.91497)

Publication date:
2020

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY-NC-SA](https://creativecommons.org/licenses/by-nc-sa/4.0/)

Citation for published version (APA):
Prawiradilaga, R. E. S., Madsen, A. O., Jørgensen, N. R., & Helge, E. W. (2020). Acute response of biochemical bone turnover markers and the associated ground reaction forces to high-impact exercise in postmenopausal women. *Biology of Sport*, 37(1), 41-48. <https://doi.org/10.5114/biolsport.2020.91497>

Acute response of biochemical bone turnover markers and the associated ground reaction forces to high-impact exercise in postmenopausal women

AUTHORS: Rizky S. Prawiradilaga^{1,2}, Anders O. Madsen¹, Niklas R. Jørgensen^{3,4}, Eva W. Helge¹

¹ Department of Nutrition, Exercise and Sports, University of Copenhagen, Denmark
Nørre Allé 51, 2200 Copenhagen N, Denmark

² Faculty of Medicine, Universitas Islam Bandung, Indonesia
Tamansari No.20, Bandung 40116, Jawa Barat, Indonesia

³ Department of Clinical Biochemistry, Rigshospitalet Glostrup, Denmark
Valdemar Hansens Vej 13, 2600 Glostrup, Denmark

⁴ OPEN, Odense Patient data Explorative Network, Odense University Hospital/Institute of Clinical Research, University of Southern Denmark, Odense, Denmark
J.B. Winsløvs Vej 4, 5000 Odense C, Denmark

ABSTRACT: The aim of the study was to examine the acute response of biochemical bone turnover markers (BTM) to high-impact jumping exercise, and to quantify the ground reaction forces (GRF) achieved during each jumping exercise, in postmenopausal women. In a randomized controlled cross-over study over three days, 29 postmenopausal women (age (mean±SD): 60.0±5.6 years) were randomly assigned to 6x10 repetitions of three different jumps: countermovement jump (CMJ), drop jump (DJ), diagonal drop jump (DDJ). A fourth day without jumping served as a control (CON). Blood samples were collected before (PRE), after (POST), and 2 hours after (2Hr) exercise. Bone turnover was evaluated by bone formation markers (procollagen type-1 amino-terminal propeptide (P1NP) and osteocalcin (OC)) and the bone resorption marker C-terminal telopeptide of type-1 collagen (CTX). Peak anteroposterior (Fx), mediolateral (Fy), and vertical (Fz) GRF were measured using a force platform. From PRE to POST, P1NP increased ($p < 0.01$) by $7.7 \pm 1.8\%$, $9.4 \pm 1.3\%$, and $10.6 \pm 1.6\%$ for CMJ, DJ, and DDJ, which were higher ($p < 0.01$) than CON. OC increased ($p < 0.05$) by $5.5 \pm 1.8\%$ for DJ, which was higher ($p < 0.05$) than CON. CTX was not significantly changed at POST. There were no significant differences in BTM Δ -values between the jumps at any time point. For the CMJ, the combined three-axis peak GRF was positively associated with the PRE to POST Δ -change in P1NP ($r = 0.71$, $p < 0.05$). The acute, jumping-induced increase in P1NP and OC without any rise in CTX may indicate increased bone formation. Moreover, the study shows a dose-response relationship between GRF and the acute P1NP response after countermovement jumps.

CITATION: Prawiradilaga RS, Madsen AO, Jørgensen NR, Helge EW. Acute response of biochemical bone turnover markers and the associated ground reaction forces to high-impact exercise in postmenopausal women. *Biol Sport*. 2020;37(1):41–48.

Received: 2019-10-03; Reviewed: 2019-11-04; Re-submitted: 2019-11-22; Accepted: 2019-11-25; Published: 2020-01-24

Corresponding author:

Rizky Suganda Prawiradilaga
Department of Nutrition
Exercise and Sports
University of Copenhagen
Nørre Allé 51, 2200
Copenhagen N, Denmark
E-mail: rsp@nexs.ku.dk
rizkysuganda@unisba.ac.id

Key words:

Bone formation
Bone resorption
Jumping
Osteogenic exercise
Weight-bearing exercise
Odd-impact

INTRODUCTION

Osteoporosis is a chronic bone disease of increasing public health concern. The disease is characterized by low bone strength due to reduced bone mass and impairment of bone micro-architecture, putting the patient at increased risk of bone fractures [1]. It is estimated to affect 200 million women worldwide, triggering more than 8.9 million fractures annually [2].

Thus, depending on the women's life stage, osteogenic exercise has different effects and aims [3]. For adolescents, the exercise is aimed at increasing peak bone mass, for premenopausal women it is aimed at increasing bone mineral density (BMD), and for postmenopausal women it is aimed at reducing the age-related bone loss [3,4]. In postmenopausal women, a combination of resistance

and high-impact or odd-impact training is found to be effective in improving bone health [3,5].

The osteogenic effect of training is mostly estimated by dual-energy x-ray absorptiometry (DXA) [6]. Additionally, the International Osteoporosis Foundation (IOF) has recommended the use of biochemical bone turnover markers (BTM) as markers of fracture risk assessment and evaluation of treatment effectiveness in clinical settings [7], and the assessment of BTM [8] is a promising method to evaluate an osteogenic response in bone turnover acutely or after only a few weeks of training.

A number of studies have been conducted examining the acute effects of exercise on BTM in adults [8–21], but the results have

been equivocal, and data in postmenopausal women are lacking. Therefore, the present randomized, controlled cross-over study compared the acute biological response of BTM to countermovement jump (CMJ), drop jump (DJ), and diagonal drop jump (DDJ) and the associated ground reaction force (GRF) in postmenopausal women. The overall aim was to investigate whether the jumps differed in the acute osteogenic response, which should be taken into consideration when planning osteogenic training to improve bone mass.

MATERIALS AND METHODS

Experimental approach

We used a randomized, controlled cross-over design comparing the BTM response after three different high-impact jumping trials and a resting control trial (CON) performed in random order in the early morning over four test days at a similar time. Every trial was separated by at least 48 hours without any moderate-vigorous activities. GRF associated with jumping was measured on a separate test day.

Ethics

All participants were fully informed of the procedures and possible discomfort associated with the study before providing their written informed consent to participate. The study was conducted following the Declaration of Helsinki and approved by the local ethics committee of the Capital Region of Denmark, H-4-2012-181.

Subjects

Healthy, sedentary postmenopausal women aged 50–70 years, who were more than two years after menopause, non-smokers, and had a body mass index (BMI) <30 kg/m², were invited to participate in the present study. To determine eligibility, prospective subjects un-

Table 1. Participant characteristics at baseline (n=29)

	Mean ± SD
Age (years)	60.0 ± 5.6
Height (cm)	165.2 ± 5.4
Weight (kg)	65.8 ± 7.7
BMI (kg/m ²)	24.1 ± 2.6
Whole-body BMD (g/cm ²)	1.099 ± 0.069
Lumbar spine BMD (g/cm ²)	1.075 ± 0.099
Total hip BMD (g/cm ²)	0.923 ± 0.086
Percent body fat (%)	34.1 ± 7.4
Total fat mass (kg)	22.3 ± 6.6
Total lean body mass (kg)	41.5 ± 3.7
VO ₂ max (ml/min/kg)	31.8 ± 5.3

BMI = body mass index, BMD = bone mineral density

derwent bone mineral density (BMD) screening at the lumbar spine and proximal femur (total hip) using DXA. If osteoporosis (T-score <-2.5 SD) or high BMD in relation to age (Z-score >1.0 SD) at either the lumbar spine or the total hip was found, the participant was excluded. In addition, the following exclusion criteria were applied: the use of medication, hormone therapy or supplements that affect bone metabolism; previous or current medical condition affecting bone health; conditions that make powerful jumping impossible; and/or engagement in regular high-impact and/or resistance training.

Thirty-five women were recruited via a local newspaper and online advertisement (Fig. 1). After the preliminary BMD screening, three




Fig. 1. Study flow chart

participants were excluded due to osteoporosis, and two participants were unable to jump properly. One participant refrained from further participation due to personal reasons. Twenty-nine of the 35 women who initially responded were, therefore, included. The baseline characteristics of the participants, including $\text{VO}_2\text{-max}$ as an indication of training status, are reported in Table 1.

Participant characteristics at baseline

Proximal femur (total hip), lumbar spine (LS), and whole body (WB) DXA scans (iDXA, Lunar Corporation, Madison, Wisconsin, USA) were performed according to standard procedures to determine BMD (g/cm^2). The regions of interest were determined automatically by the software (Encore Version 14.10.022, GE Medical Systems, Madison, Wisconsin, USA). Body composition parameters inclusive body fat percentage (%), lean mass (kg) and fat mass (kg) were derived from WB DXA. Subjects were requested to remove metal objects and void their bladder prior to DXA scanning.

Training status

To confirm the participants' training status, maximal oxygen uptake ($\text{VO}_2\text{-max}$) ($\text{ml}/\text{kg}/\text{min}$) was tested on an electronic ergometer cycle (Monark 839E, Monark Exercise AB, Vansbro, Sweden). Participants were connected to a breath-by-breath gas online analyzing system (Jaeger Oxycon Pro, VIASYS Healthcare, Höchberg, Germany), and direct measurement of $\text{VO}_2\text{-max}$ was performed according to standard procedures for a progressive test.

High-impact exercise trials

A standardized 7-minute low-impact warm-up on a 3.5 cm gymnastics mat preceded each exercise trial. Following the warm-up, the participants were instructed in the actual jump of the day: CMJ, a vertical jump with two-leg launch and landing; DJ, drop jump from a 32 cm box: the landing continued directly into a vertical two-leg jump; DDJ, as DJ, but performed diagonally forward (45°).

All jumps (6 sets of 10 repetitions) were performed on a gymnastics mat, with each set of jumps interspersed with a 90 s rest. The participants were encouraged to "do all their best" in every jump by jumping as high and powerfully as possible with an arm swing in the set-off phase and a sudden stop in the landing. Instructors gave ongoing motivation and corrections if needed.

Blood sampling and biochemical analyses

The plasma concentrations of BTM at baseline (PRE), immediately after exercise (POST) and after two hours of rest (2Hr) were measured. As dietary intake can affect BTM [22], the participants showed up in the early morning after an overnight fast and without any vigorous exercise in the preceding 48 hours. Dietary supplements were not allowed. While waiting from POST to 2Hr, participants drank a glass of water. The blood samples were collected from the antecubital vein with a butterfly needle. Each participant

had extracted $3 \times 6 \text{ ml} = 18 \text{ ml}$ of blood per test day. After each test, the plasma fractions were stored at -80°C until analysis. The bone formation markers procollagen type-1 amino-terminal propeptide (P1NP) and osteocalcin (OC), and the bone resorption marker C-terminal telopeptide of type-1 collagen (CTX), were evaluated by a fully automated immunoassay system (iSYS, Immunodiagnostic Systems Ltd., Bolton, England) by the method of chemiluminescence. The performance of the assay expressed as inter-run coefficients of variation was 9% for OC, 10% for CTX, and 8% for P1NP.

Ground reaction force

Nineteen of the 29 participants agreed to have their GRF in the three jumps measured on a separate day with an AMTI (Advanced Mechanical Technology Inc., Watertown, MA 02472–4800 USA) SGA6–4 force platform. The signal was converted using an AD-converter (National Instruments) with 16-bit resolution, and the data were accumulated using the MATLAB data acquisition toolbox (2009b; The MathWorks, Inc., Natick, USA). The GRF components were evaluated in three directions: the anteroposterior (Fx), the mediolateral (Fy), and the vertical peak GRF (Fz). Peak GRF in every direction were determined from the entire phase and normalized to body weight (BW).

The jumping set-up was standardized in accordance with the jumping trials, participants wore the same shoes as during previous jumping experiments in which BTM was measured, the same landing mat was taped on the force platform, and the techniques of each of the three jumps were initially repeated for the participants. After 2–3 attempts, GRF was measured: Each participant performed six repetitions of each jump, the highest and the lowest of the measured peak GRF were excluded, and the mean of the remaining peak GRF were used for further analyses.

Statistical analysis

SPSS, version 24.0 (IBM Corp., Armonk, NY, USA) was used for all statistical analyses. Standard descriptive statistics were used to describe the participant characteristics at baseline. Linear mixed models tested the difference of BTM between PRE, POST, and 2Hr. Repeated measures analysis of variance (ANOVA) with post hoc Bonferroni adjustments for multiple comparisons were used to test the difference in GRF between the jumps. Linear regression analyses were used to assess the correlation between the ΔBTM and the associated GRF in each type of jump. A $p\text{-value} < 0.05$ was considered significant. Unless otherwise stated, values are given as mean \pm standard error (SE).

RESULTS

Plasma BTM

Bone formation and resorption markers are presented in Table II.

For all BTM there were no statistically significant differences in Δ -values between the three types of jumps at any time point.

Compared to PRE, P1NP POST was significantly increased for all jumps: $5.4 \pm 1.6 \mu\text{g}/\text{L}$ ($p < 0.005$) for CMJ; $6.7 \pm 0.9 \mu\text{g}/\text{L}$ ($p < 0.001$)

Table 2. Bone turnover marker (BTM) concentrations at baseline (PRE), immediately after (POST), and 2 hours after exercise (2Hr)

Outcome	Jump	PRE	POST	2Hr
Markers of bone formation				
P1NP ($\mu\text{g/L}$)	CMJ	70.2 \pm 5.6	75.6 \pm 6.3***#	68.7 \pm 6.0
	DJ	71.0 \pm 5.5	77.6 \pm 5.8***##	67.5 \pm 6.0
	DDJ	73.0 \pm 6.3	80.8 \pm 6.8***##	70.2 \pm 6.0
	CON	71.9 \pm 5.3	70.1 \pm 5.6	70.6 \pm 5.4
OC ($\mu\text{g/L}$)	CMJ	31.2 \pm 2.3	32.2 \pm 2.4	28.9 \pm 2.2**
	DJ	30.7 \pm 2.2	32.4 \pm 2.5*#	28.3 \pm 2.5**
	DDJ	30.6 \pm 2.2	31.8 \pm 2.3	29.2 \pm 2.2
	CON	31.1 \pm 2.1	30.0 \pm 2.0	28.1 \pm 2.0***
Marker of bone resorption				
CTX (ng/L)	CMJ	636.0 \pm 83.4	635.5 \pm 80.3	527.9 \pm 65.7 ***
	DJ	645.2 \pm 88.3	666.2 \pm 91.0	525.5 \pm 69.0 ***
	DDJ	612.8 \pm 85.9	632.8 \pm 85.4	519.0 \pm 69.1 ***
	CON	590 \pm 73.6	582.4 \pm 74.4	501.7 \pm 65.8 ***

All values are expressed as mean \pm SE (n=29). P1NP = procollagen type-1 amino-terminal propeptide. OC = osteocalcin. CTX = C-terminal telopeptide of type-1 collagen. CMJ = countermovement jump. DJ = drop jump. DDJ = diagonal drop jump. CON = control.

* $p < 0.05$ compared to PRE. ** $p < 0.01$ compared to PRE. *** $p < 0.001$ compared to PRE. # $p < 0.05$ compared to C. ## $p < 0.01$ compared to C. ### $p < 0.001$ compared to C.




Fig. 2. Anteroposterior (Fx), mediolateral (Fy), and vertical (Fz) components of the peak ground reaction forces (GRF) normalized to body weight (BW) for the three high-impact jumping exercises. CMJ = Countermovement jump. DJ = Drop jump. DDJ = Diagonal drop jump.

Data presented as mean \pm SE (n=19).

* $p < 0.05$. ** $p < 0.01$ Statistical testing of between-group differences using repeated measures analysis of variance (ANOVA) with post hoc pairwise comparison (Bonferroni adjustment).

for DJ; and $7.8 \pm 1.1 \mu\text{g/L}$ ($p < 0.001$) for DDJ. Compared to CON, the increases were all significantly higher for all jumps: $7.2 \pm 1.3 \mu\text{g/L}$ ($p < 0.01$) for CMJ; $8.5 \pm 1.2 \mu\text{g/L}$ ($p = 0.001$) for DJ; and $9.6 \pm 1.1 \mu\text{g/L}$ ($p < 0.001$) for DDJ. P1NP 2Hr for CMJ, DJ, and DDJ did not differ significantly from PRE.

Compared to PRE, OC POST was significantly increased by $1.7 \pm 0.7 \mu\text{g/L}$ ($p < 0.05$) for DJ, but not for CMJ or DDJ. The increase was $2.8 \pm 0.8 \mu\text{g/L}$ ($p < 0.05$) higher than CON. Compared to PRE, OC 2Hr was decreased by $2.3 \pm 0.5 \mu\text{g/L}$ ($p < 0.01$) for CMJ, by $2.5 \pm 1.1 \mu\text{g/L}$ ($p < 0.005$) for DJ, and by $3.0 \pm 0.5 \mu\text{g/L}$ ($p < 0.001$) for CON.

Compared to PRE, CTX POST did not change significantly in any trial. CTX 2Hr was significantly decreased in all trials: by $108.1 \pm 22.9 \text{ ng/L}$ ($p < 0.001$) for CMJ; by $119.7 \pm 25.7 \text{ ng/L}$ ($p < 0.001$) for DJ; by $93.8 \pm 25.2 \text{ ng/L}$ ($p < 0.001$) for DDJ; and by $88.3 \pm 15.7 \text{ ng/L}$ ($p < 0.001$) for CON.

Ground reaction forces

Peak GRF in anteroposterior (Fx), mediolateral (Fy) and vertical (Fz) components of the GRF normalized to body weight, for the three jumps, are presented in Figure 2.

There was a difference ($p < 0.01$) between jumps in Fx and Fy. Subsequent paired comparisons showed that Fx for DJ was




Fig. 3. Relationship between peak ground reaction forces normalized to body weight and the change in procollagen type-1 amino-terminal propeptide immediately after exercise (Δ P1NP POST) for the countermovement jump (CMJ). (a) Vertical peak ground reaction force. (b) Combined peak ground reaction force. P1NP = procollagen type-1 amino-terminal propeptide. POST = immediately after exercise. CMJ = countermovement jump. GRF = ground reaction force. Fx = anteroposterior. Fy = mediolateral. Fz = vertical. BW = body weight.

0.211±0.05BW higher than for CMJ (0.874±0.04 vs. 0.663±0.07BW, $p < 0.005$) and 0.158±0.04BW higher than for DDJ (0.874±0.04 vs. 0.717±0.03BW, $p < 0.005$). There was no difference in Fx between CMJ and DDJ ($p = 1.00$). Moreover, Fy for DDJ was 0.122±0.03BW higher than for CMJ (0.366±0.03 vs. 0.244±0.03BW, $p < 0.005$) and 0.072±0.02BW higher than for DJ (0.366±0.03 vs. 0.294±0.02BW, $p < 0.05$). Differences between CMJ and DJ in Fy ($p = 0.056$) and Fz ($p = 0.06$) did not reach significance.

Multiple linear regression analyses of the correlations between BTM Δ -values and the three directions of peak GRF (Fx, Fy, Fz) showed that only for CMJ, Δ P1NP POST correlated significantly with Fz ($r = 0.61$, $p < 0.01$) (Fig. 3a.) and with the combined Fx, Fy, and Fz ($r = 0.71$, $p < 0.05$) (Fig. 3b.). The analyses resulted in the following equation: Δ P1NP = -17.71 + (-16.97*Fx) + 25.25*Fy + 7.33*Fz ($R^2 = 0.504$). Thus, the combination of Fx, Fy, and Fz explained 50.4% (R^2) of the Δ P1NP variation in the CMJ, while Fz explained 37.2%.

DISCUSSION

To the authors' knowledge, the present study is the first to examine the acute biological response of BTM, and the resultant GRF, of three high-impact jumping exercises in postmenopausal women.

The finding that P1NP was significantly increased at POST for both countermovement jump (7.7±1.8%), drop jump (9.4±1.3%), and diagonal drop jump (10.6±1.6%), which differed significantly from the no-exercise control trial ($p < 0.01$), seems to indicate that the P1NP response was due to exercise and not to diurnal variation. The acute increase in P1NP is consistent with the studies of Brahm et al. [10], Bowtell et al. [9], and Scott et al. [17], who respectively investigated the osteogenic response to resistance, odd-impact, and aerobic training. They found an increase from 9.5 to 31% [9,10,17], which is in a higher range compared to the present study. However, the larger response may be due to the younger age of the participants (21–36 year old men and women [10], 26–46 year old premenopausal women [9], and 28 year old men on average [17]) predisposing to larger maximal muscle strength and larger osteogenic responsiveness. In contrast to these findings, Mouzopoulos et al. [23] showed that the bone formation marker procollagen type-1 carboxy-terminal propeptide (P1CP) decreased by 8.92% after an ultra-marathon run. However, it was hypothesized that this reduction was probably due to intensive mechanical overloading that temporarily may inhibit collagen synthesis [24]. In addition, a study by Cabrera et al. showed that P1CP had less sensitivity and accuracy in response to interventions [25].

Two major strengths of the present study are the supervised and highly controlled jumping exercises and the precise timing of blood sampling on each trial day, minimizing the effect of diurnal variation when comparing BTM concentrations from the four days. It is known that diurnal variation in bone resorption markers represents up to 50% of the variation depending on the time of day [26]. However, no major day-to-day variation in OC [27], P1NP [28], or CTX [29] concentration has been observed when the samples have been collected at a similar time of day. This is supported by our findings since we did not find any significant differences between baseline samples on the individual trial days.

Our finding that P1NP concentrations returned to baseline two hours after exercise is consistent with prior studies [14,17] that showed insignificant Δ -values 2 hours after cessation of high-impact exercise in young males.

Other studies [29,30] examining the change in BTM three hours after a 60-minute bout of aerobic exercise reported that P1NP and P1CP decreased significantly in young subjects. It was speculated that the reason for the decrease could be exercise-induced acidosis that might impair osteoblast activity [31], which might also be the reason for the return of P1NP to baseline two hours after cessation of exercise in the present study. Since we found significant osteogenic improvements after the jumping trials compared to the control trial, the increase in Δ P1NP POST does not seem to be due to circadian rhythms.

OC was only significantly increased at POST for DJ ($5.48 \pm 1.76\%$, $p < 0.05$). Nevertheless, there was a non-significant increase in OC POST for DDJ ($3.84 \pm 1.42\%$, $p = 0.057$) and CMJ ($3.26 \pm 1.11\%$, $p = 0.076$), which together with the increase in P1NP might reflect an immediate anabolic effect of exercise on bone, which is supported by Maimoun *et al.* [12], who reported a significant increase (11%) in OC after 50 minutes of high-intensity resistance exercise in male cyclists. By contrast, OC decreased significantly (6.8%, $p < 0.05$) following 30 minutes of resistance exercise in young sedentary males and females [10], and decreased after a 245 km ultra-marathon run in young athletes (17.4%, $p < 0.05$) [23]. These inconsistent results confirm that, besides the musculoskeletal intensity of exercise, exercise duration and repetitiveness have an important influence on the biochemical bone marker response to acute exercise. As long exhausting exercise is known to elicit an increase in cortisol, this may induce inhibition of osteoblast function [32], leading to decreased osteocalcin concentrations after a ultra-marathon run, but also desensitization of mechanotransduction mechanisms may be responsible [33].

Our findings of significantly decreased OC concentrations at 2Hr for CMJ and DJ are in agreement with Herrmann *et al.* [30] and Scott *et al.* [17], who reported similar OC reductions after three hours of aerobic training. However, the time of the day for blood sampling was not specified in those studies, and no resting control trial was included, which makes it impossible to control for normal diurnal variation, characterized by a decline in the morning, reaching a nadir around noon, and a peak after midnight [34]. Since the 2Hr OC also decreased during the no-exercise (CON) trial in the present study, we hypothesize that the reported 2Hr OC Δ -values are due to diurnal variation, which might also be the case for the reported reductions after three hours [17,30].

CTX, generated by cathepsin K activity, is recommended as a bone resorption marker [35]. In the present study, there was no change in CTX concentrations at POST in any of the jumps, and the significant decrease in 2Hr CTX after all jumps did not differ from the control trial. Other studies [15,17] have also reported a reduced CTX concentration 2 hours following high-impact exercise in men, although with no comparison to a controlled trial to test the diurnal variation. The finding of no significant difference in 2Hr CTX between the four trials in the present study leads us to the conclusion that the CTX reductions were merely due to diurnal variation and not to exercise,

which is in agreement with Wichers *et al.* [36], who reported a decrease in CTX between 08:00 and 11:00 reaching the lowest concentration between 11:00 and 15:00.

The findings of the present study indicate that a single bout (6 sets of 10 repetitions) of high-impact jumping exercise may elicit an osteogenic effect via an increase in the rate of bone formation compared with CON; however, no differences were detected between the CMJ, DJ and DDJ at any time point.

Normally, only the vertical peak GRF (Fz) is evaluated in exercise studies on GRF, but in the present study the anteroposterior (Fx) and the mediolateral peak GRF (Fy) were also assessed. As bone strain [33] and the impact from unusual and odd directions are especially osteogenic [37,38] we considered it logical to evaluate the combined osteogenic impact from GRF in three directions. The significantly highest anteroposterior peak GRF (Fx) was reached in the DJ (0.874 ± 0.04 BW), which includes a forward movement when dropping from the box, unlike the CMJ, which is a vertical jump on the same spot, and the DDJ, which includes a forward diagonal movement. The significantly highest mediolateral peak GRF (Fy) was, as expected, reached in the DDJ (0.366 ± 0.03 BW), which also induced the highest Δ -value of P1NP POST (from 73.01 ± 6.27 to 80.78 ± 6.77 $\mu\text{g/L}$, $p < 0.001$) indicating that the DDJ had a superior osteogenic impact. However, there was no correlation ($R^2 = 0.09$, $p = 0.21$) between the Δ -value of P1NP POST and Fy in DDJ, which may be explained by the relatively low Fy in all jumps compared to especially Fz, but also to Fx. Thus, the isolated impact of Fy may only be of low osteogenic value.

Given that the vertical GRF (Fz) during the three jumping protocols averaged 4 times BW, the impact met the moderate- to high-intensity domain criteria defined by Witzke and Snow for adolescent girls [39]. However, in the present study a lower high-intensity range of GRF may be expected due to lower BMD and weaker bones in post-menopausal women. Thus, a lower GRF would be needed to induce an osteogenic strain.

The finding that Fz did not significantly differ between the jumps, even though Fz in DJ was non-significantly higher than in CMJ ($p = 0.06$), is in agreement with Smale *et al.* [40], who reported that for postmenopausal women, Fz in CMJ did not differ from Fz in DJ (second landing). However, both results are in disagreement with the study of Weeks and Beck [41] showing that DJ had a significantly higher peak vertical GRF compared to CMJ (DJ 5.5BW vs. CMJ 4.7BW). However, the participants [41] differed markedly from the present study as well as from the study of Smale *et al.* [40] by being moderately physically active, young adult males and females. Thus, the younger age of the participants could be the reason why Fz was higher than in the present study (DJ 4.0 ± 0.16 BW vs. CMJ 3.7 ± 0.16 BW) due to better coordination and muscle power in the take-off and landing phase and less variation in jumping performance in the younger group.

To evaluate the overall osteogenic effect of peak GRF in three directions we tested the correlation between the combined three-

axis peak GRF (F_x, F_y, and F_z) and Δ P1NP POST, and our finding of a significant correlation for CMJ indicates a clear dose-response relationship between GRF and the acute BTM response with GRF explaining 50.4% ($r=0.71$) of the variation in the P1NP response. A dose-response relationship is also demonstrated by Rantalainen et al. [14], but to a lesser extent. Thus, they report a significant correlation ($r=0.49$) between vertical peak GRF and acute Δ P1NP after continuous bilateral jumping using the ankle plantar flexors at 65% of maximal GRF until exhaustion [14]. We hypothesize that the higher r -value in the present study is mainly caused by the higher musculoskeletal exercise intensity and the higher age of the participants, which theoretically will give rise to a larger bone strain due to weaker bones compared to the physically active young participants [14].

Additionally, the three-directional GRF value in the present study may have an important impact on the correlation between GRF and Δ P1NP. Since R^2 obtained from multivariate models ($R^2=50.4\%$) is greater than R^2 between F_z and Δ P1NP ($R^2=37.2\%$), the impact obtained from all three directions combined seems to influence the osteogenic response positively. However, we did not find any correlation between (combined or separated) GRF and Δ BTM in the two drop-jumps (DJ and DDJ), which, based on the largest F_x in DJ and largest F_y in DDJ, was in contrast to our hypothesis. We assume that the explanation for this finding is the relative difficulty of DJ and DDJ, which was evident from the GRF curves, showing a considerable variation in GRF over the whole landing phase between the participants as well as between the six repetitions for each individual. The average difference between the minimum and maximum individual F_z in the six repetitions was 1.34BW for the DJ and 1.33BW for the DDJ compared to 1.17BW for the CMJ. This indicates that the CMJ was performed with the most stable results, maybe because it was easier for the participants to coordinate than DJ and DDJ.

A possible limitation of the present study is the fact that the measurements of GRF and BTM were not performed on the same occasion. Therefore, it cannot be ruled out that the jumping technique, and therefore the GRF, might differ slightly between the two test days. In addition, only a subgroup of participants was willing to having their GRF measured, which could introduce a bias due to self-selection. However, there was no difference in participant characteristics

between the two subgroups.

CONCLUSIONS

The evaluation of the acute biological response in plasma BTM induced by three different high-impact jumps in postmenopausal women showed stimulation of bone modelling, reflected by the significant increase in the biochemical formation markers P1NP and OC concomitant with an unchanged resorption marker (CTX). There was no difference in BTM response between the jumps, but a significantly larger increase in P1NP and OC when compared to the resting control condition. In the countermovement jump, the higher correlation between the acute P1NP response and the combined vertical, anteroposterior and mediolateral GRF than with the vertical peak GRF alone indicates that not only vertical but also mediolateral and anteroposterior peak GRF exert an osteogenic stimulus on bone turnover. This dose-response relationship may be important when planning training aimed at improving bone mass and reducing the risk of osteoporosis.

Acknowledgments

We thank all the participants for their time and effort. The authors thank Jens J. Nielsen for assistance in blood draws, Thomas Nyegaard Beck for assistance in VO₂-max testing, Therese Hornstrup Bondebjerg and Marie Hagman for help in the laboratory, Nadia Quardon for blood sample analysis, Erik B. Simonsen and Pia Melcher for guidance in using the force platform, Christian Ritz for statistical support. We thank the Department of Biomedical Sciences and Xlab at the Faculty of Health and Medical Sciences for their excellent hospitality. RSP gratefully acknowledges a Ph.D. fellowship from Lembaga Pengelola Dana Pendidikan (LPDP)/Indonesia Endowment Fund for Education. The authors have no professional relationships with companies or manufacturers that might benefit from this study.

Disclosures

No conflicts of interest, financial or otherwise, are declared by the author(s).

REFERENCES

- Kanis JA, Melton LJ, Christiansen C, Johnston CC, Khaltav N. The diagnosis of osteoporosis. *J Bone Miner Res.* 1994;9(8):1137–41.
- Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int.* 2006;17(12):1726–33.
- Xu J, Lombardi G, Jiao W, Banfi G. Effects of Exercise on Bone Status in Female Subjects, from Young Girls to Postmenopausal Women: An Overview of Systematic Reviews and Meta-Analyses. *Sports Med.* 2016;46(8):1165–82.
- Troy KL, Mancuso ME, Butler TA, Johnson JE. Exercise Early and Often: Effects of Physical Activity and Exercise on Women's Bone Health. *Int J Environ Res Public Health.* 2018; 15(5):878.
- Martyn-St James M, Carroll S. A meta-analysis of impact exercise on postmenopausal bone loss: the case for mixed loading exercise programmes. *Br J Sports Med.* 2009; 43(12):898–908.
- Miller PD, Zapalowski C, Kulak CAM, Bilezikian JP. Bone Densitometry: The Best Way to Detect Osteoporosis and to Monitor Therapy. *J Clin Endocrinol Metab.* 1999;84(6):1867–71.
- Delmas PD, Eastell R, Garnero P, Seibel MJ, Stepan J, Foundation for the C of SA of the IO. The Use of Biochemical Markers of Bone Turnover in Osteoporosis. *Osteoporos Int.* 2000;11(6):S2–17.

8. Banfi G, Lombardi G, Colombini A, Lippi G. Bone metabolism markers in sports medicine. *Sports Med Auckl NZ*. 2010;40(8):697–714.
9. Bowtell JL, Jackman SR, Scott S, Connolly LJ, Mohr M, Ermidis G, Julian R, Yousefian F, Helge EW, Jørgensen NR, Fulford J, Knapp KM, Krstrup P. Short Duration Small Sided Football and to a Lesser Extent Whole Body Vibration Exercise Induce Acute Changes in Markers of Bone Turnover. *BioMed Res Int*. 2016; 2016:e3574258.
10. Brahm H, Piehl-Aulin K, Saltin B, Ljunghall S. Net fluxes over working thigh of hormones, growth factors and biomarkers of bone metabolism during short lasting dynamic exercise. *Calcif Tissue Int*. 1997;60(2):175–80.
11. Kish K, Mezil Y, Ward WE, Klentrou P, Falk B. Effects of plyometric exercise session on markers of bone turnover in boys and young men. *Eur J Appl Physiol*. 2015;115(10):2115–24.
12. Maimoun L, Manetta J, Couret I, Dupuy AM, Mariano-Goulart D, Micallef JP, Peruchon E, Rossi M. The intensity level of physical exercise and the bone metabolism response. *Int J Sports Med*. 2006;27(2):105–11.
13. Mezil YA, Allison D, Kish K, Ditor D, Ward WE, Tsiani E, Klentrou P. Response of Bone Turnover Markers and Cytokines to High-Intensity Low-Impact Exercise. *Med Sci Sports Exerc*. 2015; 47(7):1495–502.
14. Rantalainen T, Heinonen A, Linnamo V, Komi PV, Takala TES, Kainulainen H. Short-term bone biochemical response to a single bout of high-impact exercise. *J Sports Sci Med*. 2009;8(4):553–9.
15. Rogers RS, Dawson AW, Wang Z, Thyfault JP, Hinton PS. Acute response of plasma markers of bone turnover to a single bout of resistance training or plyometrics. *J Appl Physiol Bethesda Md* 1985. 2011;111(5):1353–60.
16. Rong H, Berg U, Torring O, Sundberg CJ, Granberg B, Bucht E. Effect of acute endurance and strength exercise on circulating calcium-regulating hormones and bone markers in young healthy males. *Scand J Med Sci Sports*. 1997; 7(3):152–9.
17. Scott JPR, Sale C, Greeves JP, Casey A, Dutton J, Fraser WD. The role of exercise intensity in the bone metabolic response to an acute bout of weight-bearing exercise. *J Appl Physiol*. 2011; 110(2):423–32.
18. Wallace JD, Cuneo RC, Lundberg PA, Rosén T, Jørgensen JO, Longobardi S, Keay N, Sacca L, Christiansen JS, Bengtsson BA, Sönksen PH. Responses of markers of bone and collagen turnover to exercise, growth hormone (GH) administration, and GH withdrawal in trained adult males. *J Clin Endocrinol Metab*. 2000;85(1):124–33.
19. Welsh L, Rutherford OM, James I, Crowley C, Comer M, Wolman R. The acute effects of exercise on bone turnover. *Int J Sports Med*. 1997; 18(4):247–51.
20. Wherry S, Swanson C, Wolfe P, Wellington T, Boxer R, Schwartz R, Kohrt W. Bone Biomarker Response to Walking under Different Thermal Conditions in Older Adults. *Med Sci Sports Exerc*. 2019;51(8):1599–605.
21. Thorsen K, Kristoffersen A, Lorentzon R. The effects of brisk walking on markers of bone and calcium metabolism in postmenopausal women. *Calcif Tissue Int*. 1996;58(4):221–5.
22. Park SM, Joung JY, Cho YY, Sohn SY, Hur KY, Kim JH, Kim SW, Chung JH, Lee MK, Min Y-K. Effect of high dietary sodium on bone turnover markers and urinary calcium excretion in Korean postmenopausal women with low bone mass. *Eur J Clin Nutr*. 2015; 69(3):361–6.
23. Mouzopoulos G, Stamatakos M, Tzurbakis M, Tsernbeli A, Manti C, Safioleas M, Skandalakis P. Changes of bone turnover markers after marathon running over 245 km. *Int J Sports Med*. 2007;28(7):576–9.
24. Virtanen P, Viitasalo JT, Vuori J, Väänänen K, Takala TE. Effect of concentric exercise on serum muscle and collagen markers. *J Appl Physiol*. 1993;75(3):1272–7.
25. Cabrera CD, Henríquez MS, Traba ML, Villafañe EA, de la Piedra C. Biochemical markers of bone formation in the study of postmenopausal osteoporosis. *Osteoporos Int*. 1998;8(2):147–51.
26. Ebeling PR, Akesson K. Role of biochemical markers in the management of osteoporosis. *Best Pract Res Clin Rheumatol*. 2001;15(3):385–400.
27. Srivastava AK, Bhattacharyya S, Li X, Mohan S, Baylink DJ. Circadian and longitudinal variation of serum C-telopeptide, osteocalcin, and skeletal alkaline phosphatase in C3H/HeJ mice. *Bone*. 2001;29(4):361–7.
28. Munday K, Ginty F, Fulford A, Bates CJ. Relationships between biochemical bone turnover markers, season, and inflammatory status indices in prepubertal Gambian boys. *Calcif Tissue Int*. 2006;79(1):15–21.
29. Zittermann A, Sabatschus O, Jantzen S, Platen P, Danz A, Stehle P. Evidence for an acute rise of intestinal calcium absorption in response to aerobic exercise. *Eur J Nutr*. 2002; 41(5):189–96.
30. Herrmann M, Mueller M, Scharhag J, Sand-Hill M, Kindermann W, Herrmann W. The effect of endurance exercise-induced lactacidosis on biochemical markers of bone turnover. *Clin Chem Lab Med*. 2007; 45(10):1381–9.
31. Brandao-Burch A, Utting JC, Orriss IR, Arnett TR. Acidosis Inhibits Bone Formation by Osteoblasts In Vitro by Preventing Mineralization. *Calcif Tissue Int*. 2005;77(3):167–74.
32. Fellmann N, Bedu M, Giry J, Pharmakis-Amadiou M, Bezou MJ, Barlet JP, Coudert J. Hormonal, fluid, and electrolyte changes during a 72-h recovery from a 24-h endurance run. *Int J Sports Med*. 1989;10(6):406–12.
33. Rubin CT, Lanyon LE. Regulation of bone formation by applied dynamic loads. *J Bone Joint Surg Am*. 1984; 66(3):397–402.
34. Gundberg CM, Markowitz ME, Mizruchi M, Rosen JF. Osteocalcin in Human Serum: A Circadian Rhythm. *J Clin Endocrinol Metab*. 1985; 60(4):736–9.
35. Vasikaran S, Eastell R, Bruyère O, Foldes AJ, Garnero P, Griesmacher A, McClung M, Morris HA, Silverman S, Trenti T, Wahl DA, Cooper C, Kanis JA, IOF-IFCC Bone Marker Standards Working Group. Markers of bone turnover for the prediction of fracture risk and monitoring of osteoporosis treatment: a need for international reference standards. *Osteoporos Int*. 2011;22(2):391–420.
36. Wichers M, Schmidt E, Bidlingmaier F, Klingmüller D. Diurnal Rhythm of CrossLaps in Human Serum. *Clin Chem*. 1999;45(10):1858–60.
37. Helge EW, Andersen TR, Schmidt JF, Jørgensen NR, Hornstrup T, Krstrup P, Bangsbo J. Recreational football improves bone mineral density and bone turnover marker profile in elderly men. *Scand J Med Sci Sports*. 2014; 24 Suppl 1:98–104.
38. Nikander R, Sievänen H, Heinonen A, Daly RM, Uusi-Rasi K, Kannus P. Targeted exercise against osteoporosis: A systematic review and meta-analysis for optimising bone strength throughout life. *BMC Med*. 2010;8:47.
39. Witzke KA, Snow CM. Effects of plyometric jump training on bone mass in adolescent girls. *Med Sci Sports Exerc*. 2000;32(6):1051.
40. Smale KB, Hansen LH, Kristensen JK, Zebis MK, Andersen C, Benoit DL, Helge EW, Alkjaer T. Loading intensity of jumping exercises in post-menopausal women: Implications for osteogenic training. *Transl Sports Med*. 2018; 1(1):30–6.
41. Weeks BK, Beck BR. The BPAQ: a bone-specific physical activity assessment instrument. *Osteoporos Int*. 2008;19(11):1567–77.