



Search for long-lived neutral particles produced in pp collisions at root s=13 TeV decaying into displaced hadronic jets in the ATLAS inner detector and muon spectrometer

ATLAS Collaboration

Published in:
Physical Review D

DOI:
[10.1103/PhysRevD.101.052013](https://doi.org/10.1103/PhysRevD.101.052013)


Publication date:
2020

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
ATLAS Collaboration (2020). Search for long-lived neutral particles produced in pp collisions at root s=13 TeV decaying into displaced hadronic jets in the ATLAS inner detector and muon spectrometer. *Physical Review D*, 101(5), [052013]. <https://doi.org/10.1103/PhysRevD.101.052013>

Search for long-lived neutral particles produced in pp collisions at $\sqrt{s} = 13$ TeV decaying into displaced hadronic jets in the ATLAS inner detector and muon spectrometer

G. Aad *et al.**
(ATLAS Collaboration)

 (Received 2 December 2019; accepted 14 February 2020; published 30 March 2020)

A search is presented for pair production of long-lived neutral particles using 33 fb^{-1} of $\sqrt{s} = 13$ TeV proton–proton collision data, collected during 2016 by the ATLAS detector at the LHC. This search focuses on a topology in which one long-lived particle decays in the ATLAS inner detector and the other decays in the muon spectrometer. Special techniques are employed to reconstruct the displaced tracks and vertices in the inner detector and in the muon spectrometer. One event is observed that passes the full event selection, which is consistent with the estimated background. Limits are placed on scalar boson propagators with masses from 125 GeV to 1000 GeV decaying into pairs of long-lived hidden-sector scalars with masses from 8 GeV to 400 GeV. The limits placed on several low-mass scalars extend previous exclusion limits in the range of proper lifetimes $c\tau$ from 5 cm to 1 m.

DOI: [10.1103/PhysRevD.101.052013](https://doi.org/10.1103/PhysRevD.101.052013)

I. INTRODUCTION

Long-lived particles (LLPs) are predicted by many extensions of the Standard Model (SM), including various supersymmetric (SUSY) [1–4], hidden sector (HS) [5–7], and neutral naturalness [8–10] models that address the hierarchy problem. Decays of LLPs may go unnoticed in searches at collider experiments that are designed to identify promptly decaying particles. Searches for LLPs provide a promising avenue for the discovery of physics beyond the Standard Model (BSM). The search presented in this paper is sensitive to neutral LLPs that are pair produced, with one LLP decaying in the ATLAS inner tracking detector (ID) and the other in the muon spectrometer (MS). This particular event topology provides sensitivity to LLPs with proper lifetimes ($c\tau$) ranging from a few centimeters to several meters.

In HS models, a set of BSM particles is weakly coupled to the SM via a mediator particle. These models are intriguing because they can be built in multiple ways and can produce LLPs with little to no fine-tuning [6]. A SM Higgs boson mediator is of particular interest because the current experimental characterization of the Higgs boson allows sizable couplings of the Higgs boson to the BSM sector [11,12]. HS models are also compatible

with SUSY [6,13] and with models of neutral naturalness [14]. The results of this search are interpreted in the context of a simplified HS model, in which the SM and HS are connected via a heavy mediator Φ , which decays into a pair of long-lived neutral scalar particles s as shown in Fig. 1. The s -bosons then decay back into SM particles through their mixing with the mediator [15]. The HS model assumes an effective Yukawa coupling between the s -boson and the SM particles; therefore each s -boson decays primarily into a heavy fermion pair: $b\bar{b}$, $c\bar{c}$, $\tau^+\tau^-$. The branching ratio depends on the mass of the s -boson (m_s), but for $m_s > 25$ GeV, the branching ratio is approximately 85:5:8.¹ If the SM particles are quarks, they hadronize, resulting in jets that may be highly displaced from the interaction point (IP). The proper lifetime of the s -boson is relatively unconstrained aside from the upper limit imposed by big bang nucleosynthesis of $c\tau \lesssim 10^8$ m [18].

Searches for displaced hadronic decays have been performed by ATLAS, CMS, and LHCb in the first run of the LHC [19–26], as well as in the second run of the LHC by CMS [27] and ATLAS [16,17,28,29]. Two searches for displaced decays resulting from neutral LLPs in the HS model have been performed using the ATLAS Run 2 data set. One analysis searched for pairs of displaced hadronic jets in the calorimeter [16] (the CR analysis), and the other searched for one or two displaced

*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

¹As in recent displaced jet searches for HS models [16,17], decays of s to bosons and top quarks are not considered in this analysis.




FIG. 1. Diagram for a Higgs boson or heavy scalar Φ decaying into displaced hadronic jets via a hidden sector.

hadronic jets in the MS [17] (the MS analysis). For a SM Higgs boson mediator with $m_H = 125$ GeV, decays of neutral scalars with masses between 8 and 55 GeV have been excluded by these two analyses for $c\tau$ between 7 cm and 220 m depending on the LLP mass (assuming a 10% branching ratio of the Higgs boson into ss pairs).

This analysis uses 33.0 fb^{-1} of $\sqrt{s} = 13$ TeV proton-proton (pp) collision data collected by the ATLAS detector at the LHC and is an update to the results presented in the 8 TeV ATLAS search [21] for displaced hadronic jets in the ID and MS, and an extension of the MS analysis. In each event, one reconstructed decay vertex is required in the ID (IDVx) in addition to one in the MS (MSVx). Requiring the presence of an MSVx suppresses background, which allows looser selection requirements on the reconstructed mass and number of tracks associated with the IDVx than in other IDVx searches such as Ref. [28]. Additionally, requiring the presence of an IDVx suppresses background relative to the MS analysis, allowing greater sensitivity for LLPs with $c\tau < 1$ m. This search thus increases the sensitivity to low-mass scalars with shorter proper lifetimes relative to the combined results of the CR and MS analyses published in Ref. [16].

A signature-driven trigger that is used to collect data selects candidate events for decays of LLPs in the MS [17,30]. Standard ATLAS reconstruction methods are optimized for prompt decays and may fail to reconstruct the decays of long-lived particles. Thus, specialized tracking and vertex reconstruction algorithms are used that allow for the reconstruction of displaced decays in the ID [28,31,32] and the MS [33].

The main background source of signal-like vertices in the ID is the interactions between SM particles and the material in the inner detector, which may create multitrack vertices that are displaced from the IP. Other sources of background are hadronic jets, vertices reconstructed from fake tracks that are created from multiple unrelated energy deposits, and vertices reconstructed from random track crossings. The main source of background vertices in the MS are electromagnetic or hadronic showers that are not contained in the calorimeters (punch-through jets). Other sources of background in the MS are multijet events with mismeasured jets and noncollision backgrounds such as machine-induced background [34,35], electronic noise, and

cosmic-ray muons. A data-driven background estimation is used to determine how many events in the signal region result from background sources.

This paper is organized as follows. The ATLAS detector is described in Sec. II. The data and simulated samples used are outlined in Sec. III, and the trigger used to collect the data is detailed in Sec. IV. The algorithms used for the displaced track and vertex reconstruction are explained in Sec. V, followed by a discussion of the selection criteria for the events and vertices in Sec. VI. The background estimation procedure is described in Sec. VII, and the systematic uncertainties are summarized in Sec. VIII. Finally, the results and conclusion are presented in Secs. IX and X.

II. ATLAS DETECTOR

The ATLAS detector [36–38] at the LHC is a cylindrical multipurpose particle detector with forward–backward symmetry and nearly 4π solid angle coverage. The detector is composed of the inner tracking detector, the electromagnetic and hadronic calorimeters, and the muon spectrometer.

The ID covers the range $0.03 \text{ m} < r < 1.1 \text{ m}$ and $|z| < 3.5 \text{ m}$,² and is immersed in a 2 T axial magnetic field from a superconducting solenoid. Three ID subdetectors provide precision tracking for charged particles within the pseudorapidity region $|\eta| < 2.5$. At small radii, a silicon pixel subdetector provides high-resolution position measurements. The pixel system consists of four barrel layers positioned at radii of 33.3 mm, 50.5 mm, 88.5 mm, and 122.5 mm, and three forward disks in each end cap. The pixel detector is surrounded by the silicon microstrip tracker (SCT), which consists of four double layers in the barrel and nine forward disks in each end cap. The radial position of the first (last) SCT layer is 299 mm (514 mm). The outermost subdetector of the ID is the straw-tube transition radiation tracker (TRT), which provides an average of 30 additional two-dimensional points to tracks in the range $|\eta| < 2.0$.

The calorimeter system provides coverage over the pseudorapidity range $|\eta| < 4.9$. It consists of an electromagnetic calorimeter surrounded by a hadronic calorimeter. Within the region $|\eta| < 3.2$, the electromagnetic calorimeter consists of liquid-argon (LAr) barrel and end cap sampling calorimeters with lead absorbers. The hadronic calorimeter consists of steel/scintillator tile calorimeters within $|\eta| < 1.7$, and two copper/LAr hadronic end cap calorimeters that cover the region $1.5 < |\eta| < 3.2$. A forward calorimeter using copper and tungsten absorbers

²ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point in the center of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the center of the LHC ring, and the y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. Pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$.

with LAr completes the calorimeter coverage up to $|\eta| = 4.9$.

The MS consists of separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by superconducting air-core toroids. The muon tracking chamber system covers the region $|\eta| < 2.7$ with three layers of monitored drift tubes (MDT), complemented by cathode strip chambers (CSC) in the forward region. The MDT chambers consist of two multilayers, each of which consists of three or four layers of drift tubes. Three stations of resistive plate chambers (RPC) and thin gap chambers (TGC) are used for triggering and ϕ measurements in the MS barrel and end caps, respectively. The muon trigger system covers the range $|\eta| < 2.4$.

The ATLAS detector selects events using a two-tiered trigger system [39]. The first level (L1), which is a hardware-based system, uses coarse data collected from the calorimeters and muon detectors to reduce the event rate from the LHC crossing frequency of 40 MHz to a design value of 100 kHz. The second level, known as the high-level trigger (HLT), is a software-based system that uses information from all of the ATLAS subdetectors to reduce the rate of recorded events to approximately 1 kHz.

III. DATA AND SIMULATED EVENTS

A. Data events

This analysis uses 33.0 fb^{-1} of data collected by the ATLAS detector during the 2016 data-taking period using pp collisions at $\sqrt{s} = 13 \text{ TeV}$. The analysis is performed using selected subsets of data which underwent special reconstruction of displaced tracks and vertices.

Two sets of data are used. One set consists of events that pass a signature-driven LLP trigger, referred to here as the Muon RoI Cluster trigger (described in Sec. IV). The Muon RoI Cluster trigger is used to collect events for the signal region; these events are also required to contain reconstructed displaced vertices in the MS and the ID, described in Secs. V and VI. Events collected by the Muon RoI Cluster trigger are also used to define a validation region for the background estimation, as discussed in Sec. VII, in which case the event selection is agnostic to the presence of any reconstructed vertices in the MS.

The second set of data is used for the background estimation. Events are selected by a single-muon trigger that requires a muon with transverse momentum $p_T > 26 \text{ GeV}$ at the HLT. To reduce the signal contamination, these events are also required to contain isolated muons. This requirement is further described in Sec. VII.

All of the events that pass the triggers used by the CR analysis [16], which select events with displaced hadronic jets decaying in the hadronic calorimeter, are vetoed in this analysis. This veto is imposed in order to facilitate the combination of the results of this search with the results of the CR and MS analyses (presented in Sec. IX).

B. Simulated events

The signal Monte Carlo (MC) samples were generated with a Φ mediator connecting the SM to a hidden sector in which Φ decays into pairs of long-lived neutral scalars ss . The neutral scalars s decay back into the SM via their coupling to the mediator, assuming a Yukawa coupling between the s -boson and the SM particles. The $\Phi \rightarrow ss$ signal samples were generated using MadGraph5 [40]. Events were showered with PYTHIA 8.210 [41] using the A14 set of tuned parameters (tune) [42] and the NNPDF2.3LO set [43] of parton distribution functions (PDF). Various mass points were generated corresponding to different combinations of the mass of the Φ (m_Φ) and m_s , with $m_\Phi \in [125, 1000] \text{ GeV}$ and $m_s \in [8, 400] \text{ GeV}$. The sensitivity of the analysis to models with s -boson mass smaller than 8 GeV is limited by the ID vertex reconstruction efficiency and selections to discriminate against vertices from background.

The LLP proper lifetime in each sample was tuned so that each mass point had an approximate mean lab-frame decay length of 5 m. The mean lab-frame decay length of 5 m is used so that there are approximately equal numbers of LLP decays in the ID and in the MS, and samples with this lab-frame decay length are used for the determination of the signal versus background selection for the IDVx, as well as the IDVx and overall efficiency studies. The overall efficiency for a range of proper lifetimes is estimated by reweighting the 5 m samples using an extrapolation method, as noted in Sec. IX. The LLP proper lifetimes for each mass point were also tuned to provide a set of samples with a mean lab-frame decay length of 9 m, and these samples are used to confirm the accuracy of the extrapolation method.

Multijet samples generated with PYTHIA 8.186 are used to determine the systematic uncertainties (described in Sec. VIII) in the displaced tracking and vertex reconstruction in the ID (described in Sec. V). The A14 tune was used together with the NNPDF2.3LO PDF set.

The generated events for all MC samples described above were processed through a full simulation of the ATLAS detector geometry and response [44] using the GEANT4 [45] toolkit. To model the effect of multiple pp interactions per bunch crossing (pileup), additional simulated pp interactions were overlaid onto each simulated hard-scatter event. Pileup was simulated with PYTHIA 8.186 using the A2 set of tuned parameters and the MSTW2008LO [46] PDF set. Per-event weights are applied to all simulated events such that the mean number of interactions per bunch crossing in simulation matches that in the data.

IV. TRIGGER

Signal region events are selected with the Muon RoI Cluster trigger, which was developed to identify events

with displaced hadronic decays outside the last active layer in the hadronic calorimeter [30].

Hadronic decays after the end of the hadronic calorimeter and before the first trigger plane in the MS are characterized by multiple muon regions of interest³ (RoIs) around the LLP line of flight. The trigger is seeded by an L1 trigger that searches for two RoIs in the MS, each of which is consistent with a particle with $p_T > 10$ GeV. At the HLT, the trigger requires clusters of muon RoIs, in which a cluster is defined as a $\Delta R = 0.4$ region containing at least three (four) muon RoIs in the barrel (end cap) of the MS.

To correct for the differences in efficiency of the trigger on events in data compared with events in MC samples, scale factors are used and are determined to be 1.13 ± 0.01 for the barrel and 1.04 ± 0.02 for the end caps [17]. The difference in the scale factors between the barrel and the end caps derives from the differences in the trigger chambers used in the barrel (RPC) and the end caps (TGC).

V. RECONSTRUCTION

To reconstruct the decay products and vertices of LLPs, dedicated reconstruction algorithms are used in both the ID and the MS. In the ID, large-radius tracking (LRT) [31] is employed after standard tracking is completed in order to reconstruct those tracks that do not point to the IP, and a displaced vertex reconstruction algorithm [28,32] draws on the combined collection of standard and large-radius tracks to form displaced ID vertices.

In the MS, LLPs that decay hadronically after the last layer of the hadronic calorimeter are likely to produce narrow, high-multiplicity jets; several times as many hits are expected to be associated with the LLP decays compared with those associated with a muon. The standard algorithms in the MS are not optimized to operate in such dense environments. A special vertex reconstruction algorithm is employed for the reconstruction of MS vertices, which is described in Ref. [33], and previously used in Refs. [17,21].

A. Reconstruction of standard jets

Jets are reconstructed in the electromagnetic and hadronic calorimeters from energy deposits in neighboring calorimeter cells. Three-dimensional topological clusters of the cells containing energy significantly above a noise threshold [47,48] are used as input to the anti- k_r jet algorithm [49]. Jets are reconstructed with an $R = 0.4$ radius parameter using the FastJet 2.4.3 [50] software package. Jet energies are calibrated using the procedure described in Ref. [47].

³In the MS, L1 muon RoIs are $\Delta\eta \times \Delta\phi = 0.2 \times 0.2$ ($\Delta\eta \times \Delta\phi = 0.1 \times 0.1$) regions in the barrel (end caps) identified as containing an interesting feature.

TABLE I. Track parameter requirements for inside-out standard and large-radius tracks.

Track parameter	Standard	Large radius
Maximum $ d_0 $	10 mm	300 mm
Maximum $ z_0 $	250 mm	1500 mm
Minimum p_T	400 MeV	500 MeV
Maximum track $ \eta $	2.7	5.0
Minimum silicon hits	7	7
Minimum unshared silicon hits	6	5

B. Reconstruction of standard tracks

Tracks in the ID are reconstructed using the energy deposits, or hits, left by charged particles. The standard ATLAS tracking algorithm reconstructs *inside-out* tracks based on seeds made of three space points⁴ in the pixel and SCT detectors [51]. A window search is performed based on the seeds, and track candidates are formed by inputting the hits in the window into a Kalman filter [52]. A track candidate must pass selection requirements on the track parameters and the constituent hits, as outlined in Table I. Track candidates are passed through an ambiguity solver [51] which evaluates the track candidates by using the hits, as well as the χ^2 of the track fit. Successful track candidates are extended into the TRT, and tracks are kept whether or not the TRT extension is successful.

After the completion of the inside-out tracking pass, an *outside-in* tracking pass is performed. Standalone TRT segments are created, seeded by energy deposits in the electromagnetic calorimeter. The standalone TRT segments are extended back into the SCT and pixel detectors, using hits that were not included in tracks reconstructed in the inside-out tracking. Standalone TRT segments that fail the extension into the silicon detectors are retained. Outside-in tracks must also pass the track impact parameter requirements in Table I.

C. Reconstruction of large-radius tracks

The standard ATLAS tracking procedure is optimized for the reconstruction of tracks that originate very close to the IP, and it has strict restrictions on the impact parameters of reconstructed tracks to reduce the reconstruction of fake tracks. Tracks produced in displaced decays often have impact parameters that are larger than the maximum impact parameter allowed by the standard tracking reconstruction algorithm. In order to reconstruct these tracks, LRT is performed, taking as inputs the hits that are left over from the standard tracking. This iteration is performed with loosened requirements on the transverse (d_0) and

⁴A space point is formed from a single measurement in the pixel detector or from a pair of measurements from the double layers in the SCT, with one measurement from a strip in the axial direction and the other from a strip in the stereo direction.

longitudinal (z_0) track impact parameters, outlined in Table I, in order to provide increased efficiency for displaced tracks.

Additionally, in order to increase the efficiency of the reconstruction of the displaced decays, the requirement on the minimum number of unshared hits is relaxed.

D. Displaced vertex reconstruction in the MS

The dedicated MSVx reconstruction algorithm makes use of the two multilayers (ML) in each MDT chamber [33]. Straight-line segments are created from at least three MDT hits in each of the MLs using a minimum χ^2 fit. The segments from the two MLs are matched to form tracklets, which are used to reconstruct the MSVx positions. The vertex reconstruction algorithm proceeds separately in the barrel and the end caps of the MS, extrapolating the tracklets through the magnetic field that is present in the barrel, and extrapolating tracklets as straight lines in the end caps where the MDT chambers are not immersed in the magnetic field. In the barrel, the χ^2 probability of the vertex is required to be $\geq 5\%$. If it is less than 5%, the tracklet with the largest contribution to the χ^2 is removed and the vertex is refit. This process is repeated until the χ^2 probability of the vertex is $\geq 5\%$ or the vertex has fewer than three tracklets. In the end cap, all tracklets in the vertex must be within 30 cm of the calculated vertex position; otherwise the farthest tracklet from the vertex is removed and the vertex is refit. In both the barrel and the end caps, each MSVx is required to have at least three tracklets.

E. Displaced vertex reconstruction in the ID

A displaced vertex reconstruction algorithm is used to reconstruct the LLP decays in the ID. The algorithm takes both the standard and large-radius tracks as input and selects tracks that meet the criteria listed in the upper section of Table II.

From this collection, a set of two-track vertices is formed from all possible pairs of intersecting tracks. The tracks making up these two-track vertices are required not to have any hits in pixel or SCT layers at smaller radii than the vertex position, and must have hits on the next possible pixel or SCT layer, unless the vertex is within a few mm of

TABLE II. Track parameter requirements for the reconstruction of displaced vertices in the ID.

Track parameter	Requirement
Track $ d_0 $	$2 \text{ mm} < d_0 < 300 \text{ mm}$
Track $ z_0 $	$< 1500 \text{ mm}$
Track p_T	$> 1 \text{ GeV}$
Number of SCT hits	≥ 2
Number of pixel and TRT hits	$n_{\text{pixel}} \geq 2$ or $n_{\text{TRT}} > 0$

No hits on track may be present before the vertex.

Hits on track must be present in the layer following the vertex.

the layer. The two-track vertex seeds are merged into multitrack vertices if they are within $d/\sigma_d < 3$ (where d is the distance between the two-track vertices and σ_d is the uncertainty in the distance). Poorly associated tracks, with track $\chi^2 > 6$, are removed from the multitrack vertices, and the vertices are refit. This process is repeated until there are no more pairs of vertices satisfying $d/\sigma_d < 3$. In the final step, all vertices within 1 mm are merged and the vertex fit is recalculated.

VI. EVENT SELECTION

All events used in the analysis are required to contain a primary vertex (PV), associated with the pp hard scatter [53]. The PV must have at least two tracks, each with $p_T > 400 \text{ MeV}$. If more than one vertex exists satisfying these criteria, the PV is chosen as the vertex with the largest sum of the squares of the p_T of all tracks associated with the vertex. The events must pass the Muon RoI Cluster trigger and the veto on the triggers from the CR analysis [16]. The events must contain a good MSVx (described in Sec. VI A) matched within $\Delta R < 0.4$ to the triggering muon cluster. Finally, events are required to have a good IDVx (described in Sec. VI B), and the MSVx and IDVx must have an angular separation of $\Delta R > 0.4$.⁵

A. MS vertex selection

The primary source of background that mimics LLP decays in the MS is jets that punch through the calorimeter. In order to reduce the background from these punch-through jets, each MSVx is required to pass certain isolation requirements developed in the MS analysis [17]. The MS vertices are required to be isolated by $\Delta R > 0.3$ ($\Delta R > 0.6$) in the barrel (end caps) from jets with $p_T > 30 \text{ GeV}$ that are matched to the PV using a jet vertex tagger discriminant [54] and have $\log_{10}(E_{\text{HAD}}/E_{\text{EM}}) < 0.5$.⁶ Isolation from these jets also reduces the contamination from multijet events. To further reduce the background from multijet events, each MSVx is required to be isolated from activity in the ID. The vector sum of the transverse momenta of tracks in a $\Delta R = 0.2$ cone around the MSVx is required to be $\Sigma p_T < 10 \text{ GeV}$, and in the barrel (end caps) the MSVx must be $\Delta R > 0.3$ ($\Delta R > 0.6$) from any tracks with $p_T > 5 \text{ GeV}$. The tracks used for this isolation must point back to the PV. Additionally, they are required to have at least seven silicon hits and no shared silicon hits, or at

⁵The separation is measured from the axis of the MSVx, which is defined with respect to the detector coordinate system, and from the (η, ϕ) of the tracks associated with the IDVx.

⁶The term $\log_{10}(E_{\text{HAD}}/E_{\text{EM}})$ quantifies the ratio of the energy deposited in the hadronic calorimeter to the energy deposited in the electromagnetic calorimeter. The purpose of this requirement is to prevent highly displaced jets from being used in the isolation because LLPs that decay at the end of the hadronic calorimeter may leave a jet and also create a decay vertex in the MS.

TABLE III. MSV_x selection requirements.

	Barrel	End caps
Precision chamber hits	$300 < n_{\text{MDT}} < 3000$	
Trigger chamber hits	$n_{\text{RPC}} > 250$	$n_{\text{TGC}} > 250$
Isolation from $p_{\text{T}} > 5$ GeV tracks	$\Delta R > 0.3$	$\Delta R > 0.6$
Σp_{T} of tracks in $\Delta R = 0.2$ cone	< 10 GeV	< 10 GeV
Isolation from $p_{\text{T}} > 30$ GeV jets	$\Delta R > 0.3$	$\Delta R > 0.6$
MSV _x $ \eta $	$ \eta_{\text{MSV}_x} < 0.7$	$1.3 < \eta_{\text{MSV}_x} < 2.5$

least ten silicon hits. To reduce the contribution from electronic noise, cosmic-ray muons, and machine-induced background, each MSV_x is also required to have a minimum number of hits associated with the vertex in the MDT (n_{MDT}), RPC (n_{RPC}), and TGC (n_{TGC}), outlined in Table III. A maximum n_{MDT} selection is applied to reduce the background from coherent noise bursts in the MDTs.

MS vertices are reconstructed by different algorithms in the barrel and the end caps. When an LLP decays in the barrel-end cap transition region, $0.8 < |\eta| < 1.3$, the resulting hits will be split between the two algorithms. This leads to a low reconstruction efficiency in the transition region since neither algorithm has access to all hits, although occasionally two separate vertices will be reconstructed from a single LLP decay. The barrel-end cap transition region overlaps very closely with the barrel-end cap transition region in the hadronic calorimeter, $0.7 < |\eta| < 1.2$, in which the probability of punch-through jets is higher. Thus, each MSV_x is required to be contained in the barrel with $|\eta_{\text{MSV}_x}| < 0.7$ or in the end cap with $|\eta_{\text{MSV}_x}| > 1.3$ to remove vertices in the barrel-end cap transition region of the MS and hadronic calorimeter. A vertex that meets all of the necessary criteria is considered to be a good MSV_x.

B. ID vertex selection

One of the primary sources of background for a search for displaced hadronic decays in the ID is vertices from interactions between particles and layers of detector material. Such hadronic interactions may result in reconstructed vertices that are indistinguishable from the reconstructed vertices of signal decays in the same region of space. In order to remove this source of background, a map is created using displaced vertices found in minimum-bias data in which the decays from known long-lived hadrons have been removed, as described in Refs. [55,56]. This map is used to create a material veto, which removes vertices in regions of space that were found to contain material. The fiducial volume defined by IDV_x, $R = \sqrt{x^2 + y^2} < 300$ mm and $|z| < 300$ mm is reduced by 42% due to this material veto [28]; however, the material veto reduces the number of background events by more than a factor of 50. Some modules in the ATLAS detector are occasionally disabled. To mimic this effect in simulation, a disabled-module veto is employed that accounts for the fact that a disabled

module could cause a track not to have a hit immediately after a vertex, causing the vertex to be rejected. This disabled-module veto operates by removing vertices in regions immediately before the location of the disabled modules. The disabled-module veto leads to a minor loss in fiducial volume of 2.3% [28] and is applied in both data and simulation.

To reject poorly reconstructed vertices resulting from random track crossings, the χ^2 value of the vertex fit divided by the number of degrees of freedom is required to be less than 5.

In addition to the IDV_x reconstruction requirement that the tracks forming the vertices have $|d_0| > 2$ mm, a minimum radial distance of 4 mm is imposed between the IDV_x and the reconstructed PV to further reduce the background contribution from b -hadrons.

ID vertices are required to have a separation of $\Delta R > 0.4$ from the nearest selected MSV_x in the event in order to reduce the probability that one high-energy hadronic jet could cause a background vertex simultaneously in the ID and in the MS.

The number of charged decay products from a hadronically decaying LLP is expected to be much higher than from vertices constructed from fake tracks or from random crossings of tracks. Thus, the number of tracks associated with the IDV_x is an important discriminant between background and signal vertices. Figure 2(a) shows the distributions of the number of tracks associated with each IDV_x (n_{trk}) for reconstructed vertices in signal MC samples compared with those reconstructed in data. In Fig. 2, the signal MC events used have no additional selection requirements in order to reduce the statistical uncertainty. The background events in data are those events in the Bkg region of the background estimation method described in Sec. VII. The reconstructed vertices in the signal MC samples are required to be $dr = \sqrt{dx^2 + dy^2 + dz^2} < 5$ mm of the LLP decay position and have at least two tracks matched⁷ to particles produced in the LLP decay. The reconstructed IDV_x distributions in the background events

⁷A reconstructed track is considered matched to a particle if it has $p > 0.5$, where p represents the weighted fraction of hits in the reconstructed track that are associated with the generated particle. The hits are assigned a weight based on the subdetector in which they were found, in order to reflect the intrinsic resolution of the measurement in that subdetector.




FIG. 2. Distributions of (a) n_{trk} and (b) m_{IDVx} for simulated signal MC and data background samples. The vertices shown are required to pass all selection requirements outlined in Table IV except the one displayed in the plot. The dashed lines and arrows indicate the selection requirement on the parameter shown. The last bin includes overflow.

in data are dominated by vertices with $n_{\text{trk}} = 2$, and when two-track vertices are removed, by $n_{\text{trk}} = 3$. For an IDVx to be considered in the signal region, it is required to have $n_{\text{trk}} \geq 4$.

Each IDVx is also required to pass a minimum vertex mass (m_{IDVx}) selection of 3 GeV, where the invariant vertex mass is computed assuming that the tracks originate from charged pions. Figure 2(b) shows the distributions of m_{IDVx} in background events and several signal samples. The selection requirement of $m_{\text{IDVx}} > 3$ GeV has the greatest impact on the selection efficiency for the lowest-mass LLPs, removing approximately half of the reconstructed ID vertices in the $m_H, m_s = [125, 8]$ GeV signal MC sample that pass the other selection requirements listed in Table IV. The heavier the mass of the LLP, the less impact the selection $m_{\text{IDVx}} > 3$ GeV has on the IDVx selection efficiency. For the $m_\phi, m_s = [1000, 400]$ GeV signal MC sample, this selection removes less than 4% of the reconstructed ID vertices that pass the other signal selections. In data, the selection on m_{IDVx} removes 70% of reconstructed ID vertices which pass all the other selections.

The requirements on the signal region IDVx are summarized in Table IV.

Figure 3 shows the IDVx selection efficiency, including the reconstruction efficiency, as a function of the decay

radius of the LLP for several mass points. The efficiency is defined as the fraction of LLP decay vertices in the fiducial volume $R, |z| < 300$ mm that are within 5 mm of reconstructed vertices that have at least two tracks matched to the decay products from the LLP and meet all the selection criteria listed in Table IV.

Figure 3(a) shows that for a fixed mediator mass the IDVx selection efficiency increases with increasing LLP mass due to the larger impact of the selection on m_{IDVx} and n_{trk} at smaller LLP masses. For a fixed LLP mass, a higher boost leads to a decreased IDVx selection efficiency [Fig. 3(b)] because decay products that point back to the IP are not included in the displaced vertex reconstruction, due to the selection of $|d_0| > 2$ mm on associated tracks.

The structure in the IDVx selection efficiency as a function of the LLP decay radius R is primarily due to the impact of the material veto, as shown in Figs. 3(a) and 3(b) for one mass point by the inclusion of the selection efficiency without the material veto. The shape versus the LLP decay radius is also impacted by other factors such as the hit requirements in the track creation and vertex reconstruction and the selection requirements on the constituent track $|d_0|$ and the distance from the PV.

C. Overall selection efficiency

Table V details the total efficiency after each selection requirement applied to signal events, as well as the relative efficiency of each subsequent selection requirement. All events in signal MC samples are found to contain a PV; thus this selection requirement is not included in the table. Many of the signal MC events generated have LLPs that decay outside of the fiducial volumes of the ID and the MS, and there is no fiducial requirement on events displayed in this table. Requiring events to contain one LLP decaying in the fiducial volume of the MS and one in the fiducial volume of the ID increases the relative efficiency of every selection requirement. It is apparent from Table V that the relative efficiency of all IDVx-related selection requirements

TABLE IV. IDVx selection requirements.

Vertex parameter	Requirement
Vertex R	< 300 mm
Vertex $ z $	< 300 mm
IDVx position	Pass material veto
IDVx position	Pass disabled-module veto
Radius from PV	> 4 mm
χ^2/n_{DoF}	< 5
ΔR from nearest good MSVx	> 0.4
m_{IDVx}	> 3 GeV
n_{trk}	≥ 4




FIG. 3. IDVx selection efficiency as a function of the radial decay position for several (a) 125 GeV $H \rightarrow ss$ and (b) $\Phi \rightarrow ss$ mass points with $m_s = 50$ GeV. The dashed vertical red lines indicate the main radial material layers in the inner detector that are included in the material veto. In each figure, the IDVx selection efficiency for one signal MC sample is shown before and after the requirement of the material veto to demonstrate the impact it has on the IDVx selection efficiency.

increases with increased LLP mass but decreases with increased LLP boost (this can also be seen in Fig. 3). Events with higher LLP mass are also more likely to pass the trigger requirements and contain a good MSVx. The boost of the LLP is associated with a higher probability to pass the trigger requirements but a lower probability to contain a good MSVx, for events that pass the trigger.

VII. BACKGROUND ESTIMATION

Sources of background for ID vertices include reconstructed vertices from the interactions of particles with detector material, vertices created from fake tracks, and vertices from random track crossings. The incidence of

vertices created from fake tracks and from random crossings is correlated with the jet activity in the events; events with a greater number of jets are more likely to include vertices from background. The background vertices are predominantly removed by the IDVx selection. A data-driven background estimation method is employed to determine the residual contribution to the events in the signal region from ID vertices from all sources of background.

The number of background events in the signal region is estimated by defining a set of background events that are designed to be approximately free of signal contamination and determining the fraction of those events that contain an IDVx that passes the full IDVx selection. This fraction is applied to events that pass the full event selection except for

TABLE V. Total and relative efficiency for each selection requirement for several signal MC mass points. Here, “Pass trigger” refers to passing the Muon RoI Cluster trigger and passing the veto on the CR triggers, “Good MSVx” includes all the MSVx selection requirements described in Table III as well as being matched to the muon RoI cluster, and “IDVx” includes all selection requirements in Table IV except those on n_{trk} and m_{IDVx} , which are listed separately. All efficiencies are computed using signal MC samples with a mean lab-frame decay length of 5 m; the proper lifetime of each mass point is listed in the table.

Selection requirements Mass point [GeV]	$c\tau$ [m]	Efficiency	Pass trigger	Good MSVx	IDVx	$n_{\text{trk}} \geq 4$	$m_{\text{IDVx}} > 3$ GeV
$m_H, m_s = [125, 8]$	0.200	Total	2.71%	1.07%	0.13%	0.005%	0.003%
		Relative	2.71%	39.3%	12.5%	3.61%	63.2%
$m_H, m_s = [125, 25]$	0.760	Total	5.13%	2.23%	0.30%	0.03%	0.02%
		Relative	5.13%	43.5%	13.3%	9.15%	81.1%
$m_H, m_s = [125, 55]$	1.540	Total	1.98%	0.75%	0.11%	0.01%	0.01%
		Relative	1.98%	37.9%	14.2%	10.1%	85.4%
$m_\Phi, m_s = [200, 50]$	1.070	Total	7.06%	3.05%	0.47%	0.07%	0.06%
		Relative	7.06%	43.2%	15.3%	15.0%	83.9%
$m_\Phi, m_s = [400, 50]$	0.700	Total	13.7%	5.02%	0.73%	0.10%	0.09%
		Relative	13.7%	36.5%	14.5%	14.3%	83.5%
$m_\Phi, m_s = [600, 50]$	0.520	Total	16.4%	4.77%	0.69%	0.08%	0.07%
		Relative	16.4%	29.0%	14.5%	12.2%	78.4%

TABLE VI. The data events used in the background estimation. These events include the background events, whose selection is defined in the text, divided into all background events (Bkg), and those that contain at least one IDVx that passes the full IDVx requirements ($Bkg + IDVx$). The other events making up the plane are the signal region events (Sig) and events that pass all signal region requirements except for the inclusion of an IDVx ($Sig - IDVx$).

	Background events	Muon RoI Cluster trigger events with a good MSVx
Has IDVx passing full signal selection	$Bkg + IDVx$	Sig
Agnostic to IDVx	Bkg	$Sig - IDVx$

the requirement of an IDVx, to estimate how many background events contain an MSVx and an IDVx.

The selection of background events is designed to limit the possibility of signal contamination. The background events (Bkg), shown in the lower left in Table VI, are required to pass the single-muon trigger described in Sec. III and are required to pass a veto on the Muon RoI Cluster trigger used to collect events in the signal region. The background events are also required to contain two isolated muons with $p_T > 25$ GeV and $p_T > 20$ GeV. When applied to events in the signal MC samples, the requirements used to define the Bkg events select fewer than 0.1% of events for any given mass point, without including the requirement of reconstructing and selecting both an MSVx and an IDVx. Accounting for preexisting limits on the branching ratio and cross section for any given mass point, the signal is expected to make up less than 0.005% of the events populating the Bkg region.

The selection for the Bkg events is agnostic to the presence of an IDVx. The $Bkg + IDVx$ events, shown in the upper left in Table VI, are required to pass the same trigger and isolated muon background requirements as the Bkg events but are also required to contain at least one IDVx that passes all of the IDVx selections outlined in Table IV. The number of events in regions $Bkg + IDVx$ and Bkg ($N_{Bkg+IDVx}$ and N_{Bkg}) are used to calculate a factor

$F = N_{Bkg+IDVx}/N_{Bkg}$, which represents the probability that a given event will contain an IDVx from background that meets the selection criteria.

Signal region events (Sig) in the upper right of Table VI are those which pass the full signal selection described in Sec. VI. Events that pass the full signal selection but are not required to contain an IDVx ($Sig - IDVx$) are in the lower right of Table VI. The number of events in the signal region (N_{Sig}) which contain an IDVx from background can be estimated using the number of $Sig - IDVx$ events ($N_{Sig-IDVx}$) and the factor F defined above. Hence, the number of Sig region events which are expected to contain an IDVx from background is estimated as $N_{Sig}^{bkg\ pred.} = N_{Sig-IDVx} \times F = N_{Sig-IDVx} \times N_{Bkg+IDVx}/N_{Bkg}$.

There are 45 events in the $Bkg + IDVx$ region out of 6 099 660 events in the Bkg region, giving a factor $F = N_{Bkg+IDVx}/N_{Bkg} = (7.4 \pm 1.1(\text{stat.})) \times 10^{-6}$. The $Sig - IDVx$ region contains 156 805 events; the predicted number of background events in the signal region is then $N_{Sig}^{pred} = N_{Sig-IDVx} \times F = N_{Sig-IDVx} \times N_{Bkg+IDVx}/N_{Bkg} = 1.16 \pm 0.18(\text{stat.})$.

The validation of the background estimation is performed using two sets of validation regions, in which the predicted and observed numbers of events are compared in regions containing vertices similar to those passing the IDVx signal selection (Table VIII). This validation serves as a cross-check that the fraction of background events that contain an IDVx is not significantly different from the fraction of events passing the Muon RoI Cluster trigger, with or without a selected MSVx, which contain an IDVx from background.

The first set of validation regions, the Bkg , $2-trk$ and Val , $2-trk$ regions, are similar to the $Bkg + IDVx$ and Sig regions except that instead of containing an IDVx that meets the full IDVx selection criteria, the selection on n_{trk} is changed from $n_{trk} \geq 4$ to $n_{trk} = 2$. Vertices with $n_{trk} = 2$ are chosen because the IDVx distribution in background events is dominated by two-track vertices, as shown in Fig. 2(a), and the signal contamination is small. This allows the fraction of vertices to be examined in events that otherwise pass all signal region selections.

TABLE VII. The events used for the validation of the background estimation, alongside the events used for the background estimate. The Bkg , $2-trk$ and Val , $2-trk$ validation regions contain ID vertices that have $n_{trk} = 2$. The $Trig$, $3-trk$ and $Trig$ validation regions are events that pass the Muon RoI Cluster trigger but are agnostic to the presence of MS vertices. The Bkg , $3-trk$ and $Trig$, $3-trk$ validation regions contain ID vertices with $1 \text{ GeV} < m_{IDVx} < 3 \text{ GeV}$ and $n_{trk} = 3$.

	Background events	Muon RoI Cluster, agnostic to MSVxs	Muon RoI Cluster, with a good MSVx
Has IDVx, $n_{trk} \geq 4$, $m_{IDVx} > 3 \text{ GeV}$	$Bkg + IDVx$		Sig
Has IDVx, $n_{trk} = 3$, $1 < m_{IDVx} < 3 \text{ GeV}$	Bkg , $3-trk$	$Trig$, $3-trk$	
Has IDVx, $n_{trk} = 2$, $m_{IDVx} > 3 \text{ GeV}$	Bkg , $2-trk$		Val , $2-trk$
Agnostic to IDVx	Bkg	$Trig$	$Sig - IDVx$

The sources of background that contribute primarily to ID vertices with $n_{\text{trk}} = 2$ are not exactly the same as those that contribute to higher n_{trk} ID vertices; thus a second set of validation regions is studied. The signal contamination is non-negligible for ID vertices with $n_{\text{trk}} > 2$ in events that pass the Muon RoI Cluster trigger and also contain an MSVx. In order to study higher n_{trk} vertices, the requirement of the MSVx is removed (*Trig* and *Trig, 3-trk* regions) in the second set of validation regions. In order to further reduce the signal contamination for ID vertices with higher n_{trk} , the requirement on m_{IDVx} is modified to select an m_{IDVx} range adjacent to that used for the final signal selection. The *Bkg, 3-trk* and *Trig, 3-trk* regions contain background events and events passing the Muon RoI Cluster trigger, respectively, that have ID vertices which pass the full signal selection, with the exceptions that the m_{IDVx} selection is changed to $1 \text{ GeV} < m_{\text{IDVx}} < 3 \text{ GeV}$ and the n_{trk} selection is changed to $n_{\text{trk}} = 3$.

The predicted and observed numbers of events in the *Val, 2-trk* and *Trig, 3-trk* regions are listed in Table VIII. The predicted and observed numbers of events in the *Val, 2-trk* region agree within 2%, and the kinematic distributions for the two-track vertices in the *Val, 2-trk* and *Bkg, 2-trk* regions are also found to be in good agreement. The predicted and observed numbers of events in the *Trig, 3-trk* region agree within 25%.

The event selection used to populate the different regions impacts the jet multiplicity in each region. The probability to find an IDVx from background is highly correlated with the jet activity in the events, and the effect is more pronounced for higher n_{trk} ID vertices than for two-track ID vertices. It is found that scaling to adjust for the differences in the jet multiplicities reduces the difference in the predicted and observed numbers of events in the *Trig, 3-trk* region. However, the impact of applying the same scaling to the predicted number of background events in the final signal region changed the final background estimate by less than the statistical uncertainty, so this scaling was not ultimately applied.

The largest difference in all background systematic uncertainty studies is observed to be within 25%, so this

TABLE VIII. Numbers used in the background estimation method and for the validation of the estimate, for the observed (n_{obs}) number of events in the *Bkg, Bkg + IDVx*, and *Sig-IDVx* regions, and the predicted (n_{pred}) and observed numbers of events in the *Val, 2-trk, Trig, 3-trk*, and *Sig* regions.

	n_{obs}	
Region <i>Bkg</i>	6 099 660	
Region <i>Bkg + IDVx</i>	45	
Region <i>Sig - IDVx</i>	156 805	
	n_{pred}	n_{obs}
Region <i>Val, 2-trk</i>	$11,269 \pm 46(\text{stat.})$	11 470
Region <i>Trig, 3-trk</i>	$1750 \pm 64(\text{stat.})$	2132
Region <i>Sig</i>	$1.16 \pm 0.18(\text{stat.}) \pm 0.29(\text{syst.})$	1

value is taken to be the systematic uncertainty of the background estimation. The sensitivity of the results to this choice is tested by doubling the systematic uncertainty on the background; the resulting change in the predicted and observed limits is negligible, as the number of predicted background events is small. With a 25% systematic uncertainty, the predicted number of background events passing the final signal selection is $1.16 \pm 0.18(\text{stat.}) \pm 0.29(\text{syst.})$.

VIII. SYSTEMATIC UNCERTAINTIES IN SIGNAL PREDICTIONS

Several sources of systematic uncertainty in the signal selection efficiency are considered. The dominant uncertainty is due to the difference in performance of the LRT and ID displaced vertex reconstruction algorithms between data and MC simulation.

A. Systematic uncertainties in large-radius tracking and displaced vertex reconstruction

To assess the systematic uncertainty of the ID vertex reconstruction efficiency due to the modeling of the large-radius tracking and vertex reconstruction, the rates of displaced vertices consistent with $K_S^0 \rightarrow \pi^+ \pi^-$ decays are compared between data and multijet simulation. The uncertainty is estimated by examining the variations between data and simulation in the K_S^0 yield as a function of vertex radius.

Events in data and MC samples are selected by requiring the presence of a PV. The events used in data are in the selected subset of events that underwent the reconstruction of displaced tracks and vertices. The same reconstruction is performed on the multijet MC sample, and the simulated events are reweighted to correctly reproduce the distribution of the mean number of interactions per bunch crossing in the data. To minimize the statistical uncertainty, no additional event-level requirements are applied. From the selected events, candidate K_S^0 vertices are identified by requiring that the vertices have a decay length greater than 15 mm, exactly two tracks, and an invariant mass in the region 450 to 550 MeV. After the last two selection criteria are applied, the vast majority of selected vertices originate from K_S^0 decays, and the possibility of signal contamination is minimized. The kinematic distributions of candidate K_S^0 vertices are compared between data and MC and are found to have good agreement within statistical uncertainties.

The number of K_S^0 candidate vertices found in data and simulation are binned by their decay radius R . To achieve a better estimate of the number of K_S^0 in each bin, the background contribution is computed from the sidebands of the invariant mass distribution, from 350 to 450 MeV and from 550 to 650 MeV, and subtracted from the number of K_S^0 candidates in the region 450 to 550 MeV.

Tracks originating from a K_S^0 decay can be reconstructed by either the standard tracking or the LRT algorithm. The




FIG. 4. CL_S upper limits at 95% CL on the branching ratio $B_{H \rightarrow ss}$ for (a) $m_H = 125$ GeV, and on the cross section times branching ratio $\sigma \times B_{\phi \rightarrow ss}$ for (b) $m_\phi = 200\text{--}400$ GeV and (c) $m_\phi = 600\text{--}1000$ GeV.

data are normalized such that the number of K_S^0 vertices with two standard tracks is the same between data and simulation. This accounts for any differences that may exist between data and simulation in the total number of K_S^0 decays. The vertex yields of K_S^0 with two large-radius tracks are compared between data and MC simulation, and the largest difference in the ratio of data to MC is found to be 20%. For vertices in the signal region, the effect of this tracking inefficiency is reduced due to the high multiplicity of tracks present in the vertices. The uncertainty in the vertex reconstruction efficiency is taken to be 20% and is applied as an uncertainty of the global selection efficiency.

B. Other systematic uncertainties

The uncertainty of the integrated luminosity measurement is 2.2% [57], obtained using the LUCID-2 detector [58] for the primary luminosity measurements.

The systematic uncertainties of the Muon RoI Cluster trigger efficiency and the MSVx reconstruction efficiency are examined in Ref. [17]. The systematic uncertainty due to the trigger scale factors is evaluated by varying the scale

factors up and down by the uncertainty of the scale factor fit and comparing the trigger efficiency resulting from the modified scale factors with the trigger efficiency from the nominal scale factor. The systematic uncertainties are evaluated separately in the barrel and the end caps. Similar methods are used to determine the impact of the pileup uncertainty and the systematic uncertainty from the PDF used to generate the signal MC events on the trigger efficiency. The largest relative uncertainty in the trigger efficiency in any given mass point used in this analysis is found to be 4.8%.

These methods are also used to evaluate the impact of the pileup and PDF uncertainties on the MSVx reconstruction efficiency, and the largest relative uncertainty for any mass point in the barrel or end cap is found to be 5.5%.

IX. RESULTS

One event is observed that passes the full signal selection, consistent with the estimated background of $1.16 \pm 0.18(\text{stat.}) \pm 0.29(\text{syst.})$ events.

Upper limits at the 95% confidence level (CL) are set on the production cross section times branching ratio, for




FIG. 5. Combined limits from this analysis (ID) and the CR and MS analyses for $m_H = 125$ GeV decaying into (a) 15, (b) 25, (c) 40, and (d) 55 GeV mass scalars. The expected limit is shown as a dashed line with shading for the $\pm 1\sigma$ error band, and the observed limit is shown with a solid line. The combination of the CR and MS analyses used both the MS1 and MS2 channels for $m_H = 125$ GeV, but due to orthogonality considerations, only the MS2 channel was used when performing the combination with the ID analysis. The MS analysis did not place limits on the 55 GeV LLP mass point (d) so the combined limits use the results of the ID and CR analyses only.

various signal mass hypotheses, following the CL_s prescription [59] with a profile likelihood ratio used as the test statistic. An asymptotic approach [60] is used to compute the CL_s value. This method was tested and found to give results consistent with those obtained from ensemble tests. A Poisson probability term describing the total number of observed events is used, and the systematic uncertainties of the signal efficiency, background estimation, and luminosity are treated as nuisance parameters and are assigned Gaussian constraints.

To evaluate the efficiency of the full event selection as a function of the proper lifetime of the LLP, a reweighting procedure is used, following the method described in Ref. [16]. The extrapolated efficiency from the 5 m mean lab-frame decay length sample agrees with the efficiency from the 9 m mean lab-frame decay length sample within the combined statistical uncertainty from the efficiency and extrapolation computations.

The observed limits for all benchmark models considered are summarized in Fig. 4. For the $m_H = 125$ GeV mediator, the SM Higgs boson gluon–gluon fusion production cross section of 49 pb [61] at 13 TeV is assumed, and limits on the branching ratio $B_{H \rightarrow ss}$ are shown. The observed limits are consistent with the expected limits within $\pm 1\sigma$ and significantly extend the limits set by previous displaced jet searches [16,17] for low scalar masses and short lifetimes. The analysis is most sensitive to τ values between 0.1 m and 3 m, where equal numbers of LLP decays are found in the fiducial volumes of the ID and the MS.

A. Combination of results with other displaced jet searches

The results presented in this paper are complementary to the CR [16] and MS [17] analyses, which set limits on the same benchmark models as used in this analysis. The




FIG. 6. Combined limits from this analysis (ID) and the CR and MS analyses for benchmark models with $m_\Phi = 200$ GeV decaying into (a) 25 and (b) 50 GeV mass scalars. The expected limit is shown as a dashed line with shading for the $\pm 1\sigma$ error band, and the observed limit is shown with a solid line.

results derived above are combined with the results from the CR and MS analyses to provide increased sensitivity over a greater range of proper lifetimes.

The orthogonality of the search presented in this paper (the ID analysis) and the CR analysis is ensured by vetoing on the hadronic calorimeter triggers in both the data and signal MC events as described in Sec. VI. The MS analysis is separated into the 1-MSVx plus missing transverse momentum (MS1) and 2-MSVx (MS2) channels. To ensure orthogonality between the ID and MS analyses, only the 2-MSVx channel is used in the combination. Across all MC signal samples in this channel, only a few events are found which contain both an IDVx and two MS vertices passing all selections used by the respective analyses, and these events are explicitly vetoed for the combination. There are no events found in data which pass the full selection of both the ID and MS2 analyses. The orthogonality between the CR and MS analyses has been verified in Ref. [16]. This ensures that the final selected MC signal events and data events in all three analyses used in the combination are statistically independent.

The combination is performed using a simultaneous fit of the likelihood functions of each analysis. The signal strength is correlated between all three likelihoods, as is the nuisance parameter for the luminosity uncertainties. The signal uncertainties are treated as uncorrelated since they are dominated by different experimental uncertainties in each search. The background estimates in each analysis are derived using independent data-driven methods and are therefore not correlated.

As in the individual searches, the asymptotic approach is used to compute the CL_s value, and the limits are defined by the region excluded at 95% CL. The limits are calculated using a global fit, where the overall likelihood function is the product of the individual likelihood functions of the searches to be combined. The limits are calculated separately at each $c\tau$ point, and at each point the signal

efficiency is scaled by the result of the lifetime extrapolation procedure.

Figures 5 and 6 show the observed and expected limits for the ID analysis, as well as the combination of the CR and MS analyses both with and without their combination with the ID analysis. For the models with $m_H = 125$ GeV or $m_\Phi = 200$ GeV and $8 \text{ GeV} < m_s \leq 55 \text{ GeV}$, the ID analysis has greater sensitivity than the combination of the CR and MS analyses for proper lifetimes ranging from 0.05 m up to 0.7 m. Although the IDVx reconstruction efficiency diminishes with decreasing LLP masses, requiring both a good MSVx and a good IDVx suppresses the background in the final signal region and allows stronger limits to be set at low $c\tau$ for these mass points.

Table IX summarizes the proper lifetime ranges excluded by the combinations for the $m_H = 125$ GeV mediator, assuming a 10% branching ratio for $H \rightarrow ss$ and using the SM Higgs boson gluon–gluon fusion production cross section.

The combination of the results was also explored for signal points with $m_\Phi \geq 400$ GeV. The sensitivity of this analysis increases as the Φ - and s -boson masses increase, due to the increased fraction of reconstructed ID vertices passing the IDVx selection requirements, as shown in Fig. 2.

TABLE IX. Ranges of proper lifetimes excluded at 95% CL for the $m_H = 125$ GeV benchmark model assuming a 10% branching ratio for $H \rightarrow ss$. The $m_s = 55$ GeV exclusion range uses the results of the ID and CR analyses only.

m_s [GeV]	Excluded $c\tau$ range at 95% CL [m]
8	0.06–15
15	0.09–64
25	0.12–116
40	0.26–197
55	0.39–8.1

However, the gain in sensitivity with increasing masses is much larger for the CR analysis. Thus, the addition of the results from this analysis does not noticeably extend the combined exclusion limits for signal points with $m_\Phi \geq 400$ GeV, and they are therefore not shown in this paper.

X. CONCLUSION

This paper presents a search for pairs of long-lived particles decaying in the ATLAS inner tracking detector and muon spectrometer, using 33.0 fb^{-1} of $\sqrt{s} = 13$ TeV proton–proton collision data which were collected at the LHC in 2016. Benchmark HS models are studied, using a scalar mediator that ranges in mass from 125 to 1000 GeV, decaying into pairs of long-lived scalars ranging in mass from 8 to 400 GeV, depending on the mass of the mediator. The search presented focuses on the topology consisting of one displaced hadronic decay in the inner detector and one in the muon spectrometer. The search employs dedicated techniques to reconstruct both the displaced inner detector and muon spectrometer hadronic vertices. A data-driven background estimation is performed, which predicts approximately one event in the signal region from background sources. One event is found in the signal region, and limits are set on the various signal mass points. This search has a greater sensitivity for low-mass, long-lived scalars at shorter lifetimes than previously published searches. The limits resulting from the combination of this search with the previous ATLAS searches for long-lived particles are the most stringent thus far on the branching ratios from the Higgs boson to several low-mass scalars, and on the cross section times branching ratio for a 200 GeV mass Φ decaying into long-lived scalars with masses of 25 and 50 GeV.

ACKNOWLEDGMENTS

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently. We acknowledge the support of ANPCyT, Argentina; YerPhI,

Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; SSTC, Belarus; CNPq and FAPESP, Brazil; NSERC, NRC, and CFI, Canada; CERN; CONICYT, Chile; CAS, MOST, and NSFC, China; COLCIENCIAS, Colombia; MSMT CR, MPO CR, and VSC CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF, and MPG, Germany; GSRT, Greece; RGC and Hong Kong SAR, China; ISF and Benozio Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW and NCN, Poland; FCT, Portugal; MNE/IFA, Romania; MES of Russia and NRC KI, Russia Federation; JINR; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DST/NRF, South Africa; MINECO, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF, and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TAEK, Turkey; STFC, United Kingdom; DOE and NSF, United States. In addition, individual groups and members have received support from BCKDF, CANARIE, Compute Canada, and CRC, Canada; ERC, ERDF, Horizon 2020, Marie Skłodowska-Curie Actions, and COST, European Union; Investissements d’Avenir Labex, Investissements d’Avenir IDEX, and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales, and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and GIF, Israel; CERCA Programme Generalitat de Catalunya and PROMETEO Programme Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; and The Royal Society and Leverhulme Trust, United Kingdom. The crucial computing support from all WLCG partners is gratefully acknowledged, in particular, from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide, and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [62].

-
- [1] G.F. Giudice and R. Rattazzi, Theories with gauge-mediated supersymmetry breaking, *Phys. Rep.* **322**, 419 (1999).
 - [2] R. Barbier *et al.*, R-parity-violating supersymmetry, *Phys. Rep.* **420**, 1 (2005).
 - [3] J. Fan, M. Reece, and J. T. Ruderman, A stealth supersymmetry sampler, *J. High Energy Phys.* **07** (2012) 196.
 - [4] C. Csáki, E. Kuflik, O. Slone, and T. Volansky, Models of dynamical R-parity violation, *J. High Energy Phys.* **06** (2015) 045.
 - [5] M. J. Strassler and K. Zurek, Echoes of a hidden valley at hadron colliders, *Phys. Lett. B* **651**, 374 (2007).
 - [6] M. J. Strassler and K. M. Zurek, Discovering the Higgs through highly-displaced vertices, *Phys. Lett. B* **661**, 263 (2008).
 - [7] T. Han, Z. Si, K. Zurek, and M. J. Strassler, Phenomenology of hidden valleys at hadron colliders, *J. High Energy Phys.* **07** (2008) 008.
 - [8] Z. Chacko, H.-S. Goh, and R. Harnik, Natural Electroweak Breaking from a Mirror Symmetry, *Phys. Rev. Lett.* **96**, 231802 (2006).

- [9] H. Cai, H.-C. Cheng, and J. Terning, A quirky little Higgs model, *J. High Energy Phys.* **05** (2009) 045.
- [10] Z. Chacko, D. Curtin, and C.B. Verhaaren, A quirky probe of neutral naturalness, *Phys. Rev. D* **94**, 011504 (2016).
- [11] ATLAS Collaboration, Combined measurements of Higgs boson production and decay using up to 80 fb⁻¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS experiment, *Phys. Rev. D* **101**, 012002 (2020).
- [12] ATLAS Collaboration, Constraints on new phenomena via Higgs boson couplings and invisible decays with the ATLAS detector, *J. High Energy Phys.* **11** (2015) 206.
- [13] M.J. Strassler, Possible effects of a hidden valley on supersymmetric phenomenology, [arXiv:hep-ph/0607160](https://arxiv.org/abs/hep-ph/0607160).
- [14] D. Curtin and C.B. Verhaaren, Discovering uncolored naturalness in exotic Higgs decays, *J. High Energy Phys.* **12** (2015) 072.
- [15] D. Curtin *et al.*, Exotic decays of the 125 GeV Higgs boson, *Phys. Rev. D* **90**, 075004 (2014).
- [16] ATLAS Collaboration, Search for long-lived neutral particles in pp collisions at $\sqrt{s} = 13$ TeV that decay into displaced hadronic jets in the ATLAS calorimeter, *Eur. Phys. J. C* **79**, 481 (2019).
- [17] ATLAS Collaboration, Search for long-lived particles produced in pp collisions at $\sqrt{s} = 13$ TeV that decay into displaced hadronic jets in the ATLAS muon spectrometer, *Phys. Rev. D* **99**, 052005 (2019).
- [18] K. Jedamzik, Big bang nucleosynthesis constraints on hadronically and electromagnetically decaying relic neutral particles, *Phys. Rev. D* **74**, 103509 (2006).
- [19] ATLAS Collaboration, Search for a Light Higgs Boson Decaying to Long-Lived Weakly Interacting Particles in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector, *Phys. Rev. Lett.* **108**, 251801 (2012).
- [20] LHCb Collaboration, Search for long-lived particles decaying to jet pairs, *Eur. Phys. J. C* **75**, 152 (2015).
- [21] ATLAS Collaboration, Search for long-lived, weakly interacting particles that decay to displaced hadronic jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *Phys. Rev. D* **92**, 012010 (2015).
- [22] ATLAS Collaboration, Search for massive, long-lived particles using multitrack displaced vertices or displaced lepton pairs in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *Phys. Rev. D* **92**, 072004 (2015).
- [23] CMS Collaboration, Search for long-lived neutral particles decaying to quark-antiquark pairs in proton-proton collisions at $\sqrt{s} = 8$ TeV, *Phys. Rev. D* **91**, 012007 (2015).
- [24] ATLAS Collaboration, Search for pair-produced long-lived neutral particles decaying to jets in the ATLAS hadronic calorimeter in pp collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett. B* **743**, 15 (2015).
- [25] LHCb Collaboration, Updated search for long-lived particles decaying to jet pairs, *Eur. Phys. J. C* **77**, 812 (2017).
- [26] LHCb Collaboration, Search for massive long-lived particles decaying semileptonically in the LHCb detector, *Eur. Phys. J. C* **77**, 224 (2017).
- [27] CMS Collaboration, Search for new long-lived particles at $\sqrt{s} = 13$ TeV, *Phys. Lett. B* **780**, 432 (2018).
- [28] ATLAS Collaboration, Search for long-lived, massive particles in events with displaced vertices and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector, *Phys. Rev. D* **97**, 052012 (2018).
- [29] ATLAS Collaboration, Search for long-lived particles in final states with displaced dimuon vertices in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. D* **99**, 012001 (2019).
- [30] ATLAS Collaboration, Triggers for displaced decays of long-lived neutral particles in the ATLAS detector, *J. Instrum.* **8**, P07015 (2013).
- [31] ATLAS Collaboration, Performance of the reconstruction of large impact parameter tracks in the inner detector of ATLAS, CERN Report No. ATL-PHYS-PUB-2017-014, 2017, <https://cds.cern.ch/record/2275635>.
- [32] ATLAS Collaboration, A study of the material in the ATLAS inner detector using secondary hadronic interactions, *J. Instrum.* **7**, P01013 (2012).
- [33] ATLAS Collaboration, Standalone vertex finding in the ATLAS muon spectrometer, *J. Instrum.* **9**, P02001 (2014).
- [34] R. Bruce *et al.*, Sources of machine-induced background in the ATLAS and CMS detectors at the CERN Large Hadron Collider, *Nucl. Instrum. Methods Phys. Res., Sect. A* **729**, 825 (2013).
- [35] ATLAS Collaboration, Comparison between simulated and observed LHC beam backgrounds in the ATLAS experiment at $E_{\text{beam}} = 4$ TeV, *J. Instrum.* **13**, P12006 (2018).
- [36] ATLAS Collaboration, The ATLAS Experiment at the CERN Large Hadron Collider, *J. Instrum.* **3**, S08003 (2008).
- [37] ATLAS Collaboration, ATLAS insertable B-layer technical design report, CERN Report No. ATLAS-TDR-19, 2010, <https://cds.cern.ch/record/1291633>; ATLAS insertable B-layer technical design report Addendum, CERN Report No. ATLAS-TDR-19-ADD-1, 2012, <https://cds.cern.ch/record/1451888>.
- [38] B. Abbott *et al.*, Production and integration of the ATLAS insertable B-layer, *J. Instrum.* **13**, T05008 (2018).
- [39] ATLAS Collaboration, Performance of the ATLAS trigger system in 2015, *Eur. Phys. J. C* **77**, 317 (2017).
- [40] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *J. High Energy Phys.* **07** (2014) 079.
- [41] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, An introduction to PYTHIA 8.2, *Comput. Phys. Commun.* **191**, 159 (2015).
- [42] ATLAS Collaboration, ATLAS PYTHIA 8 tunes to 7 TeV data, CERN Report No. ATL-PHYS-PUB-2014-021, 2014, <https://cds.cern.ch/record/1966419>.
- [43] R. D. Ball *et al.*, Parton distributions with LHC data, *Nucl. Phys.* **B867**, 244 (2013).
- [44] ATLAS Collaboration, The ATLAS simulation infrastructure, *Eur. Phys. J. C* **70**, 823 (2010).
- [45] S. Agostinelli *et al.*, GEANT4—A simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).

- [46] R. D. Ball *et al.*, Parton distributions for the LHC Run II, *J. High Energy Phys.* **04** (2015) 040.
- [47] ATLAS Collaboration, Jet energy scale measurements and their systematic uncertainties in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. D* **96**, 072002 (2017).
- [48] ATLAS Collaboration, Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1, *Eur. Phys. J. C* **77**, 490 (2017).
- [49] M. Cacciari, G. P. Salam, and G. Soyez, The anti- k_r jet clustering algorithm, *J. High Energy Phys.* **04** (2008) 063.
- [50] M. Cacciari, G. P. Salam, and G. Soyez, Fastjet user manual, *Eur. Phys. J. C* **72**, 1896 (2012).
- [51] ATLAS Collaboration, Performance of the ATLAS silicon pattern recognition algorithm in data and simulation at $\sqrt{s} = 7$ TeV, Report No. ATLAS-CONF-2010-072, 2010, <https://cds.cern.ch/record/1281363>.
- [52] R. Frühwirth, Application of Kalman filtering to track and vertex fitting, *Nucl. Instrum. Methods Phys. Res., Sect. A* **262**, 444 (1987).
- [53] ATLAS Collaboration, Vertex reconstruction performance of the ATLAS detector at $\sqrt{s} = 13$ TeV, CERN Report No. ATL-PHYS-PUB-2015-026, 2015, <https://cds.cern.ch/record/2037717>.
- [54] ATLAS Collaboration, Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector, *Eur. Phys. J. C* **76**, 581 (2016).
- [55] ATLAS Collaboration, A measurement of material in the ATLAS tracker using secondary hadronic interactions in 7 TeV pp collisions, *J. Instrum.* **11**, P11020 (2016).
- [56] ATLAS Collaboration, Study of the material of the ATLAS inner detector for Run 2 of the LHC, *J. Instrum.* **12**, P12009 (2017).
- [57] ATLAS Collaboration, Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC, CERN Tech. Reports No. ATLAS-CONF-2019-021 and No. CERN, 2019, <https://cds.cern.ch/record/2677054>.
- [58] G. Avoni *et al.*, The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS, *J. Instrum.* **13**, P07017 (2018).
- [59] A. L. Read, Presentation of search results: The CLS technique, *J. Phys. G* **28**, 2693 (2002).
- [60] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, *Eur. Phys. J. C* **71**, 1554 (2011); **73**, 2501(E) (2013).
- [61] D. de Florian *et al.*, *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, CERN Yellow Reports: Monographs Vol. 2 (CERN, Geneva, Switzerland, 2017).
- [62] ATLAS Collaboration, ATLAS computing acknowledgements, CERN Report No. ATL-GEN-PUB-2016-002, <https://cds.cern.ch/record/2202407>.

G. Aad,¹⁰² B. Abbott,¹²⁹ D. C. Abbott,¹⁰³ A. Abed Abud,³⁶ K. Abeling,⁵³ D. K. Abhayasinghe,⁹⁴ S. H. Abidi,¹⁶⁷ O. S. AbouZeid,⁴⁰ N. L. Abraham,¹⁵⁶ H. Abramowicz,¹⁶¹ H. Abreu,¹⁶⁰ Y. Abulaiti,⁶ B. S. Acharya,^{67a,67b,b} B. Achkar,⁵³ S. Adachi,¹⁶³ L. Adam,¹⁰⁰ C. Adam Bourdarios,⁵ L. Adamczyk,^{84a} L. Adamek,¹⁶⁷ J. Adelman,¹²¹ M. Adersberger,¹¹⁴ A. Adiguzel,^{12c} S. Adorni,⁵⁴ T. Adye,¹⁴⁴ A. A. Affolder,¹⁴⁶ Y. Afik,¹⁶⁰ C. Agapopoulou,⁶⁵ M. N. Agaras,³⁸ A. Aggarwal,¹¹⁹ C. Agheorghiesei,^{27c} J. A. Aguilar-Saavedra,^{140f,140a,c} F. Ahmadov,⁸⁰ W. S. Ahmed,¹⁰⁴ X. Ai,¹⁸ G. Aielli,^{74a,74b} S. Akatsuka,⁸⁶ T. P. A. Åkesson,⁹⁷ E. Akilli,⁵⁴ A. V. Akimov,¹¹¹ K. Al Khoury,⁶⁵ G. L. Alberghi,^{23b,23a} J. Albert,¹⁷⁶ M. J. Alconada Verzini,¹⁶¹ S. Alderweireldt,³⁶ M. Aleksa,³⁶ I. N. Aleksandrov,⁸⁰ C. Alexa,^{27b} T. Alexopoulos,¹⁰ A. Alfonsi,¹²⁰ F. Alfonsi,^{23b,23a} M. Alhroob,¹²⁹ B. Ali,¹⁴² M. Aliev,¹⁶⁶ G. Alimonti,^{69a} S. P. Alkire,¹⁴⁸ C. Allaire,⁶⁵ B. M. M. Allbrooke,¹⁵⁶ B. W. Allen,¹³² P. P. Allport,²¹ A. Aloisio,^{70a,70b} F. Alonso,⁸⁹ C. Alpigiani,¹⁴⁸ A. A. Alshehri,⁵⁷ M. Alvarez Estevez,⁹⁹ M. G. Alviggi,^{70a,70b} Y. Amaral Coutinho,^{81b} A. Ambler,¹⁰⁴ L. Ambroz,¹³⁵ C. Amelung,²⁶ D. Amidei,¹⁰⁶ S. P. Amor Dos Santos,^{140a} S. Amoroso,⁴⁶ C. S. Amrouche,⁵⁴ F. An,⁷⁹ C. Anastopoulos,¹⁴⁹ N. Andari,¹⁴⁵ T. Andeen,¹¹ C. F. Anders,^{61b} J. K. Anders,²⁰ A. Andreazza,^{69a,69b} V. Andrei,^{61a} C. R. Anelli,¹⁷⁶ S. Angelidakis,³⁸ A. Angerami,³⁹ A. V. Anisenkov,^{122b,122a} A. Annovi,^{72a} C. Antel,⁵⁴ M. T. Anthony,¹⁴⁹ E. Antipov,¹³⁰ M. Antonelli,⁵¹ D. J. A. Antrim,¹⁷¹ F. Anulli,^{73a} M. Aoki,⁸² J. A. Aparisi Pozo,¹⁷⁴ L. Aperio Bella,^{15a} J. P. Araque,^{140a} V. Araujo Ferraz,^{81b} R. Araujo Pereira,^{81b} C. Arcangeletti,⁵¹ A. T. H. Arce,⁴⁹ F. A. Arduh,⁸⁹ J-F. Arguin,¹¹⁰ S. Argyropoulos,⁷⁸ J.-H. Arling,⁴⁶ A. J. Armbruster,³⁶ A. Armstrong,¹⁷¹ O. Arnaez,¹⁶⁷ H. Arnold,¹²⁰ Z. P. Arrubarrena Tame,¹¹⁴ G. Artoni,¹³⁵ S. Artz,¹⁰⁰ S. Asai,¹⁶³ T. Asawatavonvanich,¹⁶⁵ N. Asbah,⁵⁹ E. M. Asimakopoulou,¹⁷² L. Asquith,¹⁵⁶ J. Assahsah,^{35d} K. Assamagan,²⁹ R. Astalos,^{28a} R. J. Atkin,^{33a} M. Atkinson,¹⁷³ N. B. Atlay,¹⁹ H. Atmani,⁶⁵ K. Augsten,¹⁴² G. Avolio,³⁶ R. Avramidou,^{60a} M. K. Ayoub,^{15a} A. M. Azoulay,^{168b} G. Azuelos,^{110,d} H. Bachacou,¹⁴⁵ K. Bachas,^{68a,68b} M. Backes,¹³⁵ F. Backman,^{45a,45b} P. Bagnaia,^{73a,73b} M. Bahmani,⁸⁵ H. Bahrasemani,¹⁵² A. J. Bailey,¹⁷⁴ V. R. Bailey,¹⁷³ J. T. Baines,¹⁴⁴ C. Bakalis,¹⁰ O. K. Baker,¹⁸³ P. J. Bakker,¹²⁰ D. Bakshi Gupta,⁸ S. Balaji,¹⁵⁷ E. M. Baldin,^{122b,122a} P. Balek,¹⁸⁰ F. Balli,¹⁴⁵ W. K. Balunas,¹³⁵ J. Balz,¹⁰⁰ E. Banas,⁸⁵ A. Bandyopadhyay,²⁴ Sw. Banerjee,^{181,e} A. A. E. Bannoura,¹⁸² L. Barak,¹⁶¹ W. M. Barbe,³⁸ E. L. Barberio,¹⁰⁵ D. Barberis,^{55b,55a} M. Barbero,¹⁰² G. Barbour,⁹⁵ T. Barillari,¹¹⁵ M-S. Barisits,³⁶ J. Barkeloo,¹³² T. Barklow,¹⁵³ R. Barnea,¹⁶⁰ S. L. Barnes,^{60c} B. M. Barnett,¹⁴⁴ R. M. Barnett,¹⁸ Z. Barnovska-Blenessy,^{60a}

A. Baroncelli,^{60a} G. Barone,²⁹ A. J. Barr,¹³⁵ L. Barranco Navarro,^{45a,45b} F. Barreiro,⁹⁹ J. Barreiro Guimarães da Costa,^{15a} S. Barsov,¹³⁸ R. Bartoldus,¹⁵³ G. Bartolini,¹⁰² A. E. Barton,⁹⁰ P. Bartos,^{28a} A. Basalaeu,⁴⁶ A. Basan,¹⁰⁰ A. Bassalat,^{65,f} M. J. Basso,¹⁶⁷ R. L. Bates,⁵⁷ S. Batlamous,^{35e} J. R. Batley,³² B. Batool,¹⁵¹ M. Battaglia,¹⁴⁶ M. Bauce,^{73a,73b} F. Bauer,¹⁴⁵ K. T. Bauer,¹⁷¹ H. S. Bawa,³¹ J. B. Beacham,⁴⁹ T. Beau,¹³⁶ P. H. Beauchemin,¹⁷⁰ F. Becherer,⁵² P. Bechtle,²⁴ H. C. Beck,⁵³ H. P. Beck,^{20,g} K. Becker,¹⁷⁸ C. Becot,⁴⁶ A. Beddall,^{12d} A. J. Beddall,^{12a} V. A. Bednyakov,⁸⁰ M. Bedognetti,¹²⁰ C. P. Bee,¹⁵⁵ T. A. Beermann,¹⁸² M. Begalli,^{81b} M. Begel,²⁹ A. Behera,¹⁵⁵ J. K. Behr,⁴⁶ F. Beisiegel,²⁴ A. S. Bell,⁹⁵ G. Bella,¹⁶¹ L. Bellagamba,^{23b} A. Bellerive,³⁴ P. Bellos,⁹ K. Beloborodov,^{122b,122a} K. Belotskiy,¹¹² N. L. Belyaev,¹¹² D. Bencheekroun,^{35a} N. Benekos,¹⁰ Y. Benhammou,¹⁶¹ D. P. Benjamin,⁶ M. Benoit,⁵⁴ J. R. Bensinger,²⁶ S. Bentvelsen,¹²⁰ L. Beresford,¹³⁵ M. Beretta,⁵¹ D. Berge,⁴⁶ E. Bergeaas Kuutmann,¹⁷² N. Berger,⁵ B. Bergmann,¹⁴² L. J. Bergsten,²⁶ J. Beringer,¹⁸ S. Berlendis,⁷ G. Bernardi,¹³⁶ C. Bernius,¹⁵³ F. U. Bernlochner,²⁴ T. Berry,⁹⁴ P. Berta,¹⁰⁰ C. Bertella,^{15a} I. A. Bertram,⁹⁰ O. Bessidskaia Bylund,¹⁸² N. Besson,¹⁴⁵ A. Bethani,¹⁰¹ S. Bethke,¹¹⁵ A. Betti,⁴² A. J. Bevan,⁹³ J. Beyer,¹¹⁵ D. S. Bhattacharya,¹⁷⁷ P. Bhattarai,²⁶ R. Bi,¹³⁹ R. M. Bianchi,¹³⁹ O. Biebel,¹¹⁴ D. Biedermann,¹⁹ R. Bielski,³⁶ K. Bierwagen,¹⁰⁰ N. V. Biesuz,^{72a,72b} M. Biglietti,^{75a} T. R. V. Billoud,¹¹⁰ M. Bindi,⁵³ A. Bingul,^{12d} C. Bini,^{73a,73b} S. Biondi,^{23b,23a} M. Birman,¹⁸⁰ T. Bisanz,⁵³ J. P. Biswal,¹⁶¹ D. Biswas,^{181,e} A. Bitadze,¹⁰¹ C. Bittrich,⁴⁸ K. Bjørke,¹³⁴ T. Blazek,^{28a} I. Bloch,⁴⁶ C. Blocker,²⁶ A. Blue,⁵⁷ U. Blumenschein,⁹³ G. J. Bobbink,¹²⁰ V. S. Bobrovnikov,^{122b,122a} S. S. Bocchetta,⁹⁷ A. Bocci,⁴⁹ D. Boerner,⁴⁶ D. Bogavac,¹⁴ A. G. Bogdanchikov,^{122b,122a} C. Bohm,^{45a} V. Boisvert,⁹⁴ P. Bokan,^{53,172} T. Bold,^{84a} A. S. Boldyrev,¹¹³ A. E. Bolz,^{61b} M. Bomben,¹³⁶ M. Bona,⁹³ J. S. Bonilla,¹³² M. Boonekamp,¹⁴⁵ C. D. Booth,⁹⁴ H. M. Borecka-Bielska,⁹¹ A. Borisov,¹²³ G. Borissov,⁹⁰ J. Bortfeldt,³⁶ D. Bortoletto,¹³⁵ D. Boscherini,^{23b} M. Bosman,¹⁴ J. D. Bossio Sola,¹⁰⁴ K. Bouaouda,^{35a} J. Boudreau,¹³⁹ E. V. Bouhova-Thacker,⁹⁰ D. Boumediene,³⁸ S. K. Boutle,⁵⁷ A. Boveia,¹²⁷ J. Boyd,³⁶ D. Boye,^{33b,h} I. R. Boyko,⁸⁰ A. J. Bozson,⁹⁴ J. Bracinik,²¹ N. Brahimi,¹⁰² G. Brandt,¹⁸² O. Brandt,³² F. Braren,⁴⁶ B. Brau,¹⁰³ J. E. Brau,¹³² W. D. Breaden Madden,⁵⁷ K. Brendlinger,⁴⁶ L. Brenner,⁴⁶ R. Brenner,¹⁷² S. Bressler,¹⁸⁰ B. Brickwedde,¹⁰⁰ D. L. Briglin,²¹ D. Britton,⁵⁷ D. Britzger,¹¹⁵ I. Brock,²⁴ R. Brock,¹⁰⁷ G. Brooijmans,³⁹ W. K. Brooks,^{147c} E. Brost,¹²¹ J. H. Broughton,²¹ P. A. Bruckman de Renstrom,⁸⁵ D. Bruncko,^{28b} A. Bruni,^{23b} G. Bruni,^{23b} L. S. Bruni,¹²⁰ S. Bruno,^{74a,74b} M. Bruschi,^{23b} N. Bruscinò,^{73a,73b} P. Bryant,³⁷ L. Bryngemark,⁹⁷ T. Buanes,¹⁷ Q. Buat,³⁶ P. Buchholz,¹⁵¹ A. G. Buckley,⁵⁷ I. A. Budagov,⁸⁰ M. K. Bugge,¹³⁴ F. Bühner,⁵² O. Bulekov,¹¹² T. J. Burch,¹²¹ S. Burdin,⁹¹ C. D. Burgard,¹²⁰ A. M. Burger,¹³⁰ B. Burghgrave,⁸ J. T. P. Burr,⁴⁶ C. D. Burton,¹¹ J. C. Burzynski,¹⁰³ V. Büscher,¹⁰⁰ E. Buschmann,⁵³ P. J. Bussey,⁵⁷ J. M. Butler,²⁵ C. M. Buttar,⁵⁷ J. M. Butterworth,⁹⁵ P. Butti,³⁶ W. Buttinger,³⁶ C. J. Buxo Vazquez,¹⁰⁷ A. Buzatu,¹⁵⁸ A. R. Buzykaev,^{122b,122a} G. Cabras,^{23b,23a} S. Cabrera Urbán,¹⁷⁴ D. Caforio,⁵⁶ H. Cai,¹⁷³ V. M. M. Cairo,¹⁵³ O. Cakir,^{4a} N. Calace,³⁶ P. Calafiura,¹⁸ A. Calandri,¹⁰² G. Calderini,¹³⁶ P. Calfayan,⁶⁶ G. Callea,⁵⁷ L. P. Caloba,^{81b} A. Caltabiano,^{74a,74b} S. Calvente Lopez,⁹⁹ D. Calvet,³⁸ S. Calvet,³⁸ T. P. Calvet,¹⁵⁵ M. Calvetti,^{72a,72b} R. Camacho Toro,¹³⁶ S. Camarda,³⁶ D. Camarero Munoz,⁹⁹ P. Camarri,^{74a,74b} D. Cameron,¹³⁴ R. Caminal Armadans,¹⁰³ C. Camincher,³⁶ S. Campana,³⁶ M. Campanelli,⁹⁵ A. Camplani,⁴⁰ A. Campoverde,¹⁵¹ V. Canale,^{70a,70b} A. Canesse,¹⁰⁴ M. Cano Bret,^{60c} J. Cantero,¹³⁰ T. Cao,¹⁶¹ Y. Cao,¹⁷³ M. D. M. Capeans Garrido,³⁶ M. Capua,^{41b,41a} R. Cardarelli,^{74a} F. Cardillo,¹⁴⁹ G. Carducci,^{41b,41a} I. Carli,¹⁴³ T. Carli,³⁶ G. Carlino,^{70a} B. T. Carlson,¹³⁹ L. Carminati,^{69a,69b} R. M. D. Carney,¹⁵³ S. Caron,¹¹⁹ E. Carquin,^{147c} S. Carrá,⁴⁶ J. W. S. Carter,¹⁶⁷ M. P. Casado,^{14,i} A. F. Casha,¹⁶⁷ D. W. Casper,¹⁷¹ R. Castelijns,¹²⁰ F. L. Castillo,¹⁷⁴ L. Castillo Garcia,¹⁴ V. Castillo Gimenez,¹⁷⁴ N. F. Castro,^{140a,140e} A. Catinaccio,³⁶ J. R. Catmore,¹³⁴ A. Cattai,³⁶ V. Cavaliere,²⁹ E. Cavallaro,¹⁴ M. Cavalli-Sforza,¹⁴ V. Cavalzani,^{72a,72b} E. Celebi,^{12b} L. Cerda Alberich,¹⁷⁴ K. Cerny,¹³¹ A. S. Cerqueira,^{81a} A. Cerri,¹⁵⁶ L. Cerrito,^{74a,74b} F. Cerutti,¹⁸ A. Cervelli,^{23b,23a} S. A. Cetin,^{12b} Z. Chadi,^{35a} D. Chakraborty,¹²¹ W. S. Chan,¹²⁰ W. Y. Chan,⁹¹ J. D. Chapman,³² B. Chargeishvili,^{159b} D. G. Charlton,²¹ T. P. Charman,⁹³ C. C. Chau,³⁴ S. Che,¹²⁷ S. Chekanov,⁶ S. V. Chekulaev,^{168a} G. A. Chelkov,^{80j} M. A. Chelstowska,³⁶ B. Chen,⁷⁹ C. Chen,^{60a} C. H. Chen,⁷⁹ H. Chen,²⁹ J. Chen,^{60a} J. Chen,³⁹ J. Chen,²⁶ S. Chen,¹³⁷ S. J. Chen,^{15c} X. Chen,^{15b} Y.-H. Chen,⁴⁶ H. C. Cheng,^{63a} H. J. Cheng,^{15a} A. Cheplakov,⁸⁰ E. Cheremushkina,¹²³ R. Cherkaoui El Moursli,^{35e} E. Cheu,⁷ K. Cheung,⁶⁴ T. J. A. Chevaléras,¹⁴⁵ L. Chevalier,¹⁴⁵ V. Chiarella,⁵¹ G. Chiarelli,^{72a} G. Chiodini,^{68a} A. S. Chisholm,²¹ A. Chitan,^{27b} I. Chiu,¹⁶³ Y. H. Chiu,¹⁷⁶ M. V. Chizhov,⁸⁰ K. Choi,¹¹ A. R. Chomont,^{73a,73b} S. Chouridou,¹⁶² Y. S. Chow,¹²⁰ M. C. Chu,^{63a} X. Chu,^{15a,15d} J. Chudoba,¹⁴¹ A. J. Chuinard,¹⁰⁴ J. J. Chwastowski,⁸⁵ L. Chytka,¹³¹ D. Cieri,¹¹⁵ K. M. Ciesla,⁸⁵ D. Cinca,⁴⁷ V. Cindro,⁹² I. A. Cioară,^{27b} A. Ciocio,¹⁸ F. Ciotto,^{70a,70b} Z. H. Citron,^{180,k} M. Citterio,^{69a} D. A. Ciubotaru,^{27b} B. M. Ciungu,¹⁶⁷ A. Clark,⁵⁴ M. R. Clark,³⁹ P. J. Clark,⁵⁰ C. Clement,^{45a,45b} Y. Coadou,¹⁰² M. Cobal,^{67a,67c} A. Coccaro,^{55b} J. Cochran,⁷⁹ H. Cohen,¹⁶¹ A. E. C. Coimbra,³⁶ L. Colasurdo,¹¹⁹ B. Cole,³⁹ A. P. Colijn,¹²⁰ J. Collot,⁵⁸ P. Conde Muiño,^{140a,140h} S. H. Connell,^{33b}

I. A. Connelly,⁵⁷ S. Constantinescu,^{27b} F. Conventi,^{70a,1} A. M. Cooper-Sarkar,¹³⁵ F. Cormier,¹⁷⁵ K. J. R. Cormier,¹⁶⁷
L. D. Corpe,⁹⁵ M. Corradi,^{73a,73b} E. E. Corrigan,⁹⁷ F. Corriveau,^{104,m} A. Cortes-Gonzalez,³⁶ M. J. Costa,¹⁷⁴ F. Costanza,⁵
D. Costanzo,¹⁴⁹ G. Cowan,⁹⁴ J. W. Cowley,³² J. Crane,¹⁰¹ K. Cranmer,¹²⁵ S. J. Crawley,⁵⁷ R. A. Creager,¹³⁷
S. Crépé-Renaudin,⁵⁸ F. Crescioli,¹³⁶ M. Cristinziani,²⁴ V. Croft,¹⁷⁰ G. Crosetti,^{41b,41a} A. Cueto,⁵
T. Cuhadar Donszelmann,¹⁴⁹ A. R. Cukierman,¹⁵³ W. R. Cunningham,⁵⁷ S. Czekerda,⁸⁵ P. Czodrowski,³⁶
M. J. Da Cunha Sargedas De Sousa,^{60b} J. V. Da Fonseca Pinto,^{81b} C. Da Via,¹⁰¹ W. Dabrowski,^{84a} F. Dachs,³⁶ T. Dado,^{28a}
S. Dahbi,^{33d} T. Dai,¹⁰⁶ C. Dallapiccola,¹⁰³ M. Dam,⁴⁰ G. D'amen,²⁹ V. D'Amico,^{75a,75b} J. Damp,¹⁰⁰ J. R. Dandoy,¹³⁷
M. F. Daneri,³⁰ N. P. Dang,^{181,e} N. S. Dann,¹⁰¹ M. Danninger,¹⁷⁵ V. Dao,³⁶ G. Darbo,^{55b} O. Dartsis,⁵ A. Dattagupta,¹³²
T. Daubney,⁴⁶ S. D'Auria,^{69a,69b} C. David,^{168b} T. Davidek,¹⁴³ D. R. Davis,⁴⁹ I. Dawson,¹⁴⁹ K. De,⁸ R. De Asmundis,^{70a}
M. De Beurs,¹²⁰ S. De Castro,^{23b,23a} S. De Cecco,^{73a,73b} N. De Groot,¹¹⁹ P. de Jong,¹²⁰ H. De la Torre,¹⁰⁷ A. De Maria,^{15c}
D. De Pedis,^{73a} A. De Salvo,^{73a} U. De Sanctis,^{74a,74b} M. De Santis,^{74a,74b} A. De Santo,¹⁵⁶ K. De Vasconcelos Corga,¹⁰²
J. B. De Vivie De Regie,⁶⁵ C. Debenedetti,¹⁴⁶ D. V. Dedovich,⁸⁰ A. M. Deiana,⁴² J. Del Peso,⁹⁹ Y. Delabat Diaz,⁴⁶
D. Delgove,⁶⁵ F. Deliot,^{145,n} C. M. Delitzsch,⁷ M. Della Pietra,^{70a,70b} D. Della Volpe,⁵⁴ A. Dell'Acqua,³⁶ L. Dell'Asta,^{74a,74b}
M. Delmastro,⁵ C. Delporte,⁶⁵ P. A. Delsart,⁵⁸ D. A. DeMarco,¹⁶⁷ S. Demers,¹⁸³ M. Demichev,⁸⁰ G. Demontigny,¹¹⁰
S. P. Denisov,¹²³ L. D'Eramo,¹³⁶ D. Derendarz,⁸⁵ J. E. Derkaoui,^{35d} F. Derue,¹³⁶ P. Dervan,⁹¹ K. Desch,²⁴ C. Deterre,⁴⁶
K. Dette,¹⁶⁷ C. Deutsch,²⁴ M. R. Devesa,³⁰ P. O. Deviveiros,³⁶ A. Dewhurst,¹⁴⁴ F. A. Di Bello,⁵⁴ A. Di Ciaccio,^{74a,74b}
L. Di Ciaccio,⁵ W. K. Di Clemente,¹³⁷ C. Di Donato,^{70a,70b} A. Di Girolamo,³⁶ G. Di Gregorio,^{72a,72b} B. Di Micco,^{75a,75b}
R. Di Nardo,^{75a,75b} K. F. Di Petrillo,⁵⁹ R. Di Sipio,¹⁶⁷ D. Di Valentino,³⁴ C. Diaconu,¹⁰² F. A. Dias,⁴⁰ T. Dias Do Vale,^{140a}
M. A. Diaz,^{147a} J. Dickinson,¹⁸ E. B. Diehl,¹⁰⁶ J. Dietrich,¹⁹ S. Díez Cornell,⁴⁶ A. Dimitrievska,¹⁸ W. Ding,^{15b}
J. Dingfelder,²⁴ F. Dittus,³⁶ F. Djama,¹⁰² T. Djobava,^{159b} J. I. Djuvsland,¹⁷ M. A. B. Do Vale,^{81c} M. Dobre,^{27b}
D. Dodsworth,²⁶ C. Doglioni,⁹⁷ J. Dolejsi,¹⁴³ Z. Dolezal,¹⁴³ M. Donadelli,^{81d} B. Dong,^{60c} J. Donini,³⁸ A. D'onofrio,^{15c}
M. D'Onofrio,⁹¹ J. Dopke,¹⁴⁴ A. Doria,^{70a} M. T. Dova,⁸⁹ A. T. Doyle,⁵⁷ E. Drechsler,¹⁵² E. Dreyer,¹⁵² T. Dreyer,⁵³
A. S. Drobac,¹⁷⁰ D. Du,^{60b} Y. Duan,^{60b} F. Dubinin,¹¹¹ M. Dubovsky,^{28a} A. Dubreuil,⁵⁴ E. Duchovni,¹⁸⁰ G. Duckeck,¹¹⁴
A. Ducourthial,¹³⁶ O. A. Ducu,¹¹⁰ D. Duda,¹¹⁵ A. Dudarev,³⁶ A. C. Dudder,¹⁰⁰ E. M. Duffield,¹⁸ L. Dufлот,⁶⁵ M. Dührssen,³⁶
C. Dülsen,¹⁸² M. Dumancic,¹⁸⁰ A. E. Dumitriu,^{27b} A. K. Duncan,⁵⁷ M. Dunford,^{61a} A. Duperrin,¹⁰² H. Duran Yildiz,^{4a}
M. Düren,⁵⁶ A. Durglishvili,^{159b} D. Duschinger,⁴⁸ B. Dutta,⁴⁶ D. Duvnjak,¹ G. I. Dyckes,¹³⁷ M. Dyndal,³⁶ S. Dysch,¹⁰¹
B. S. Dziedzic,⁸⁵ K. M. Ecker,¹¹⁵ R. C. Edgar,¹⁰⁶ M. G. Eggleston,⁴⁹ T. Eifert,⁸ G. Eigen,¹⁷ K. Einsweiler,¹⁸ T. Ekelof,¹⁷²
H. El Jarrari,^{35e} R. El Kosseifi,¹⁰² V. Ellajosyula,¹⁷² M. Ellert,¹⁷² F. Ellinghaus,¹⁸² A. A. Elliot,⁹³ N. Ellis,³⁶ J. Elmsheuser,²⁹
M. Elsing,³⁶ D. Emeliyanov,¹⁴⁴ A. Emerman,³⁹ Y. Enari,¹⁶³ M. B. Epland,⁴⁹ J. Erdmann,⁴⁷ A. Ereditato,²⁰ P. A. Erland,⁸⁵
M. Errenst,³⁶ M. Escalier,⁶⁵ C. Escobar,¹⁷⁴ O. Estrada Pastor,¹⁷⁴ E. Etzion,¹⁶¹ H. Evans,⁶⁶ A. Ezhilov,¹³⁸ F. Fabbri,⁵⁷
L. Fabbri,^{23b,23a} V. Fabiani,¹¹⁹ G. Facini,⁹⁵ R. M. Faisca Rodrigues Pereira,^{140a} R. M. Fakhruddinov,¹²³ S. Falciano,^{73a}
P. J. Falke,⁵ S. Falke,⁵ J. Faltova,¹⁴³ Y. Fang,^{15a} Y. Fang,^{15a} G. Fanourakis,⁴⁴ M. Fanti,^{69a,69b} M. Faraj,^{67a,67c,o} A. Farbin,⁸
A. Farilla,^{75a} E. M. Farina,^{71a,71b} T. Farooque,¹⁰⁷ S. Farrell,¹⁸ S. M. Farrington,⁵⁰ P. Farthouat,³⁶ F. Fassi,^{35e} P. Fassnacht,³⁶
D. Fassouliotis,⁹ M. Fauci Giannelli,⁵⁰ W. J. Fawcett,³² L. Fayard,⁶⁵ O. L. Fedin,^{138,p} W. Fedorko,¹⁷⁵ A. Fehr,²⁰
M. Feickert,¹⁷³ L. Feligioni,¹⁰² A. Fell,¹⁴⁹ C. Feng,^{60b} M. Feng,⁴⁹ M. J. Fenton,⁵⁷ A. B. Fenyuk,¹²³ S. W. Ferguson,⁴³
J. Ferrando,⁴⁶ A. Ferrante,¹⁷³ A. Ferrari,¹⁷² P. Ferrari,¹²⁰ R. Ferrari,^{71a} D. E. Ferreira de Lima,^{61b} A. Ferrer,¹⁷⁴ D. Ferrere,⁵⁴
C. Ferretti,¹⁰⁶ F. Fiedler,¹⁰⁰ A. Filipčič,⁹² F. Filthaut,¹¹⁹ K. D. Finelli,²⁵ M. C. N. Fiolhais,^{140a,140c,q} L. Fiorini,¹⁷⁴ F. Fischer,¹¹⁴
W. C. Fisher,¹⁰⁷ I. Fleck,¹⁵¹ P. Fleischmann,¹⁰⁶ R. R. M. Fletcher,¹³⁷ T. Flick,¹⁸² B. M. Flierl,¹¹⁴ L. Flores,¹³⁷
L. R. Flores Castillo,^{63a} F. M. Follega,^{76a,76b} N. Fomin,¹⁷ J. H. Foo,¹⁶⁷ G. T. Forcolin,^{76a,76b} A. Formica,¹⁴⁵ F. A. Förster,¹⁴
A. C. Forti,¹⁰¹ A. G. Foster,²¹ M. G. Foti,¹³⁵ D. Fournier,⁶⁵ H. Fox,⁹⁰ P. Francavilla,^{72a,72b} S. Francescato,^{73a,73b}
M. Franchini,^{23b,23a} S. Franchino,^{61a} D. Francis,³⁶ L. Franconi,²⁰ M. Franklin,⁵⁹ A. N. Fray,⁹³ P. M. Freeman,²¹ B. Freund,¹¹⁰
W. S. Freund,^{81b} E. M. Freundlich,⁴⁷ D. C. Frizzell,¹²⁹ D. Froidevaux,³⁶ J. A. Frost,¹³⁵ C. Fukunaga,¹⁶⁴
E. Fullana Torregrosa,¹⁷⁴ E. Fumagalli,^{55b,55a} T. Fusayasu,¹¹⁶ J. Fuster,¹⁷⁴ A. Gabrielli,^{23b,23a} A. Gabrielli,¹⁸ S. Gadatsch,⁵⁴
P. Gadow,¹¹⁵ G. Gagliardi,^{55b,55a} L. G. Gagnon,¹¹⁰ C. Galea,^{27b} B. Galhardo,^{140a} G. E. Gallardo,¹³⁵ E. J. Gallas,¹³⁵
B. J. Gallop,¹⁴⁴ G. Galster,⁴⁰ R. Gamboa Goni,⁹³ K. K. Gan,¹²⁷ S. Ganguly,¹⁸⁰ J. Gao,^{60a} Y. Gao,⁵⁰ Y. S. Gao,^{31,r} C. García,¹⁷⁴
J. E. García Navarro,¹⁷⁴ J. A. García Pascual,^{15a} C. Garcia-Argos,⁵² M. Garcia-Sciveres,¹⁸ R. W. Gardner,³⁷ N. Garelli,¹⁵³
S. Gargiulo,⁵² C. A. Garner,¹⁶⁷ V. Garonne,¹³⁴ S. J. Gasiorowski,¹⁴⁸ P. Gaspar,^{81b} A. Gaudiello,^{55b,55a} G. Gaudio,^{71a}
I. L. Gavrilenko,¹¹¹ A. Gavriilyuk,¹²⁴ C. Gay,¹⁷⁵ G. Gaycken,⁴⁶ E. N. Gazis,¹⁰ A. A. Geanta,^{27b} C. M. Gee,¹⁴⁶ C. N. P. Gee,¹⁴⁴
J. Geisen,⁵³ M. Geisen,¹⁰⁰ C. Gemme,^{55b} M. H. Genest,⁵⁸ C. Geng,¹⁰⁶ S. Gentile,^{73a,73b} S. George,⁹⁴ T. Gerialis,⁴⁴

L. O. Gerlach,⁵³ P. Gessinger-Befurt,¹⁰⁰ G. Gessner,⁴⁷ S. Ghasemi,¹⁵¹ M. Ghasemi Bostanabad,¹⁷⁶ M. Ghneimat,¹⁵¹ A. Ghosh,⁶⁵ A. Ghosh,⁷⁸ B. Giacobbe,^{23b} S. Giagu,^{73a,73b} N. Giangiacomi,^{23b,23a} P. Giannetti,^{72a} A. Giannini,^{70a,70b} G. Giannini,¹⁴ S. M. Gibson,⁹⁴ M. Gignac,¹⁴⁶ D. Gillberg,³⁴ G. Gilles,¹⁸² D. M. Gingrich,^{3,d} M. P. Giordani,^{67a,67c} P. F. Giraud,¹⁴⁵ G. Giugliarelli,^{67a,67c} D. Giugni,^{69a} F. Giuli,^{74a,74b} S. Gkaitatzis,¹⁶² I. Gkialas,^{9,s} E. L. Gkoukousis,¹⁴ P. Gkoutoumis,¹⁰ L. K. Gladilin,¹¹³ C. Glasman,⁹⁹ J. Glatzer,¹⁴ P. C. F. Glaysheer,⁴⁶ A. Glazov,⁴⁶ G. R. Gledhill,¹³² M. Goblirsch-Kolb,²⁶ D. Godin,¹¹⁰ S. Goldfarb,¹⁰⁵ T. Golling,⁵⁴ D. Golubkov,¹²³ A. Gomes,^{140a,140b} R. Goncalves Gama,⁵³ R. Gonçalves,^{140a} G. Gonella,⁵² L. Gonella,²¹ A. Gongadze,⁸⁰ F. Gonnella,²¹ J. L. Gonski,³⁹ S. González de la Hoz,¹⁷⁴ S. Gonzalez-Sevilla,⁵⁴ G. R. Gonzalvo Rodriguez,¹⁷⁴ L. Goossens,³⁶ N. A. Gorasia,²¹ P. A. Gorbounov,¹²⁴ H. A. Gordon,²⁹ B. Gorini,³⁶ E. Gorini,^{68a,68b} A. Gorišek,⁹² A. T. Goshaw,⁴⁹ M. I. Gostkin,⁸⁰ C. A. Gottardo,¹¹⁹ M. Goughri,^{35b} A. G. Goussiou,¹⁴⁸ N. Govender,^{33b} C. Goy,⁵ E. Gozani,¹⁶⁰ I. Grabowska-Bold,^{84a} E. C. Graham,⁹¹ J. Gramling,¹⁷¹ E. Gramstad,¹³⁴ S. Grancagnolo,¹⁹ M. Grandi,¹⁵⁶ V. Gratchev,¹³⁸ P. M. Gravila,^{27f} F. G. Gravili,^{68a,68b} C. Gray,⁵⁷ H. M. Gray,¹⁸ C. Greife,²⁴ K. Gregersen,⁹⁷ I. M. Gregor,⁴⁶ P. Grenier,¹⁵³ K. Grevtsov,⁴⁶ C. Grieco,¹⁴ N. A. Grieser,¹²⁹ A. A. Grillo,¹⁴⁶ K. Grimm,^{31,t} S. Grinstein,^{14,u} J.-F. Grivaz,⁶⁵ S. Groh,¹⁰⁰ E. Gross,¹⁸⁰ J. Grosse-Knetter,⁵³ Z. J. Grout,⁹⁵ C. Grud,¹⁰⁶ A. Grummer,¹¹⁸ L. Guan,¹⁰⁶ W. Guan,¹⁸¹ C. Gubbels,¹⁷⁵ J. Guenther,³⁶ A. Guerguichon,⁶⁵ J. G. R. Guerrero Rojas,¹⁷⁴ F. Guescini,¹¹⁵ D. Guest,¹⁷¹ R. Gugel,⁵² T. Guillemin,⁵ S. Guindon,³⁶ U. Gul,⁵⁷ J. Guo,^{60c} W. Guo,¹⁰⁶ Y. Guo,^{60a,v} Z. Guo,¹⁰² R. Gupta,⁴⁶ S. Gurbuz,^{12c} G. Gustavino,¹²⁹ M. Guth,⁵² P. Gutierrez,¹²⁹ C. Gutschow,⁹⁵ C. Guyot,¹⁴⁵ C. Gwenlan,¹³⁵ C. B. Gwilliam,⁹¹ A. Haas,¹²⁵ C. Haber,¹⁸ H. K. Hadavand,⁸ A. Hadeif,^{60a} S. Hageböck,³⁶ M. Haleem,¹⁷⁷ J. Haley,¹³⁰ G. Halladjian,¹⁰⁷ G. D. Hallewell,¹⁰² K. Hamacher,¹⁸² P. Hamal,¹³¹ K. Hamano,¹⁷⁶ H. Hamdaoui,^{35e} M. Hamer,²⁴ G. N. Hamity,⁵⁰ K. Han,^{60a,w} L. Han,^{60a} S. Han,^{15a} Y. F. Han,¹⁶⁷ K. Hanagaki,^{82,x} M. Hance,¹⁴⁶ D. M. Handl,¹¹⁴ B. Haney,¹³⁷ R. Hankache,¹³⁶ E. Hansen,⁹⁷ J. B. Hansen,⁴⁰ J. D. Hansen,⁴⁰ M. C. Hansen,²⁴ P. H. Hansen,⁴⁰ E. C. Hanson,¹⁰¹ K. Hara,¹⁶⁹ T. Harenberg,¹⁸² S. Harkusha,¹⁰⁸ P. F. Harrison,¹⁷⁸ N. M. Hartmann,¹¹⁴ Y. Hasegawa,¹⁵⁰ A. Hasib,⁵⁰ S. Hassani,¹⁴⁵ S. Haug,²⁰ R. Hauser,¹⁰⁷ L. B. Havener,³⁹ M. Havranek,¹⁴² C. M. Hawkes,²¹ R. J. Hawkings,³⁶ D. Hayden,¹⁰⁷ C. Hayes,¹⁰⁶ R. L. Hayes,¹⁷⁵ C. P. Hays,¹³⁵ J. M. Hays,⁹³ H. S. Hayward,⁹¹ S. J. Haywood,¹⁴⁴ F. He,^{60a} M. P. Heath,⁵⁰ V. Hedberg,⁹⁷ S. Heer,²⁴ K. K. Heidegger,⁵² W. D. Heidorn,⁷⁹ J. Heilman,³⁴ S. Heim,⁴⁶ T. Heim,¹⁸ B. Heinemann,^{46,y} J. J. Heinrich,¹³² L. Heinrich,³⁶ J. Hejbal,¹⁴¹ L. Helary,^{61b} A. Held,¹⁷⁵ S. Hellesund,¹³⁴ C. M. Helling,¹⁴⁶ S. Hellman,^{45a,45b} C. Helsens,³⁶ R. C. W. Henderson,⁹⁰ Y. Heng,¹⁸¹ L. Henkelmann,^{61a} S. Henkelmann,¹⁷⁵ A. M. Henriques Correia,³⁶ H. Herde,²⁶ V. Herget,¹⁷⁷ Y. Hernández Jiménez,^{33d} H. Herr,¹⁰⁰ M. G. Herrmann,¹¹⁴ T. Herrmann,⁴⁸ G. Herten,⁵² R. Hertenberger,¹¹⁴ L. Hervas,³⁶ T. C. Herwig,¹³⁷ G. G. Hesketh,⁹⁵ N. P. Hessey,^{168a} A. Higashida,¹⁶³ S. Higashino,⁸² E. Higón-Rodríguez,¹⁷⁴ K. Hildebrand,³⁷ J. C. Hill,³² K. K. Hill,²⁹ K. H. Hiller,⁴⁶ S. J. Hillier,²¹ M. Hils,⁴⁸ I. Hinchliffe,¹⁸ F. Hinterkeuser,²⁴ M. Hirose,¹³³ S. Hirose,⁵² D. Hirschbuehl,¹⁸² B. Hiti,⁹² O. Hladik,¹⁴¹ D. R. Hlaluku,^{33d} X. Hoad,⁵⁰ J. Hobbs,¹⁵⁵ N. Hod,¹⁸⁰ M. C. Hodgkinson,¹⁴⁹ A. Hoecker,³⁶ D. Hohn,⁵² D. Hohov,⁶⁵ T. Holm,²⁴ T. R. Holmes,³⁷ M. Holzbock,¹¹⁴ L. B. A. H. Hommels,³² S. Honda,¹⁶⁹ T. M. Hong,¹³⁹ J. C. Honig,⁵² A. Hönle,¹¹⁵ B. H. Hooberman,¹⁷³ W. H. Hopkins,⁶ Y. Horii,¹¹⁷ P. Horn,⁴⁸ L. A. Horyn,³⁷ S. Hou,¹⁵⁸ A. Hoummada,^{35a} J. Howarth,¹⁰¹ J. Hoya,⁸⁹ M. Hrabovsky,¹³¹ J. Hrdinka,⁷⁷ I. Hristova,¹⁹ J. Hrivnac,⁶⁵ A. Hrynevich,¹⁰⁹ T. Hryn'ova,⁵ P. J. Hsu,⁶⁴ S.-C. Hsu,¹⁴⁸ Q. Hu,²⁹ S. Hu,^{60c} Y. F. Hu,^{15a,15d} D. P. Huang,⁹⁵ Y. Huang,^{60a} Y. Huang,^{15a} Z. Hubacek,¹⁴² F. Hubaut,¹⁰² M. Huebner,²⁴ F. Huegging,²⁴ T. B. Huffman,¹³⁵ M. Huhtinen,³⁶ R. F. H. Hunter,³⁴ P. Huo,¹⁵⁵ N. Huseynov,^{80,z} J. Huston,¹⁰⁷ J. Huth,⁵⁹ R. Hyneman,¹⁰⁶ S. Hyrych,^{28a} G. Iacobucci,⁵⁴ G. Iakovidis,²⁹ I. Ibragimov,¹⁵¹ L. Iconomidou-Fayard,⁶⁵ P. Iengo,³⁶ R. Ignazzi,⁴⁰ O. Igonkina,^{120,a,aa} R. Iguchi,¹⁶³ T. Iizawa,⁵⁴ Y. Ikegami,⁸² M. Ikeno,⁸² D. Iliadis,¹⁶² N. Ilic,^{119,167,m} F. Iltzsche,⁴⁸ G. Introzzi,^{71a,71b} M. Iodice,^{75a} K. Iordanidou,^{168a} V. Ippolito,^{73a,73b} M. F. Isacson,¹⁷² M. Ishino,¹⁶³ W. Islam,¹³⁰ C. Issever,^{19,46} S. Istin,¹⁶⁰ F. Ito,¹⁶⁹ J. M. Iturbe Ponce,^{63a} R. Iuppa,^{76a,76b} A. Ivina,¹⁸⁰ H. Iwasaki,⁸² J. M. Izen,⁴³ V. Izzo,^{70a} P. Jacka,¹⁴¹ P. Jackson,¹ R. M. Jacobs,²⁴ B. P. Jaeger,¹⁵² V. Jain,² G. Jäkel,¹⁸² K. B. Jakobi,¹⁰⁰ K. Jakobs,⁵² T. Jakoubek,¹⁴¹ J. Jamieson,⁵⁷ K. W. Janas,^{84a} R. Jansky,⁵⁴ J. Janssen,²⁴ M. Janus,⁵³ P. A. Janus,^{84a} G. Jarlskog,⁹⁷ N. Javadov,^{80,z} T. Javůrek,³⁶ M. Javurkova,¹⁰³ F. Jeanneau,¹⁴⁵ L. Jeanty,¹³² J. Jejelava,^{159a} A. Jelinskas,¹⁷⁸ P. Jenni,^{52,bb} J. Jeong,⁴⁶ N. Jeong,⁴⁶ S. Jézéquel,⁵ H. Ji,¹⁸¹ J. Jia,¹⁵⁵ H. Jiang,⁷⁹ Y. Jiang,^{60a} Z. Jiang,^{153,cc} S. Jiggins,⁵² F. A. Jimenez Morales,³⁸ J. Jimenez Pena,¹¹⁵ S. Jin,^{15c} A. Jinaru,^{27b} O. Jinnouchi,¹⁶⁵ H. Jivan,^{33d} P. Johansson,¹⁴⁹ K. A. Johns,⁷ C. A. Johnson,⁶⁶ R. W. L. Jones,⁹⁰ S. D. Jones,¹⁵⁶ S. Jones,⁷ T. J. Jones,⁹¹ J. Jongmanns,^{61a} P. M. Jorge,^{140a} J. Jovicevic,³⁶ X. Ju,¹⁸ J. J. Junggeburth,¹¹⁵ A. Juste Rozas,^{14,u} A. Kaczmarska,⁸⁵ M. Kado,^{73a,73b} H. Kagan,¹²⁷ M. Kagan,¹⁵³ A. Kahn,³⁹ C. Kahra,¹⁰⁰ T. Kaji,¹⁷⁹ E. Kajomovitz,¹⁶⁰ C. W. Kalderon,⁹⁷ A. Kaluza,¹⁰⁰ A. Kamenshchikov,¹²³ M. Kaneda,¹⁶³ N. J. Kang,¹⁴⁶ S. Kang,⁷⁹ L. Kanjir,⁹² Y. Kano,¹¹⁷ V. A. Kantserov,¹¹² J. Kanzaki,⁸²

L. S. Kaplan,¹⁸¹ D. Kar,^{33d} K. Karava,¹³⁵ M. J. Kareem,^{168b} S. N. Karpov,⁸⁰ Z. M. Karpova,⁸⁰ V. Kartvelishvili,⁹⁰
A. N. Karyukhin,¹²³ L. Kashif,¹⁸¹ R. D. Kass,¹²⁷ A. Kastanas,^{45a,45b} C. Kato,^{60d,60c} J. Katzy,⁴⁶ K. Kawade,¹⁵⁰ K. Kawagoe,⁸⁸
T. Kawaguchi,¹¹⁷ T. Kawamoto,¹⁴⁵ G. Kawamura,⁵³ E. F. Kay,¹⁷⁶ V. F. Kazanin,^{122b,122a} R. Keeler,¹⁷⁶ R. Kehoe,⁴²
J. S. Keller,³⁴ E. Kellermann,⁹⁷ D. Kelsey,¹⁵⁶ J. J. Kempster,²¹ J. Kendrick,²¹ K. E. Kennedy,³⁹ O. Kepka,¹⁴¹ S. Kersten,¹⁸²
B. P. Kerševan,⁹² S. Ketabchi Haghghat,¹⁶⁷ M. Khader,¹⁷³ F. Khalil-Zada,¹³ M. Khandoga,¹⁴⁵ A. Khanov,¹³⁰
A. G. Kharlamov,^{122b,122a} T. Kharlamova,^{122b,122a} E. E. Khoda,¹⁷⁵ A. Khodinov,¹⁶⁶ T. J. Khoo,⁵⁴ E. Khramov,⁸⁰
J. Khubua,^{159b} S. Kido,⁸³ M. Kiehn,⁵⁴ C. R. Kilby,⁹⁴ Y. K. Kim,³⁷ N. Kimura,⁹⁵ O. M. Kind,¹⁹ B. T. King,^{91.a}
D. Kirchmeier,⁴⁸ J. Kirk,¹⁴⁴ A. E. Kiryunin,¹¹⁵ T. Kishimoto,¹⁶³ D. P. Kisliuk,¹⁶⁷ V. Kitali,⁴⁶ O. Kivernyk,⁵
T. Klapdor-Kleingrothaus,⁵² M. Klassen,^{61a} C. Klein,³² M. H. Klein,¹⁰⁶ M. Klein,⁹¹ U. Klein,⁹¹ K. Kleinknecht,¹⁰⁰
P. Klimek,¹²¹ A. Klimentov,²⁹ T. Klingl,²⁴ T. Klioutchnikova,³⁶ F. F. Klitzner,¹¹⁴ P. Kluit,¹²⁰ S. Kluth,¹¹⁵ E. Kneringer,⁷⁷
E. B. F. G. Knoop,¹⁰² A. Knue,⁵² D. Kobayashi,⁸⁸ T. Kobayashi,¹⁶³ M. Kobel,⁴⁸ M. Kocian,¹⁵³ T. Kodama,¹⁶³ P. Kodys,¹⁴³
P. T. Koenig,²⁴ T. Koffas,³⁴ N. M. Köhler,³⁶ T. Koi,¹⁵³ M. Kolb,¹⁴⁵ I. Koletsou,⁵ T. Komarek,¹³¹ T. Kondo,⁸² K. Köneke,⁵²
A. X. Y. Kong,¹ A. C. König,¹¹⁹ T. Kono,¹²⁶ R. Konoplich,^{125,dd} V. Konstantinides,⁹⁵ N. Konstantinidis,⁹⁵ B. Konya,⁹⁷
R. Kopeliansky,⁶⁶ S. Koperny,^{84a} K. Korcyl,⁸⁵ K. Kordas,¹⁶² G. Koren,¹⁶¹ A. Korn,⁹⁵ I. Korolkov,¹⁴ E. V. Korolkova,¹⁴⁹
N. Korotkova,¹¹³ O. Kortner,¹¹⁵ S. Kortner,¹¹⁵ T. Kosek,¹⁴³ V. V. Kostyukhin,^{149,166} A. Kotskechagia,⁶⁵ A. Kotwal,⁴⁹
A. Koulouris,¹⁰ A. Kourkoumeli-Charalampidi,^{71a,71b} C. Kourkoumelis,⁹ E. Kourlitis,¹⁴⁹ V. Kouskoura,²⁹
A. B. Kowalewska,⁸⁵ R. Kowalewski,¹⁷⁶ W. Kozanecki,¹⁴⁵ A. S. Kozhin,¹²³ V. A. Kramarenko,¹¹³ G. Kramberger,⁹²
D. Krasnopevtsev,^{60a} M. W. Krasny,¹³⁶ A. Krasznahorkay,³⁶ D. Krauss,¹¹⁵ J. A. Kremer,^{84a} J. Kretzschmar,⁹¹ P. Krieger,¹⁶⁷
F. Krieter,¹¹⁴ A. Krishnan,^{61b} K. Krizka,¹⁸ K. Kroeninger,⁴⁷ H. Kroha,¹¹⁵ J. Kroll,¹⁴¹ J. Kroll,¹³⁷ K. S. Krowpman,¹⁰⁷
J. Krstic,¹⁶ U. Kruchonak,⁸⁰ H. Krüger,²⁴ N. Krumnack,⁷⁹ M. C. Kruse,⁴⁹ J. A. Krzysiak,⁸⁵ T. Kubota,¹⁰⁵ O. Kuchinskaia,¹⁶⁶
S. Kunday,^{4b} J. T. Kuechler,⁴⁶ S. Kuehn,³⁶ A. Kugel,^{61a} T. Kuhl,⁴⁶ V. Kukhtin,⁸⁰ R. Kukla,¹⁰² Y. Kulchitsky,^{108,ee}
S. Kuleshov,^{147c} Y. P. Kulinich,¹⁷³ M. Kuna,⁵⁸ T. Kunigo,⁸⁶ A. Kupco,¹⁴¹ T. Kupfer,⁴⁷ O. Kuprash,⁵² H. Kurashige,⁸³
L. L. Kurchaninov,^{168a} Y. A. Kurochkin,¹⁰⁸ A. Kurova,¹¹² M. G. Kurth,^{15a,15d} E. S. Kuwertz,³⁶ M. Kuze,¹⁶⁵ A. K. Kvam,¹⁴⁸
J. Kvita,¹³¹ T. Kwan,¹⁰⁴ A. La Rosa,¹¹⁵ L. La Rotonda,^{41b,41a} F. La Ruffa,^{41b,41a} C. Lacasta,¹⁷⁴ F. Lacava,^{73a,73b}
D. P. J. Lack,¹⁰¹ H. Lacker,¹⁹ D. Lacour,¹³⁶ E. Ladygin,⁸⁰ R. Lafaye,⁵ B. Laforge,¹³⁶ T. Lagouri,^{33d} S. Lai,⁵³
I. K. Lakomic,^{84a} S. Lammers,⁶⁶ W. Lampl,⁷ C. Lampoudis,¹⁶² E. Lançon,²⁹ U. Landgraf,⁵² M. P. J. Landon,⁹³
M. C. Lanfermann,⁵⁴ V. S. Lang,⁴⁶ J. C. Lange,⁵³ R. J. Langenberg,¹⁰³ A. J. Lankford,¹⁷¹ F. Lanni,²⁹ K. Lantzsch,²⁴
A. Lanza,^{71a} A. Lapertosa,^{55b,55a} S. Laplace,¹³⁶ J. F. Laporte,¹⁴⁵ T. Lari,^{69a} F. Lasagni Manghi,^{23b,23a} M. Lassnig,³⁶
T. S. Lau,^{63a} A. Laudrain,⁶⁵ A. Laurier,³⁴ M. Lavorgna,^{70a,70b} S. D. Lawlor,⁹⁴ M. Lazzaroni,^{69a,69b} B. Le,¹⁰⁵ E. Le Guirriec,¹⁰²
M. LeBlanc,⁷ T. LeCompte,⁶ F. Ledroit-Guillon,⁵⁸ A. C. A. Lee,⁹⁵ C. A. Lee,²⁹ G. R. Lee,¹⁷ L. Lee,⁵⁹ S. C. Lee,¹⁵⁸
S. J. Lee,³⁴ S. Lee,⁷⁹ B. Lefebvre,^{168a} H. P. Lefebvre,⁹⁴ M. Lefebvre,¹⁷⁶ C. Leggett,¹⁸ K. Lehmann,¹⁵² N. Lehmann,¹⁸²
G. Lehmann Miotto,³⁶ W. A. Leight,⁴⁶ A. Leisos,^{162,ff} M. A. L. Leite,^{81d} C. E. Leitgeb,¹¹⁴ R. Leitner,¹⁴³ D. Lellouch,^{180,a}
K. J. C. Leney,⁴² T. Lenz,²⁴ R. Leone,⁷ S. Leone,^{72a} C. Leonidopoulos,⁵⁰ A. Leopold,¹³⁶ C. Leroy,¹¹⁰ R. Les,¹⁶⁷
C. G. Lester,³² M. Levchenko,¹³⁸ J. Levêque,⁵ D. Levin,¹⁰⁶ L. J. Levinson,¹⁸⁰ D. J. Lewis,²¹ B. Li,^{15b} B. Li,¹⁰⁶ C-Q. Li,^{60a}
F. Li,^{60c} H. Li,^{60a} H. Li,^{60b} J. Li,^{60c} K. Li,¹⁴⁸ L. Li,^{60c} M. Li,^{15a,15d} Q. Li,^{15a,15d} Q. Y. Li,^{60a} S. Li,^{60d,60c} X. Li,⁴⁶ Y. Li,⁴⁶ Z. Li,^{60b}
Z. Liang,^{15a} B. Liberti,^{74a} A. Liblong,¹⁶⁷ K. Lie,^{63c} S. Lim,²⁹ C. Y. Lin,³² K. Lin,¹⁰⁷ T. H. Lin,¹⁰⁰ R. A. Linck,⁶⁶
J. H. Lindon,²¹ A. L. Lioni,⁵⁴ E. Lipeles,¹³⁷ A. Lipniacka,¹⁷ T. M. Liss,^{173,gg} A. Lister,¹⁷⁵ J. D. Little,⁸ B. Liu,⁷⁹ B. L. Liu,⁶
H. B. Liu,²⁹ H. Liu,¹⁰⁶ J. B. Liu,^{60a} J. K. K. Liu,³⁷ K. Liu,¹³⁶ M. Liu,^{60a} P. Liu,¹⁸ Y. Liu,^{15a,15d} Y. L. Liu,¹⁰⁶ Y. W. Liu,^{60a}
M. Livan,^{71a,71b} A. Lleres,⁵⁸ J. Llorente Merino,¹⁵² S. L. Lloyd,⁹³ C. Y. Lo,^{63b} F. Lo Sterzo,⁴² E. M. Lobodzinska,⁴⁶ P. Loch,⁷
S. Loffredo,^{74a,74b} T. Lohse,¹⁹ K. Lohwasser,¹⁴⁹ M. Lokajicek,¹⁴¹ J. D. Long,¹⁷³ R. E. Long,⁹⁰ L. Longo,³⁶ K. A. Looper,¹²⁷
J. A. Lopez,^{147c} I. Lopez Paz,¹⁰¹ A. Lopez Solis,¹⁴⁹ J. Lorenz,¹¹⁴ N. Lorenzo Martinez,⁵ A. M. Lory,¹¹⁴ M. Losada,²²
P. J. Lösel,¹¹⁴ A. Lösle,⁵² X. Lou,⁴⁶ X. Lou,^{15a} A. Lounis,⁶⁵ J. Love,⁶ P. A. Love,⁹⁰ J. J. Lozano Bahilo,¹⁷⁴ M. Lu,^{60a}
Y. J. Lu,⁶⁴ H. J. Lubatti,¹⁴⁸ C. Luci,^{73a,73b} A. Lucotte,⁵⁸ C. Luedtke,⁵² F. Luehring,⁶⁶ I. Luise,¹³⁶ L. Luminari,^{73a}
B. Lund-Jensen,¹⁵⁴ M. S. Lutz,¹⁰³ D. Lynn,²⁹ H. Lyons,⁹¹ R. Lysak,¹⁴¹ E. Lytken,⁹⁷ F. Lyu,^{15a} V. Lyubushkin,⁸⁰
T. Lyubushkina,⁸⁰ H. Ma,²⁹ L. L. Ma,^{60b} Y. Ma,^{60b} G. Maccarrone,⁵¹ A. Macchiolo,¹¹⁵ C. M. Macdonald,¹⁴⁹
J. Machado Miguens,¹³⁷ D. Madaffari,¹⁷⁴ R. Madar,³⁸ W. F. Mader,⁴⁸ M. Madugoda Ralalage Don,¹³⁰ N. Madysa,⁴⁸
J. Maeda,⁸³ T. Maeno,²⁹ M. Maerker,⁴⁸ A. S. Maevskiy,¹¹³ V. Magerl,⁵² N. Magini,⁷⁹ D. J. Mahon,³⁹ C. Maidantchik,^{81b}
T. Maier,¹¹⁴ A. Maio,^{140a,140b,140d} K. Maj,^{84a} O. Majersky,^{28a} S. Majewski,¹³² Y. Makida,⁸² N. Makovec,⁶⁵ B. Malaescu,¹³⁶
Pa. Malecki,⁸⁵ V. P. Maleev,¹³⁸ F. Malek,⁵⁸ U. Mallik,⁷⁸ D. Malon,⁶ C. Malone,³² S. Maltezos,¹⁰ S. Malyukov,⁸⁰

J. Mamuzic,¹⁷⁴ G. Mancini,⁵¹ I. Mandić,⁹² L. Manhaes de Andrade Filho,^{81a} I. M. Maniatis,¹⁶² J. Manjarres Ramos,⁴⁸
 K. H. Mankinen,⁹⁷ A. Mann,¹¹⁴ A. Manousos,⁷⁷ B. Mansoulie,¹⁴⁵ I. Manthos,¹⁶² S. Manzoni,¹²⁰ A. Marantis,¹⁶²
 G. Marceca,³⁰ L. Marchese,¹³⁵ G. Marchiori,¹³⁶ M. Marcisovsky,¹⁴¹ L. Marcoccia,^{74a,74b} C. Marcon,⁹⁷ C. A. Marin Tobon,³⁶
 M. Marjanovic,¹²⁹ Z. Marshall,¹⁸ M. U. F. Martensson,¹⁷² S. Marti-Garcia,¹⁷⁴ C. B. Martin,¹²⁷ T. A. Martin,¹⁷⁸ V. J. Martin,⁵⁰
 B. Martin dit Latour,¹⁷ L. Martinelli,^{75a,75b} M. Martinez,^{14,u} V. I. Martinez Outschoorn,¹⁰³ S. Martin-Haugh,¹⁴⁴
 V. S. Martoiu,^{27b} A. C. Martyniuk,⁹⁵ A. Marzin,³⁶ S. R. Maschek,¹¹⁵ L. Masetti,¹⁰⁰ T. Mashimo,¹⁶³ R. Mashinistov,¹¹¹
 J. Masik,¹⁰¹ A. L. Maslennikov,^{122b,122a} L. Massa,^{74a,74b} P. Massarotti,^{70a,70b} P. Mastrandrea,^{72a,72b} A. Mastroberardino,^{41b,41a}
 T. Masubuchi,¹⁶³ D. Matakias,¹⁰ A. Matic,¹¹⁴ N. Matsuzawa,¹⁶³ P. Mättig,²⁴ J. Maurer,^{27b} B. Maček,⁹²
 D. A. Maximov,^{122b,122a} R. Mazini,¹⁵⁸ I. Maznas,¹⁶² S. M. Mazza,¹⁴⁶ S. P. Mc Kee,¹⁰⁶ T. G. McCarthy,¹¹⁵
 W. P. McCormack,¹⁸ E. F. McDonald,¹⁰⁵ J. A. MCFayden,³⁶ G. Mchedlidze,^{159b} M. A. McKay,⁴² K. D. McLean,¹⁷⁶
 S. J. McMahan,¹⁴⁴ P. C. McNamara,¹⁰⁵ C. J. McNicol,¹⁷⁸ R. A. McPherson,^{176,m} J. E. Mdhului,^{33d} Z. A. Meadows,¹⁰³
 S. Meehan,³⁶ T. Megy,⁵² S. Mehlhase,¹¹⁴ A. Mehta,⁹¹ T. Meideck,⁵⁸ B. Meirose,⁴³ D. Melini,¹⁶⁰ B. R. Mellado Garcia,^{33d}
 J. D. Mellenthin,⁵³ M. Melo,^{28a} F. Meloni,⁴⁶ A. Melzer,²⁴ S. B. Menary,¹⁰¹ E. D. Mendes Gouveia,^{140a,140e} L. Meng,³⁶
 X. T. Meng,¹⁰⁶ S. Menke,¹¹⁵ E. Meoni,^{41b,41a} S. Mergelmeyer,¹⁹ S. A. M. Merkt,¹³⁹ C. Merlassino,¹³⁵ P. Mermod,⁵⁴
 L. Merola,^{70a,70b} C. Meroni,^{69a} G. Merz,¹⁰⁶ O. Meshkov,^{113,111} J. K. R. Meshreki,¹⁵¹ A. Messina,^{73a,73b} J. Metcalfe,⁶
 A. S. Mete,⁶ C. Meyer,⁶⁶ J-P. Meyer,¹⁴⁵ H. Meyer Zu Theenhausen,^{61a} F. Miano,¹⁵⁶ M. Michetti,¹⁹ R. P. Middleton,¹⁴⁴
 L. Mijović,⁵⁰ G. Mikenberg,¹⁸⁰ M. Mikestikova,¹⁴¹ M. Mikuž,⁹² H. Mildner,¹⁴⁹ M. Milesi,¹⁰⁵ A. Milic,¹⁶⁷ D. A. Millar,⁹³
 D. W. Miller,³⁷ A. Milov,¹⁸⁰ D. A. Milstead,^{45a,45b} R. A. Mina,¹⁵³ A. A. Minaenko,¹²³ M. Miñano Moya,¹⁷⁴
 I. A. Minashvili,^{159b} A. I. Mincer,¹²⁵ B. Mindur,^{84a} M. Mineev,⁸⁰ Y. Minegishi,¹⁶³ L. M. Mir,¹⁴ A. Mirto,^{68a,68b}
 K. P. Mistry,¹³⁷ T. Mitani,¹⁷⁹ J. Mitrevski,¹¹⁴ V. A. Mitsou,¹⁷⁴ M. Mittal,^{60c} O. Miu,¹⁶⁷ A. Miucci,²⁰ P. S. Miyagawa,¹⁴⁹
 A. Mizukami,⁸² J. U. Mjörnmark,⁹⁷ T. Mkrtchyan,^{61a} M. Mlynarikova,¹⁴³ T. Moa,^{45a,45b} K. Mochizuki,¹¹⁰ P. Mogg,⁵²
 S. Mohapatra,³⁹ R. Moles-Valls,²⁴ M. C. Mondragon,¹⁰⁷ K. Mönig,⁴⁶ J. Monk,⁴⁰ E. Monnier,¹⁰² A. Montalbano,¹⁵²
 J. Montejo Berlingen,³⁶ M. Montella,⁹⁵ F. Monticelli,⁸⁹ S. Monzani,^{69a} N. Morange,⁶⁵ D. Moreno,²² M. Moreno Llácer,¹⁷⁴
 C. Moreno Martinez,¹⁴ P. Morettini,^{55b} M. Morgenstern,¹²⁰ S. Morgenstern,⁴⁸ D. Mori,¹⁵² M. Morii,⁵⁹ M. Morinaga,¹⁷⁹
 V. Morisbak,¹³⁴ A. K. Morley,³⁶ G. Mornacchi,³⁶ A. P. Morris,⁹⁵ L. Morvaj,¹⁵⁵ P. Moschovakos,³⁶ B. Moser,¹²⁰
 M. Mosidze,^{159b} T. Moskalets,¹⁴⁵ H. J. Moss,¹⁴⁹ J. Moss,^{31,hh} E. J. W. Moyses,¹⁰³ S. Muanza,¹⁰² J. Mueller,¹³⁹
 R. S. P. Mueller,¹¹⁴ D. Muenstermann,⁹⁰ G. A. Mullier,⁹⁷ D. P. Mungo,^{69a,69b} J. L. Munoz Martinez,¹⁴
 F. J. Munoz Sanchez,¹⁰¹ P. Murin,^{28b} W. J. Murray,^{178,144} A. Murrone,^{69a,69b} M. Muškinja,¹⁸ C. Mwewa,^{33a}
 A. G. Myagkov,^{123,ii} A. A. Myers,¹³⁹ J. Myers,¹³² M. Myska,¹⁴² B. P. Nachman,¹⁸ O. Nackenhorst,⁴⁷ A. Nag Nag,⁴⁸
 K. Nagai,¹³⁵ K. Nagano,⁸² Y. Nagasaka,⁶² J. L. Nagle,²⁹ E. Nagy,¹⁰² A. M. Nairz,³⁶ Y. Nakahama,¹¹⁷ K. Nakamura,⁸²
 T. Nakamura,¹⁶³ I. Nakano,¹²⁸ H. Nanjo,¹³³ F. Napolitano,^{61a} R. F. Naranjo Garcia,⁴⁶ R. Narayan,⁴² I. Naryshkin,¹³⁸
 T. Naumann,⁴⁶ G. Navarro,²² P. Y. Nechaeva,¹¹¹ F. Nechansky,⁴⁶ T. J. Neep,²¹ A. Negri,^{71a,71b} M. Negrini,^{23b} C. Nellist,⁵³
 M. E. Nelson,^{45a,45b} S. Nemecek,¹⁴¹ P. Nemethy,¹²⁵ M. Nessi,^{36,ij} M. S. Neubauer,¹⁷³ F. Neuhaus,¹⁰⁰ M. Neumann,¹⁸²
 R. Newhouse,¹⁷⁵ P. R. Newman,²¹ C. W. Ng,¹³⁹ Y. S. Ng,¹⁹ Y. W. Y. Ng,¹⁷¹ B. Ngair,^{35e} H. D. N. Nguyen,¹⁰²
 T. Nguyen Manh,¹¹⁰ E. Nibigira,³⁸ R. B. Nickerson,¹³⁵ R. Nicolaidou,¹⁴⁵ D. S. Nielsen,⁴⁰ J. Nielsen,¹⁴⁶ N. Nikiforou,¹¹
 V. Nikolaenko,^{123,ii} I. Nikolic-Audit,¹³⁶ K. Nikolopoulos,²¹ P. Nilsson,²⁹ H. R. Nindhito,⁵⁴ Y. Ninomiya,⁸² A. Nisati,^{73a}
 N. Nishu,^{60c} R. Nisius,¹¹⁵ I. Nitsche,⁴⁷ T. Nitta,¹⁷⁹ T. Nobe,¹⁶³ Y. Noguchi,⁸⁶ I. Nomidis,¹³⁶ M. A. Nomura,²⁹ M. Nordberg,³⁶
 N. Norjoharuddeen,¹³⁵ T. Novak,⁹² O. Novgorodova,⁴⁸ R. Novotny,¹⁴² L. Nozka,¹³¹ K. Ntekas,¹⁷¹ E. Nurse,⁹⁵
 F. G. Oakham,^{34,d} H. Oberlack,¹¹⁵ J. Ocariz,¹³⁶ A. Ochi,⁸³ I. Ochoa,³⁹ J. P. Ochoa-Ricoux,^{147a} K. O'Connor,²⁶ S. Oda,⁸⁸
 S. Odaka,⁸² S. Oerdek,⁵³ A. Ogrodnik,^{84a} A. Oh,¹⁰¹ S. H. Oh,⁴⁹ C. C. Ohm,¹⁵⁴ H. Oide,¹⁶⁵ M. L. Ojeda,¹⁶⁷ H. Okawa,¹⁶⁹
 Y. Okazaki,⁸⁶ M. W. O'Keefe,⁹¹ Y. Okumura,¹⁶³ T. Okuyama,⁸² A. Olariu,^{27b} L. F. Oleiro Seabra,^{140a} S. A. Olivares Pino,^{147a}
 D. Oliveira Damazio,²⁹ J. L. Oliver,¹ M. J. R. Olsson,¹⁷¹ A. Olszewski,⁸⁵ J. Olszowska,⁸⁵ D. C. O'Neil,¹⁵² A. P. O'Neill,¹³⁵
 A. Onofre,^{140a,140e} P. U. E. Onyisi,¹¹ H. Oppen,¹³⁴ M. J. Oreglia,³⁷ G. E. Orellana,⁸⁹ D. Orestano,^{75a,75b} N. Orlando,¹⁴
 R. S. Orr,¹⁶⁷ V. O'Shea,⁵⁷ R. Ospanov,^{60a} G. Otero y Garzon,³⁰ H. Otono,⁸⁸ P. S. Ott,^{61a} M. Ouchrif,^{35d} J. Ouellette,²⁹
 F. Ould-Saada,¹³⁴ A. Ouraou,¹⁴⁵ Q. Ouyang,^{15a} M. Owen,⁵⁷ R. E. Owen,²¹ V. E. Ozcan,^{12c} N. Ozturk,⁸ J. Pacalt,¹³¹
 H. A. Pacey,³² K. Pachal,⁴⁹ A. Pacheco Pages,¹⁴ C. Padilla Aranda,¹⁴ S. Pagan Griso,¹⁸ M. Paganini,¹⁸³ G. Palacino,⁶⁶
 S. Palazzo,⁵⁰ S. Palestini,³⁶ M. Palka,^{84b} D. Pallin,³⁸ I. Panagoulas,¹⁰ C. E. Pandini,³⁶ J. G. Panduro Vazquez,⁹⁴ P. Pani,⁴⁶
 G. Panizzo,^{67a,67c} L. Paolozzi,⁵⁴ C. Papadatos,¹¹⁰ K. Papageorgiou,^{9,s} S. Parajuli,⁴³ A. Paramonov,⁶ D. Paredes Hernandez,^{63b}
 S. R. Paredes Saenz,¹³⁵ B. Parida,¹⁶⁶ T. H. Park,¹⁶⁷ A. J. Parker,³¹ M. A. Parker,³² F. Parodi,^{55b,55a} E. W. Parrish,¹²¹

J. A. Parsons,³⁹ U. Parzefall,⁵² L. Pascual Dominguez,¹³⁶ V. R. Pascuzzi,¹⁶⁷ J. M. P. Pasner,¹⁴⁶ F. Pasquali,¹²⁰
 E. Pasqualucci,^{73a} S. Passaggio,^{55b} F. Pastore,⁹⁴ P. Pasuwan,^{45a,45b} S. Patarraia,¹⁰⁰ J. R. Pater,¹⁰¹ A. Pathak,^{181,e} J. Patton,⁹¹
 T. Pauly,³⁶ J. Pearkes,¹⁵³ B. Pearson,¹¹⁵ M. Pedersen,¹³⁴ L. Pedraza Diaz,¹¹⁹ R. Pedro,^{140a} T. Peiffer,⁵³
 S. V. Peleganchuk,^{122b,122a} O. Penc,¹⁴¹ H. Peng,^{60a} B. S. Peralva,^{81a} M. M. Perego,⁶⁵ A. P. Pereira Peixoto,^{140a}
 D. V. Perepelitsa,²⁹ F. Peri,¹⁹ L. Perini,^{69a,69b} H. Pernegger,³⁶ S. Perrella,^{140a} A. Perrevoort,¹²⁰ K. Peters,⁴⁶ R. F. Y. Peters,¹⁰¹
 B. A. Petersen,³⁶ T. C. Petersen,⁴⁰ E. Petit,¹⁰² A. Petridis,¹ C. Petridou,¹⁶² P. Petroff,⁶⁵ M. Petrov,¹³⁵ F. Petrucci,^{75a,75b}
 M. Pettee,¹⁸³ N. E. Pettersson,¹⁰³ K. Petukhova,¹⁴³ A. Peyaud,¹⁴⁵ R. Pezoa,^{147c} L. Pezzotti,^{71a,71b} T. Pham,¹⁰⁵
 F. H. Phillips,¹⁰⁷ P. W. Phillips,¹⁴⁴ M. W. Phipps,¹⁷³ G. Piacquadio,¹⁵⁵ E. Pianori,¹⁸ A. Picazio,¹⁰³ R. H. Pickles,¹⁰¹
 R. Piegaia,³⁰ D. Pietreanu,^{27b} J. E. Pilcher,³⁷ A. D. Pilkington,¹⁰¹ M. Pinamonti,^{67a,67c} J. L. Pinfold,³ M. Pitt,¹⁶¹
 L. Pizzimento,^{74a,74b} M.-A. Pleier,²⁹ V. Pleskot,¹⁴³ E. Plotnikova,⁸⁰ P. Podberezko,^{122b,122a} R. Poettgen,⁹⁷ R. Poggi,⁵⁴
 L. Poggioli,⁶⁵ I. Pogrebnyak,¹⁰⁷ D. Pohl,²⁴ I. Pokharel,⁵³ G. Polesello,^{71a} A. Poley,¹⁸ A. Policicchio,^{73a,73b} R. Polifka,¹⁴³
 A. Polini,^{23b} C. S. Pollard,⁴⁶ V. Polychronakos,²⁹ D. Ponomarenko,¹¹² L. Pontecorvo,³⁶ S. Popa,^{27a} G. A. Popeneciu,^{27d}
 L. Portales,⁵ D. M. Portillo Quintero,⁵⁸ S. Pospisil,¹⁴² K. Potamianos,⁴⁶ I. N. Potrap,⁸⁰ C. J. Potter,³² H. Potti,¹¹ T. Poulsen,⁹⁷
 J. Poveda,³⁶ T. D. Powell,¹⁴⁹ G. Pownall,⁴⁶ M. E. Pozo Astigarraga,³⁶ P. Pralavorio,¹⁰² S. Prell,⁷⁹ D. Price,¹⁰¹
 M. Primavera,^{68a} S. Prince,¹⁰⁴ M. L. Proffitt,¹⁴⁸ N. Proklova,¹¹² K. Prokofiev,^{63c} F. Prokoshin,⁸⁰ S. Protopopescu,²⁹
 J. Proudfoot,⁶ M. Przybycien,^{84a} D. Pudzha,¹³⁸ A. Puri,¹⁷³ P. Puzo,⁶⁵ J. Qian,¹⁰⁶ Y. Qin,¹⁰¹ A. Quadt,⁵³
 M. Queitsch-Maitland,³⁶ A. Qureshi,¹ M. Racko,^{28a} F. Ragusa,^{69a,69b} G. Rahal,⁹⁸ J. A. Raine,⁵⁴ S. Rajagopalan,²⁹
 A. Ramirez Morales,⁹³ K. Ran,^{15a,15d} T. Rashid,⁶⁵ S. Raspopov,⁵ D. M. Rauch,⁴⁶ F. Rauscher,¹¹⁴ S. Rave,¹⁰⁰ B. Ravina,¹⁴⁹
 I. Ravinovich,¹⁸⁰ J. H. Rawling,¹⁰¹ M. Raymond,³⁶ A. L. Read,¹³⁴ N. P. Readioff,⁵⁸ M. Reale,^{68a,68b} D. M. Rebuffi,^{71a,71b}
 A. Redelbach,¹⁷⁷ G. Redlinger,²⁹ K. Reeves,⁴³ L. Rehnisch,¹⁹ J. Reichert,¹³⁷ D. Reikher,¹⁶¹ A. Reiss,¹⁰⁰ A. Rej,¹⁵¹
 C. Rembser,³⁶ A. Renardi,⁴⁶ M. Renda,^{27b} M. Rescigno,^{73a} S. Resconi,^{69a} E. D. Resseguie,¹³⁷ S. Rettie,¹⁷⁵ B. Reynolds,¹²⁷
 E. Reynolds,²¹ O. L. Rezanova,^{122b,122a} P. Reznicek,¹⁴³ E. Ricci,^{76a,76b} R. Richter,¹¹⁵ S. Richter,⁴⁶ E. Richter-Was,^{84b}
 O. Ricken,²⁴ M. Ridel,¹³⁶ P. Rieck,¹¹⁵ O. Rifki,⁴⁶ M. Rijssenbeek,¹⁵⁵ A. Rimoldi,^{71a,71b} M. Rimoldi,⁴⁶ L. Rinaldi,^{23b}
 G. Ripellino,¹⁵⁴ I. Riu,¹⁴ J. C. Rivera Vergara,¹⁷⁶ F. Rizatdinova,¹³⁰ E. Rizvi,⁹³ C. Rizzi,³⁶ R. T. Roberts,¹⁰¹
 S. H. Robertson,^{104,m} M. Robin,⁴⁶ D. Robinson,³² J. E. M. Robinson,⁴⁶ C. M. Robles Gajardo,^{147c} M. Robles Manzano,¹⁰⁰
 A. Robson,⁵⁷ A. Rocchi,^{74a,74b} E. Rocco,¹⁰⁰ C. Roda,^{72a,72b} S. Rodriguez Bosca,¹⁷⁴ A. Rodriguez Perez,¹⁴
 D. Rodriguez Rodriguez,¹⁷⁴ A. M. Rodríguez Vera,^{168b} S. Roe,³⁶ O. Røhne,¹³⁴ R. Röhrig,¹¹⁵ R. A. Rojas,^{147c} B. Roland,⁵²
 C. P. A. Roland,⁶⁶ J. Roloff,²⁹ A. Romaniouk,¹¹² M. Romano,^{23b,23a} N. Rompotis,⁹¹ M. Ronzani,¹²⁵ L. Roos,¹³⁶ S. Rosati,^{73a}
 G. Rosin,¹⁰³ B. J. Rosser,¹³⁷ E. Rossi,⁴⁶ E. Rossi,^{75a,75b} E. Rossi,^{70a,70b} L. P. Rossi,^{55b} L. Rossini,^{69a,69b} R. Rosten,¹⁴
 M. Rotaru,^{27b} J. Rothberg,¹⁴⁸ B. Rottler,⁵² D. Rousseau,⁶⁵ G. Rovelli,^{71a,71b} A. Roy,¹¹ D. Roy,^{33d} A. Rozanov,¹⁰² Y. Rozen,¹⁶⁰
 X. Ruan,^{33d} F. Rühr,⁵² A. Ruiz-Martinez,¹⁷⁴ A. Rummler,³⁶ Z. Rurikova,⁵² N. A. Rusakovich,⁸⁰ H. L. Russell,¹⁰⁴
 L. Rustige,^{38,47} J. P. Rutherford,⁷ E. M. Rüttinger,¹⁴⁹ M. Rybar,³⁹ G. Rybkin,⁶⁵ E. B. Rye,¹³⁴ A. Ryzhov,¹²³
 J. A. Sabater Iglesias,⁴⁶ P. Sabatini,⁵³ G. Sabato,¹²⁰ S. Sacerdoti,⁶⁵ H. F.-W. Sadrozinski,¹⁴⁶ R. Sadykov,⁸⁰ F. Safai Tehrani,^{73a}
 B. Safarzadeh Samani,¹⁵⁶ M. Safdari,¹⁵³ P. Saha,¹²¹ S. Saha,¹⁰⁴ M. Sahinsoy,^{61a} A. Sahu,¹⁸² M. Saimpert,⁴⁶ M. Saito,¹⁶³
 T. Saito,¹⁶³ H. Sakamoto,¹⁶³ A. Sakharov,^{125,dd} D. Salamani,⁵⁴ G. Salamanna,^{75a,75b} J. E. Salazar Loyola,^{147c} A. Salnikov,¹⁵³
 J. Salt,¹⁷⁴ D. Salvatore,^{41b,41a} F. Salvatore,¹⁵⁶ A. Salvucci,^{63a,63b,63c} A. Salzburger,³⁶ J. Samarati,³⁶ D. Sammel,⁵²
 D. Sampsonidis,¹⁶² D. Sampsonidou,¹⁶² J. Sánchez,¹⁷⁴ A. Sanchez Pineda,^{67a,36,67c} H. Sandaker,¹³⁴ C. O. Sander,⁴⁶
 I. G. Sanderswood,⁹⁰ M. Sandhoff,¹⁸² C. Sandoval,²² D. P. C. Sankey,¹⁴⁴ M. Sannino,^{55b,55a} Y. Sano,¹¹⁷ A. Sansoni,⁵¹
 C. Santoni,³⁸ H. Santos,^{140a,140b} S. N. Santpur,¹⁸ A. Santra,¹⁷⁴ A. Saponov,⁸⁰ J. G. Saraiva,^{140a,140d} O. Sasaki,⁸² K. Sato,¹⁶⁹
 F. Sauerburger,⁵² E. Sauvan,⁵ P. Savard,^{167,d} R. Sawada,¹⁶³ C. Sawyer,¹⁴⁴ L. Sawyer,^{96,kk} C. Sbarra,^{23b} A. Sbrizzi,^{23a}
 T. Scanlon,⁹⁵ J. Schaarschmidt,¹⁴⁸ P. Schacht,¹¹⁵ B. M. Schachtner,¹¹⁴ D. Schaefer,³⁷ L. Schaefer,¹³⁷ J. Schaeffer,¹⁰⁰
 S. Schaepe,³⁶ U. Schäfer,¹⁰⁰ A. C. Schaffer,⁶⁵ D. Schaile,¹¹⁴ R. D. Schamberger,¹⁵⁵ N. Scharmberg,¹⁰¹ V. A. Schegelsky,¹³⁸
 D. Scheirich,¹⁴³ F. Schenck,¹⁹ M. Schernau,¹⁷¹ C. Schiavi,^{55b,55a} S. Schier,¹⁴⁶ L. K. Schildgen,²⁴ Z. M. Schillaci,²⁶
 E. J. Schioppa,³⁶ M. Schioppa,^{41b,41a} K. E. Schleicher,⁵² S. Schlenker,³⁶ K. R. Schmidt-Sommerfeld,¹¹⁵ K. Schmieden,³⁶
 C. Schmitt,¹⁰⁰ S. Schmitt,⁴⁶ S. Schmitz,¹⁰⁰ J. C. Schmoedel,⁴⁶ U. Schnoor,⁵² L. Schoeffel,¹⁴⁵ A. Schoening,^{61b}
 P. G. Scholer,⁵² E. Schopf,¹³⁵ M. Schott,¹⁰⁰ J. F. P. Schouwenberg,¹¹⁹ J. Schovancova,³⁶ S. Schramm,⁵⁴ F. Schroeder,¹⁸²
 A. Schulte,¹⁰⁰ H.-C. Schultz-Coulon,^{61a} M. Schumacher,⁵² B. A. Schumm,¹⁴⁶ Ph. Schune,¹⁴⁵ A. Schwartzman,¹⁵³
 T. A. Schwarz,¹⁰⁶ Ph. Schwemling,¹⁴⁵ R. Schwiendhorst,¹⁰⁷ A. Sciandra,¹⁴⁶ G. Sciolla,²⁶ M. Scodreggio,⁴⁶
 M. Scornajenghi,^{41b,41a} F. Scuri,^{72a} F. Scutti,¹⁰⁵ L. M. Scyboz,¹¹⁵ C. D. Sebastiani,^{73a,73b} P. Seema,¹⁹ S. C. Seidel,¹¹⁸

A. Seiden,¹⁴⁶ B. D. Seidlitz,²⁹ T. Seiss,³⁷ J. M. Seixas,^{81b} G. Sekhniaidze,^{70a} K. Sekhon,¹⁰⁶ S. J. Sekula,⁴²
 N. Semprini-Cesari,^{23b,23a} S. Sen,⁴⁹ C. Serfon,⁷⁷ L. Serin,⁶⁵ L. Serkin,^{67a,67b} M. Sessa,^{60a} H. Severini,¹²⁹ S. Sevova,¹⁵³
 T. Šfiligoj,⁹² F. Sforza,^{55b,55a} A. Sfyrla,⁵⁴ E. Shabalina,⁵³ J. D. Shahinian,¹⁴⁶ N. W. Shaikh,^{45a,45b} D. Shaked Renous,¹⁸⁰
 L. Y. Shan,^{15a} J. T. Shank,²⁵ M. Shapiro,¹⁸ A. Sharma,¹³⁵ A. S. Sharma,¹ P. B. Shatalov,¹²⁴ K. Shaw,¹⁵⁶ S. M. Shaw,¹⁰¹
 M. Shehade,¹⁸⁰ Y. Shen,¹²⁹ A. D. Sherman,²⁵ P. Sherwood,⁹⁵ L. Shi,¹⁵⁸ S. Shimizu,⁸² C. O. Shimmin,¹⁸³ Y. Shimogama,¹⁷⁹
 M. Shimojima,¹¹⁶ I. P. J. Shipsey,¹³⁵ S. Shirabe,¹⁶⁵ M. Shiyakova,^{80,11} J. Shlomi,¹⁸⁰ A. Shmeleva,¹¹¹ M. J. Shochet,³⁷
 J. Shojaii,¹⁰⁵ D. R. Shope,¹²⁹ S. Shrestha,¹²⁷ E. M. Shrif,^{33d} E. Shulga,¹⁸⁰ P. Sicho,¹⁴¹ A. M. Sickles,¹⁷³ P. E. Sidebo,¹⁵⁴
 E. Sideras Haddad,^{33d} O. Sidiropoulou,³⁶ A. Sidoti,^{23b,23a} F. Siegert,⁴⁸ Dj. Sijacki,¹⁶ M. Silva Jr.,¹⁸¹ M. V. Silva Oliveira,^{81a}
 S. B. Silverstein,^{45a} S. Simion,⁶⁵ R. Simoniello,¹⁰⁰ S. Simsek,^{12b} P. Sinervo,¹⁶⁷ V. Sinetckii,¹¹³ N. B. Sinev,¹³² S. Singh,¹⁵²
 M. Sioli,^{23b,23a} I. Siral,¹³² S. Yu. Sivoklov,¹¹³ J. Sjölin,^{45a,45b} E. Skorda,⁹⁷ P. Skubic,¹²⁹ M. Slawinska,⁸⁵ K. Sliwa,¹⁷⁰
 R. Slovak,¹⁴³ V. Smakhtin,¹⁸⁰ B. H. Smart,¹⁴⁴ J. Smiesko,^{28a} N. Smirnov,¹¹² S. Yu. Smirnov,¹¹² Y. Smirnov,¹¹²
 L. N. Smirnova,^{113,mm} O. Smirnova,⁹⁷ J. W. Smith,⁵³ M. Smizanska,⁹⁰ K. Smolek,¹⁴² A. Smykiewicz,⁸⁵ A. A. Snesev,¹¹¹
 H. L. Snoek,¹²⁰ I. M. Snyder,¹³² S. Snyder,²⁹ R. Sobie,^{176,m} A. Soffer,¹⁶¹ A. Sjøgaard,⁵⁰ F. Sohns,⁵³ C. A. Solans Sanchez,³⁶
 E. Yu. Soldatov,¹¹² U. Soldevila,¹⁷⁴ A. A. Solodkov,¹²³ A. Soloshenko,⁸⁰ O. V. Solovyanov,¹²³ V. Solovyeu,¹³⁸ P. Sommer,¹⁴⁹
 H. Son,¹⁷⁰ W. Song,¹⁴⁴ W. Y. Song,^{168b} A. Sopczak,¹⁴² A. L. Soppio,⁹⁵ F. Sopkova,^{28b} C. L. Sotiropoulou,^{72a,72b}
 S. Sottocornola,^{71a,71b} R. Soualah,^{67a,67c,mm} A. M. Soukharev,^{122b,122a} D. South,⁴⁶ S. Spagnolo,^{68a,68b} M. Spalla,¹¹⁵
 M. Spangenberg,¹⁷⁸ F. Spanò,⁹⁴ D. Sperlich,⁵² T. M. Spieker,^{61a} G. Spigo,³⁶ M. Spina,¹⁵⁶ D. P. Spiteri,⁵⁷ M. Spousta,¹⁴³
 A. Stabile,^{69a,69b} B. L. Stamas,¹²¹ R. Stamen,^{61a} M. Stamenkovic,¹²⁰ E. Stanecka,⁸⁵ B. Stanislaus,¹³⁵ M. M. Stanitzki,⁴⁶
 M. Stankaityte,¹³⁵ B. Stapf,¹²⁰ E. A. Starchenko,¹²³ G. H. Stark,¹⁴⁶ J. Stark,⁵⁸ P. Staroba,¹⁴¹ P. Starovoitov,^{61a} S. Stärz,¹⁰⁴
 R. Staszewski,⁸⁵ G. Stavropoulos,⁴⁴ M. Stegler,⁴⁶ P. Steinberg,²⁹ A. L. Steinhebel,¹³² B. Stelzer,¹⁵² H. J. Stelzer,¹³⁹
 O. Stelzer-Chilton,^{168a} H. Stenzel,⁵⁶ T. J. Stevenson,¹⁵⁶ G. A. Stewart,³⁶ M. C. Stockton,³⁶ G. Stoicea,^{27b} M. Stolarski,^{140a}
 S. Stonjek,¹¹⁵ A. Straessner,⁴⁸ J. Strandberg,¹⁵⁴ S. Strandberg,^{45a,45b} M. Strauss,¹²⁹ P. Strizenec,^{28b} R. Ströhmer,¹⁷⁷
 D. M. Strom,¹³² R. Stroynowski,⁴² A. Strubig,⁵⁰ S. A. Stucci,²⁹ B. Stugu,¹⁷ J. Stupak,¹²⁹ N. A. Styles,⁴⁶ D. Su,¹⁵³ W. Su,^{60c}
 S. Suchek,^{61a} V. V. Sulini,¹¹¹ M. J. Sullivan,⁹¹ D. M. S. Sultan,⁵⁴ S. Sultansoy,^{4c} T. Sumida,⁸⁶ S. Sun,¹⁰⁶ X. Sun,³
 K. Suruliz,¹⁵⁶ C. J. E. Suster,¹⁵⁷ M. R. Sutton,¹⁵⁶ S. Suzuki,⁸² M. Svatos,¹⁴¹ M. Swiatlowski,³⁷ S. P. Swift,² T. Swirski,¹⁷⁷
 A. Sydorenko,¹⁰⁰ I. Sykora,^{28a} M. Sykora,¹⁴³ T. Sykora,¹⁴³ D. Ta,¹⁰⁰ K. Tackmann,^{46,oo} J. Taenzer,¹⁶¹ A. Taffard,¹⁷¹
 R. Tafirout,^{168a} H. Takai,²⁹ R. Takashima,⁸⁷ K. Takeda,⁸³ T. Takeshita,¹⁵⁰ E. P. Takeva,⁵⁰ Y. Takubo,⁸² M. Talby,¹⁰²
 A. A. Talyshev,^{122b,122a} N. M. Tamir,¹⁶¹ J. Tanaka,¹⁶³ M. Tanaka,¹⁶⁵ R. Tanaka,⁶⁵ S. Tapia Araya,¹⁷³ S. Tapprogge,¹⁰⁰
 A. Tarek Abouelfadl Mohamed,¹³⁶ S. Tarem,¹⁶⁰ K. Tariq,^{60b} G. Tarna,^{27b,pp} G. F. Tartarelli,^{69a} P. Tas,¹⁴³ M. Tasevsky,¹⁴¹
 T. Tashiro,⁸⁶ E. Tassi,^{41b,41a} A. Tavares Delgado,^{140a} Y. Tayalati,^{35e} A. J. Taylor,⁵⁰ G. N. Taylor,¹⁰⁵ W. Taylor,^{168b} A. S. Tee,⁹⁰
 R. Teixeira De Lima,¹⁵³ P. Teixeira-Dias,⁹⁴ H. Ten Kate,³⁶ J. J. Teoh,¹²⁰ S. Terada,⁸² K. Terashi,¹⁶³ J. Terron,⁹⁹ S. Terzo,¹⁴
 M. Testa,⁵¹ R. J. Teuscher,^{167,m} S. J. Thais,¹⁸³ T. Thevenaux-Pelzer,⁴⁶ F. Thiele,⁴⁰ D. W. Thomas,⁹⁴ J. O. Thomas,⁴²
 J. P. Thomas,²¹ P. D. Thompson,²¹ L. A. Thomsen,¹⁸³ E. Thomson,¹³⁷ E. J. Thorpe,⁹³ R. E. Ticse Torres,⁵³
 V. O. Tikhomirov,^{111,qq} Yu. A. Tikhonov,^{122b,122a} S. Timoshenko,¹¹² P. Tipton,¹⁸³ S. Tisserant,¹⁰² K. Todome,^{23b,23a}
 S. Todorova-Nova,⁵ S. Todt,⁴⁸ J. Tojo,⁸⁸ S. Tokár,^{28a} K. Tokushuku,⁸² E. Tolley,¹²⁷ K. G. Tomiwa,^{33d} M. Tomoto,¹¹⁷
 L. Tompkins,^{153,cc} B. Tong,⁵⁹ P. Tornambe,¹⁰³ E. Torrence,¹³² H. Torres,⁴⁸ E. Torró Pastor,¹⁴⁸ C. Toscirri,¹³⁵ J. Toth,^{102,rr}
 D. R. Tovey,¹⁴⁹ A. Traet,¹⁷ C. J. Treado,¹²⁵ T. Trefzger,¹⁷⁷ F. Tresoldi,¹⁵⁶ A. Tricoli,²⁹ I. M. Trigger,^{168a}
 S. Trincaz-Duvoid,¹³⁶ D. T. Trischuk,¹⁷⁵ W. Trischuk,¹⁶⁷ B. Trocmé,⁵⁸ A. Trofymov,¹⁴⁵ C. Troncon,^{69a} M. Trovatelli,¹⁷⁶
 F. Trovato,¹⁵⁶ L. Truong,^{33b} M. Trzebinski,⁸⁵ A. Trzupek,⁸⁵ F. Tsai,⁴⁶ J. C-L. Tseng,¹³⁵ P. V. Tsiarehka,^{108,ee}
 A. Tsirigotis,^{162,ff} V. Tsiskaridze,¹⁵⁵ E. G. Tskhadadze,^{159a} M. Tsopoulou,¹⁶² I. I. Tsukerman,¹²⁴ V. Tsulaia,¹⁸ S. Tsuno,⁸²
 D. Tsybychev,¹⁵⁵ Y. Tu,^{63b} A. Tudorache,^{27b} V. Tudorache,^{27b} T. T. Tulbure,^{27a} A. N. Tuna,⁵⁹ S. Turchikhin,⁸⁰
 D. Turgeman,¹⁸⁰ I. Turk Cakir,^{4b,ss} R. J. Turner,²¹ R. T. Turra,^{69a} P. M. Tuts,³⁹ S. Tzamarias,¹⁶² E. Tzovara,¹⁰⁰ G. Ucchielli,⁴⁷
 K. Uchida,¹⁶³ I. Ueda,⁸² F. Ukegawa,¹⁶⁹ G. Unal,³⁶ A. Undrus,²⁹ G. Unel,¹⁷¹ F. C. Ungaro,¹⁰⁵ Y. Unno,⁸² K. Uno,¹⁶³
 J. Urban,^{28b} P. Urquijo,¹⁰⁵ G. Usai,⁸ Z. Uysal,^{12d} V. Vacek,¹⁴² B. Vachon,¹⁰⁴ K. O. H. Vadla,¹³⁴ A. Vaidya,⁹⁵ C. Valderanis,¹¹⁴
 E. Valdes Santurio,^{45a,45b} M. Valente,⁵⁴ S. Valentinetti,^{23b,23a} A. Valero,¹⁷⁴ L. Valéry,⁴⁶ R. A. Vallance,²¹ A. Vallier,³⁶
 J. A. Valls Ferrer,¹⁷⁴ T. R. Van Daalen,¹⁴ P. Van Gemmeren,⁶ I. Van Vulpen,¹²⁰ M. Vanadia,^{74a,74b} W. Vandelli,³⁶
 M. Vandenbroucke,¹⁴⁵ E. R. Vandewall,¹³⁰ A. Vaniachine,¹⁶⁶ D. Vannicola,^{73a,73b} R. Vari,^{73a} E. W. Varnes,⁷ C. Varni,^{55b,55a}
 T. Varol,¹⁵⁸ D. Varouchas,⁶⁵ K. E. Varvell,¹⁵⁷ M. E. Vasile,^{27b} G. A. Vasquez,¹⁷⁶ F. Vazeille,³⁸ D. Vazquez Furelos,¹⁴
 T. Vazquez Schroeder,³⁶ J. Veatch,⁵³ V. Vecchio,^{75a,75b} M. J. Veen,¹²⁰ L. M. Veloce,¹⁶⁷ F. Veloso,^{140a,140c} S. Veneziano,^{73a}

A. Ventura,^{68a,68b} N. Venturi,³⁶ A. Verbytskyi,¹¹⁵ V. Vercesi,^{71a} M. Verducci,^{72a,72b} C. M. Vergel Infante,⁷⁹ C. Vergis,²⁴ W. Verkerke,¹²⁰ A. T. Vermeulen,¹²⁰ J. C. Vermeulen,¹²⁰ M. C. Vetterli,^{152,d} N. Viaux Maira,^{147c} M. Vicente Barreto Pinto,⁵⁴ T. Vickey,¹⁴⁹ O. E. Vickey Boeriu,¹⁴⁹ G. H. A. Viehhauser,¹³⁵ L. Vignani,^{61b} M. Villa,^{23b,23a} M. Villaplana Perez,³ E. Vilucchi,⁵¹ M. G. Vincker,³⁴ G. S. Virdee,²¹ A. Vishwakarma,⁴⁶ C. Vittori,^{23b,23a} I. Vivarelli,¹⁵⁶ M. Vogel,¹⁸² P. Vokac,¹⁴² S. E. von Buddenbrock,^{33d} E. Von Toerne,²⁴ V. Vorobel,¹⁴³ K. Vorobev,¹¹² M. Vos,¹⁷⁴ J. H. Vosseveld,⁹¹ M. Vozak,¹⁰¹ N. Vranjes,¹⁶ M. Vranjes Milosavljevic,¹⁶ V. Vrba,¹⁴² M. Vreeswijk,¹²⁰ R. Vuillermet,³⁶ I. Vukotic,³⁷ P. Wagner,²⁴ W. Wagner,¹⁸² J. Wagner-Kuhr,¹¹⁴ S. Wahdan,¹⁸² H. Wahlberg,⁸⁹ V. M. Walbrecht,¹¹⁵ J. Walder,⁹⁰ R. Walker,¹¹⁴ S. D. Walker,⁹⁴ W. Walkowiak,¹⁵¹ V. Wallangen,^{45a,45b} A. M. Wang,⁵⁹ A. Z. Wang,¹⁸¹ C. Wang,^{60c} F. Wang,¹⁸¹ H. Wang,¹⁸ H. Wang,³ J. Wang,^{63a} J. Wang,^{61b} P. Wang,⁴² Q. Wang,¹²⁹ R.-J. Wang,¹⁰⁰ R. Wang,^{60a} R. Wang,⁶ S. M. Wang,¹⁵⁸ W. T. Wang,^{60a} W. Wang,^{15c} W. X. Wang,^{60a} Y. Wang,^{60a} Z. Wang,^{60c} C. Wanotayaroj,⁴⁶ A. Warburton,¹⁰⁴ C. P. Ward,³² D. R. Wardrope,⁹⁵ N. Warrack,⁵⁷ A. Washbrook,⁵⁰ A. T. Watson,²¹ M. F. Watson,²¹ G. Watts,¹⁴⁸ B. M. Waugh,⁹⁵ A. F. Webb,¹¹ S. Webb,¹⁰⁰ C. Weber,¹⁸³ M. S. Weber,²⁰ S. A. Weber,³⁴ S. M. Weber,^{61a} A. R. Weidberg,¹³⁵ J. Weingarten,⁴⁷ M. Weirich,¹⁰⁰ C. Weiser,⁵² P. S. Wells,³⁶ T. Wenaus,²⁹ T. Wengler,³⁶ S. Wenig,³⁶ N. Vermes,²⁴ M. D. Werner,⁷⁹ M. Wessels,^{61a} T. D. Weston,²⁰ K. Whalen,¹³² N. L. Whallon,¹⁴⁸ A. M. Wharton,⁹⁰ A. S. White,¹⁰⁶ A. White,⁸ M. J. White,¹ D. Whiteson,¹⁷¹ B. W. Whitmore,⁹⁰ W. Wiedenmann,¹⁸¹ C. Wiel,⁴⁸ M. Wielers,¹⁴⁴ N. Wieseotte,¹⁰⁰ C. Wiglesworth,⁴⁰ L. A. M. Wiik-Fuchs,⁵² H. G. Wilkens,³⁶ L. J. Wilkins,⁹⁴ H. H. Williams,¹³⁷ S. Williams,³² C. Willis,¹⁰⁷ S. Willocq,¹⁰³ J. A. Wilson,²¹ I. Wingerter-Seez,⁵ E. Winkels,¹⁵⁶ F. Winklmeier,¹³² O. J. Winston,¹⁵⁶ B. T. Winter,⁵² M. Wittgen,¹⁵³ M. Wobisch,⁹⁶ A. Wolf,¹⁰⁰ T. M. H. Wolf,¹²⁰ R. Wolff,¹⁰² R. W. Wölker,¹³⁵ J. Wollrath,⁵² M. W. Wolter,⁸⁵ H. Wolters,^{140a,140c} V. W. S. Wong,¹⁷⁵ N. L. Woods,¹⁴⁶ S. D. Worm,²¹ B. K. Wosiek,⁸⁵ K. W. Woźniak,⁸⁵ K. Wraight,⁵⁷ S. L. Wu,¹⁸¹ X. Wu,⁵⁴ Y. Wu,^{60a} T. R. Wyatt,¹⁰¹ B. M. Wynne,⁵⁰ S. Xella,⁴⁰ Z. Xi,¹⁰⁶ L. Xia,¹⁷⁸ X. Xiao,¹⁰⁶ I. Xiotidis,¹⁵⁶ D. Xu,^{15a} H. Xu,^{60a} H. Xu,^{60a} L. Xu,²⁹ T. Xu,¹⁴⁵ W. Xu,¹⁰⁶ Z. Xu,^{60b} Z. Xu,¹⁵³ B. Yabsley,¹⁵⁷ S. Yacoob,^{33a} K. Yajima,¹³³ D. P. Yallup,⁹⁵ N. Yamaguchi,⁸⁸ Y. Yamaguchi,¹⁶⁵ A. Yamamoto,⁸² M. Yamatani,¹⁶³ T. Yamazaki,¹⁶³ Y. Yamazaki,⁸³ Z. Yan,²⁵ H. J. Yang,^{60c,60d} H. T. Yang,¹⁸ S. Yang,^{60a} T. Yang,^{63c} X. Yang,^{60b,58} Y. Yang,¹⁶³ W.-M. Yao,¹⁸ Y. C. Yap,⁴⁶ Y. Yasu,⁸² E. Yatsenko,^{60c,60d} H. Ye,^{15c} J. Ye,⁴² S. Ye,²⁹ I. Yeletsikh,⁸⁰ M. R. Yexley,⁹⁰ E. Yigitbasi,²⁵ K. Yorita,¹⁷⁹ K. Yoshihara,¹³⁷ C. J. S. Young,³⁶ C. Young,¹⁵³ J. Yu,⁷⁹ R. Yuan,^{60b,tt} X. Yue,^{61a} S. P. Y. Yuen,²⁴ M. Zaazoua,^{35e} B. Zabinski,⁸⁵ G. Zacharis,¹⁰ E. Zaffaroni,⁵⁴ J. Zahreddine,¹³⁶ A. M. Zaitsev,^{123,ii} T. Zakareishvili,^{159b} N. Zakharchuk,³⁴ S. Zambito,⁵⁹ D. Zanzi,³⁶ D. R. Zaripovas,⁵⁷ S. V. Zeiβner,⁴⁷ C. Zeitnitz,¹⁸² G. Zemaityte,¹³⁵ J. C. Zeng,¹⁷³ O. Zenin,¹²³ T. Ženiš,^{28a} D. Zerwas,⁶⁵ M. Zgubič,¹³⁵ B. Zhang,^{15c} D. F. Zhang,^{15b} G. Zhang,^{15b} H. Zhang,^{15c} J. Zhang,⁶ L. Zhang,^{15c} L. Zhang,^{60a} M. Zhang,¹⁷³ R. Zhang,¹⁸¹ S. Zhang,¹⁰⁶ X. Zhang,^{60b} Y. Zhang,^{15a,15d} Z. Zhang,^{63a} Z. Zhang,⁶⁵ P. Zhao,⁴⁹ Y. Zhao,^{60b} Z. Zhao,^{60a} A. Zhemchugov,⁸⁰ Z. Zheng,¹⁰⁶ D. Zhong,¹⁷³ B. Zhou,¹⁰⁶ C. Zhou,¹⁸¹ M. S. Zhou,^{15a,15d} M. Zhou,¹⁵⁵ N. Zhou,^{60c} Y. Zhou,⁷ C. G. Zhu,^{60b} C. Zhu,^{15a,15d} H. L. Zhu,^{60a} H. Zhu,^{15a} J. Zhu,¹⁰⁶ Y. Zhu,^{60a} X. Zhuang,^{15a} K. Zhukov,¹¹¹ V. Zhulanov,^{122b,122a} D. Zieminska,⁶⁶ N. I. Zimine,⁸⁰ S. Zimmermann,⁵² Z. Zinonos,¹¹⁵ M. Ziolkowski,¹⁵¹ L. Živković,¹⁶ G. Zobernig,¹⁸¹ A. Zoccoli,^{23b,23a} K. Zoch,⁵³ T. G. Zorbas,¹⁴⁹ R. Zou,³⁷ and L. Zwalinski³⁶

(ATLAS Collaboration)

¹Department of Physics, University of Adelaide, Adelaide, Australia²Physics Department, SUNY Albany, Albany New York, USA³Department of Physics, University of Alberta, Edmonton AB, Canada^{4a}Department of Physics, Ankara University, Ankara, Turkey^{4b}Istanbul Aydin University, Istanbul, Turkey^{4c}Division of Physics, TOBB University of Economics and Technology, Ankara, Turkey⁵LAPP, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy, France⁶High Energy Physics Division, Argonne National Laboratory, Argonne Illinois, USA⁷Department of Physics, University of Arizona, Tucson Arizona, USA⁸Department of Physics, University of Texas at Arlington, Arlington Texas, USA⁹Physics Department, National and Kapodistrian University of Athens, Athens, Greece¹⁰Physics Department, National Technical University of Athens, Zografou, Greece¹¹Department of Physics, University of Texas at Austin, Austin Texas, USA^{12a}Bahcesehir University, Faculty of Engineering and Natural Sciences, Istanbul, Turkey^{12b}Istanbul Bilgi University, Faculty of Engineering and Natural Sciences, Istanbul, Turkey^{12c}Department of Physics, Bogazici University, Istanbul, Turkey

- ^{12d}*Department of Physics Engineering, Gaziantep University, Gaziantep, Turkey*
- ¹³*Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan*
- ¹⁴*Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona, Spain*
- ^{15a}*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China*
- ^{15b}*Physics Department, Tsinghua University, Beijing, China*
- ^{15c}*Department of Physics, Nanjing University, Nanjing, China*
- ^{15d}*University of Chinese Academy of Science (UCAS), Beijing, China*
- ¹⁶*Institute of Physics, University of Belgrade, Belgrade, Serbia*
- ¹⁷*Department for Physics and Technology, University of Bergen, Bergen, Norway*
- ¹⁸*Physics Division, Lawrence Berkeley National Laboratory and University of California, Berkeley California, USA*
- ¹⁹*Institut für Physik, Humboldt Universität zu Berlin, Berlin, Germany*
- ²⁰*Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern, Switzerland*
- ²¹*School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*
- ²²*Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogota, Colombia*
- ^{23a}*INFN Bologna and Università di Bologna, Dipartimento di Fisica, Italy*
- ^{23b}*INFN Sezione di Bologna, Italy*
- ²⁴*Physikalisches Institut, Universität Bonn, Bonn, Germany*
- ²⁵*Department of Physics, Boston University, Boston Massachusetts, USA*
- ²⁶*Department of Physics, Brandeis University, Waltham Massachusetts, USA*
- ^{27a}*Transilvania University of Brasov, Brasov, Romania*
- ^{27b}*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania*
- ^{27c}*Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi, Romania*
- ^{27d}*National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca, Romania*
- ^{27e}*University Politehnica Bucharest, Bucharest, Romania*
- ^{27f}*West University in Timisoara, Timisoara, Romania*
- ^{28a}*Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovak Republic*
- ^{28b}*Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice, Slovak Republic*
- ²⁹*Physics Department, Brookhaven National Laboratory, Upton New York, USA*
- ³⁰*Departamento de Física, Universidad de Buenos Aires, Buenos Aires, Argentina*
- ³¹*California State University, California, USA*
- ³²*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ^{33a}*Department of Physics, University of Cape Town, Cape Town, South Africa*
- ^{33b}*Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg, South Africa*
- ^{33c}*University of South Africa, Department of Physics, Pretoria, South Africa*
- ^{33d}*School of Physics, University of the Witwatersrand, Johannesburg, South Africa*
- ³⁴*Department of Physics, Carleton University, Ottawa ON, Canada*
- ^{35a}*Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies—Université Hassan II, Casablanca, Morocco*
- ^{35b}*Faculté des Sciences, Université Ibn-Tofail, Kénitra, Morocco*
- ^{35c}*Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech, Morocco*
- ^{35d}*Faculté des Sciences, Université Mohamed Premier and LPTPM, Oujda, Morocco*
- ^{35e}*Faculté des sciences, Université Mohammed V, Rabat, Morocco*
- ³⁶*CERN, Geneva, Switzerland*
- ³⁷*Enrico Fermi Institute, University of Chicago, Chicago Illinois, USA*
- ³⁸*LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand, France*
- ³⁹*Nevis Laboratory, Columbia University, Irvington New York, USA*
- ⁴⁰*Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark*
- ^{41a}*Dipartimento di Fisica, Università della Calabria, Rende, Italy*
- ^{41b}*INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati, Italy*
- ⁴²*Physics Department, Southern Methodist University, Dallas Texas, USA*
- ⁴³*Physics Department, University of Texas at Dallas, Richardson Texas, USA*
- ⁴⁴*National Centre for Scientific Research “Demokritos”, Agia Paraskevi, Greece*
- ^{45a}*Department of Physics, Stockholm University, Sweden*
- ^{45b}*Oskar Klein Centre, Stockholm, Sweden*

- ⁴⁶Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen, Germany
- ⁴⁷Lehrstuhl für Experimentelle Physik IV, Technische Universität Dortmund, Dortmund, Germany
- ⁴⁸Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden, Germany
- ⁴⁹Department of Physics, Duke University, Durham North Carolina, USA
- ⁵⁰SUPA—School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
- ⁵¹INFN e Laboratori Nazionali di Frascati, Frascati, Italy
- ⁵²Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg, Germany
- ⁵³II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany
- ⁵⁴Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève, Switzerland
- ^{55a}Dipartimento di Fisica, Università di Genova, Genova, Italy
- ^{55b}INFN Sezione di Genova, Italy
- ⁵⁶II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen, Germany
- ⁵⁷SUPA—School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
- ⁵⁸LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble, France
- ⁵⁹Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge Massachusetts, USA
- ^{60a}Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei, China
- ^{60b}Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao, China
- ^{60c}School of Physics and Astronomy, Shanghai Jiao Tong University, KLPPAC-MoE, SKLPPC, Shanghai, China
- ^{60d}Tsung-Dao Lee Institute, Shanghai, China
- ^{61a}Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ^{61b}Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ⁶²Faculty of Applied Information Science, Hiroshima Institute of Technology, Hiroshima, Japan
- ^{63a}Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong, China
- ^{63b}Department of Physics, University of Hong Kong, Hong Kong, China
- ^{63c}Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China
- ⁶⁴Department of Physics, National Tsing Hua University, Hsinchu, Taiwan
- ⁶⁵IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay, France
- ⁶⁶Department of Physics, Indiana University, Bloomington Indiana, USA
- ^{67a}INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine, Italy
- ^{67b}ICTP, Trieste, Italy
- ^{67c}Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine, Italy
- ^{68a}INFN Sezione di Lecce, Italy
- ^{68b}Dipartimento di Matematica e Fisica, Università del Salento, Lecce, Italy
- ^{69a}INFN Sezione di Milano, Italy
- ^{69b}Dipartimento di Fisica, Università di Milano, Milano, Italy
- ^{70a}INFN Sezione di Napoli, Italy
- ^{70b}Dipartimento di Fisica, Università di Napoli, Napoli, Italy
- ^{71a}INFN Sezione di Pavia, Italy
- ^{71b}Dipartimento di Fisica, Università di Pavia, Pavia, Italy
- ^{72a}INFN Sezione di Pisa, Italy
- ^{72b}Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa, Italy
- ^{73a}INFN Sezione di Roma, Italy
- ^{73b}Dipartimento di Fisica, Sapienza Università di Roma, Roma, Italy
- ^{74a}INFN Sezione di Roma Tor Vergata, Italy
- ^{74b}Dipartimento di Fisica, Università di Roma Tor Vergata, Roma, Italy
- ^{75a}INFN Sezione di Roma Tre, Italy
- ^{75b}Dipartimento di Matematica e Fisica, Università Roma Tre, Roma, Italy
- ^{76a}INFN-TIFPA, Italy
- ^{76b}Università degli Studi di Trento, Trento, Italy
- ⁷⁷Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität, Innsbruck, Austria
- ⁷⁸University of Iowa, Iowa City Iowa, USA
- ⁷⁹Department of Physics and Astronomy, Iowa State University, Ames Iowa, USA
- ⁸⁰Joint Institute for Nuclear Research, Dubna, Russia
- ^{81a}Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora, Brazil
- ^{81b}Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro, Brazil

- ^{81c}*Universidade Federal de São João del Rei (UFSJ), São João del Rei, Brazil*
- ^{81d}*Instituto de Física, Universidade de São Paulo, São Paulo, Brazil*
- ⁸²*KEK, High Energy Accelerator Research Organization, Tsukuba, Japan*
- ⁸³*Graduate School of Science, Kobe University, Kobe, Japan*
- ^{84a}*AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow, Poland*
- ^{84b}*Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow, Poland*
- ⁸⁵*Institute of Nuclear Physics Polish Academy of Sciences, Krakow, Poland*
- ⁸⁶*Faculty of Science, Kyoto University, Kyoto, Japan*
- ⁸⁷*Kyoto University of Education, Kyoto, Japan*
- ⁸⁸*Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka, Japan*
- ⁸⁹*Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata, Argentina*
- ⁹⁰*Physics Department, Lancaster University, Lancaster, United Kingdom*
- ⁹¹*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁹²*Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana, Slovenia*
- ⁹³*School of Physics and Astronomy, Queen Mary University of London, London, United Kingdom*
- ⁹⁴*Department of Physics, Royal Holloway University of London, Egham, United Kingdom*
- ⁹⁵*Department of Physics and Astronomy, University College London, London, United Kingdom*
- ⁹⁶*Louisiana Tech University, Ruston Louisiana, USA*
- ⁹⁷*Fysiska institutionen, Lunds universitet, Lund, Sweden*
- ⁹⁸*Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules (IN2P3), Villeurbanne, France*
- ⁹⁹*Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid, Spain*
- ¹⁰⁰*Institut für Physik, Universität Mainz, Mainz, Germany*
- ¹⁰¹*School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ¹⁰²*CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France*
- ¹⁰³*Department of Physics, University of Massachusetts, Amherst Massachusetts, USA*
- ¹⁰⁴*Department of Physics, McGill University, Montreal QC, Canada*
- ¹⁰⁵*School of Physics, University of Melbourne, Victoria, Australia*
- ¹⁰⁶*Department of Physics, University of Michigan, Ann Arbor Michigan, USA*
- ¹⁰⁷*Department of Physics and Astronomy, Michigan State University, East Lansing Michigan, USA*
- ¹⁰⁸*B.I. Stepanov Institute of Physics, National Academy of Sciences of Belarus, Minsk, Belarus*
- ¹⁰⁹*Research Institute for Nuclear Problems of Byelorussian State University, Minsk, Belarus*
- ¹¹⁰*Group of Particle Physics, University of Montreal, Montreal QC, Canada*
- ¹¹¹*P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow, Russia*
- ¹¹²*National Research Nuclear University MEPhI, Moscow, Russia*
- ¹¹³*D.V. Skobeltsyn Institute of Nuclear Physics, M.V. Lomonosov Moscow State University, Moscow, Russia*
- ¹¹⁴*Fakultät für Physik, Ludwig-Maximilians-Universität München, München, Germany*
- ¹¹⁵*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München, Germany*
- ¹¹⁶*Nagasaki Institute of Applied Science, Nagasaki, Japan*
- ¹¹⁷*Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya, Japan*
- ¹¹⁸*Department of Physics and Astronomy, University of New Mexico, Albuquerque New Mexico, USA*
- ¹¹⁹*Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen, Netherlands*
- ¹²⁰*Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam, Netherlands*
- ¹²¹*Department of Physics, Northern Illinois University, DeKalb Illinois, USA*
- ^{122a}*Budker Institute of Nuclear Physics and NSU, SB RAS, Novosibirsk, Russia*
- ^{122b}*Novosibirsk State University Novosibirsk, Russia*
- ¹²³*Institute for High Energy Physics of the National Research Centre Kurchatov Institute, Protvino, Russia*
- ¹²⁴*Institute for Theoretical and Experimental Physics named by A.I. Alikhanov of National Research Centre "Kurchatov Institute", Moscow, Russia*
- ¹²⁵*Department of Physics, New York University, New York New York, USA*
- ¹²⁶*Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo, Japan*
- ¹²⁷*Ohio State University, Columbus Ohio, USA*
- ¹²⁸*Faculty of Science, Okayama University, Okayama, Japan*
- ¹²⁹*Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman Oklahoma, USA*

- ¹³⁰*Department of Physics, Oklahoma State University, Stillwater Oklahoma, USA*
- ¹³¹*Palacký University, RCPTM, Joint Laboratory of Optics, Olomouc, Czech Republic*
- ¹³²*Center for High Energy Physics, University of Oregon, Eugene Oregon, USA*
- ¹³³*Graduate School of Science, Osaka University, Osaka, Japan*
- ¹³⁴*Department of Physics, University of Oslo, Oslo, Norway*
- ¹³⁵*Department of Physics, Oxford University, Oxford, United Kingdom*
- ¹³⁶*LPNHE, Sorbonne Université, Université de Paris, CNRS/IN2P3, Paris, France*
- ¹³⁷*Department of Physics, University of Pennsylvania, Philadelphia Pennsylvania, USA*
- ¹³⁸*Konstantinov Nuclear Physics Institute of National Research Centre “Kurchatov Institute”,
PNPI, St. Petersburg, Russia*
- ¹³⁹*Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh Pennsylvania, USA*
- ^{140a}*Laboratório de Instrumentação e Física Experimental de Partículas—LIP, Lisboa, Portugal*
- ^{140b}*Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal*
- ^{140c}*Departamento de Física, Universidade de Coimbra, Coimbra, Portugal*
- ^{140d}*Centro de Física Nuclear da Universidade de Lisboa, Lisboa, Portugal*
- ^{140e}*Departamento de Física, Universidade do Minho, Braga, Portugal*
- ^{140f}*Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain), Spain*
- ^{140g}*Dep Física and CEFITEC of Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa,
Caparica, Portugal*
- ^{140h}*Instituto Superior Técnico, Universidade de Lisboa, Lisboa, Portugal*
- ¹⁴¹*Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic*
- ¹⁴²*Czech Technical University in Prague, Prague, Czech Republic*
- ¹⁴³*Charles University, Faculty of Mathematics and Physics, Prague, Czech Republic*
- ¹⁴⁴*Particle Physics Department, Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ¹⁴⁵*IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France*
- ¹⁴⁶*Santa Cruz Institute for Particle Physics, University of California Santa Cruz,
Santa Cruz California, USA*
- ^{147a}*Departamento de Física, Pontificia Universidad Católica de Chile, Santiago, Chile*
- ^{147b}*Universidad Andres Bello, Department of Physics, Santiago, Chile*
- ^{147c}*Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso, Chile*
- ¹⁴⁸*Department of Physics, University of Washington, Seattle Washington, USA*
- ¹⁴⁹*Department of Physics and Astronomy, University of Sheffield, Sheffield, United Kingdom*
- ¹⁵⁰*Department of Physics, Shinshu University, Nagano, Japan*
- ¹⁵¹*Department Physik, Universität Siegen, Siegen, Germany*
- ¹⁵²*Department of Physics, Simon Fraser University, Burnaby BC, Canada*
- ¹⁵³*SLAC National Accelerator Laboratory, Stanford California, USA*
- ¹⁵⁴*Physics Department, Royal Institute of Technology, Stockholm, Sweden*
- ¹⁵⁵*Departments of Physics and Astronomy, Stony Brook University, Stony Brook New York, USA*
- ¹⁵⁶*Department of Physics and Astronomy, University of Sussex, Brighton, United Kingdom*
- ¹⁵⁷*School of Physics, University of Sydney, Sydney, Australia*
- ¹⁵⁸*Institute of Physics, Academia Sinica, Taipei, Taiwan*
- ^{159a}*E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi, Georgia*
- ^{159b}*High Energy Physics Institute, Tbilisi State University, Tbilisi, Georgia*
- ¹⁶⁰*Department of Physics, Technion, Israel Institute of Technology, Haifa, Israel*
- ¹⁶¹*Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv, Israel*
- ¹⁶²*Department of Physics, Aristotle University of Thessaloniki, Thessaloniki, Greece*
- ¹⁶³*International Center for Elementary Particle Physics and Department of Physics, University of Tokyo,
Tokyo, Japan*
- ¹⁶⁴*Graduate School of Science and Technology, Tokyo Metropolitan University, Tokyo, Japan*
- ¹⁶⁵*Department of Physics, Tokyo Institute of Technology, Tokyo, Japan*
- ¹⁶⁶*Tomsk State University, Tomsk, Russia*
- ¹⁶⁷*Department of Physics, University of Toronto, Toronto ON, Canada*
- ^{168a}*TRIUMF, Vancouver BC, Canada*
- ^{168b}*Department of Physics and Astronomy, York University, Toronto ON, Canada*
- ¹⁶⁹*Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied
Sciences, University of Tsukuba, Tsukuba, Japan*
- ¹⁷⁰*Department of Physics and Astronomy, Tufts University, Medford Massachusetts, USA*
- ¹⁷¹*Department of Physics and Astronomy, University of California Irvine, Irvine California, USA*
- ¹⁷²*Department of Physics and Astronomy, University of Uppsala, Uppsala, Sweden*
- ¹⁷³*Department of Physics, University of Illinois, Urbana Illinois, USA*

¹⁷⁴*Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia—CSIC, Valencia, Spain*

¹⁷⁵*Department of Physics, University of British Columbia, Vancouver BC, Canada*

¹⁷⁶*Department of Physics and Astronomy, University of Victoria, Victoria BC, Canada*

¹⁷⁷*Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg, Germany*

¹⁷⁸*Department of Physics, University of Warwick, Coventry, United Kingdom*

¹⁷⁹*Waseda University, Tokyo, Japan*

¹⁸⁰*Department of Particle Physics, Weizmann Institute of Science, Rehovot, Israel*

¹⁸¹*Department of Physics, University of Wisconsin, Madison Wisconsin, USA*

¹⁸²*Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik,
Bergische Universität Wuppertal, Wuppertal, Germany*

¹⁸³*Department of Physics, Yale University, New Haven Connecticut, USA*

^aDeceased.

^bAlso at Department of Physics, King's College London, London, United Kingdom.

^cAlso at Instituto de Física Teórica, IFT-UAM/CSIC, Madrid, Spain.

^dAlso at TRIUMF, Vancouver BC, Canada.

^eAlso at Department of Physics and Astronomy, University of Louisville, Louisville, Kentucky, USA.

^fAlso at Physics Department, An-Najah National University, Nablus, Palestine.

^gAlso at Department of Physics, University of Fribourg, Fribourg, Switzerland.

^hAlso at Physics Dept, University of South Africa, Pretoria, South Africa.

ⁱAlso at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona, Spain.

^jAlso at Tomsk State University, Tomsk, and Moscow Institute of Physics and Technology State University, Dolgoprudny, Russia.

^kAlso at Department of Physics, Ben Gurion University of the Negev, Beer Sheva, Israel.

^lAlso at Università di Napoli Parthenope, Napoli, Italy.

^mAlso at Institute of Particle Physics (IPP), Vancouver, Canada.

ⁿAlso at Department of Physics, University of Adelaide, Adelaide, Australia.

^oAlso at Dipartimento di Matematica, Informatica e Fisica, Università di Udine, Udine, Italy.

^pAlso at Department of Physics, St. Petersburg State Polytechnical University, St. Petersburg, Russia.

^qAlso at Borough of Manhattan Community College, City University of New York, New York New York, USA.

^rAlso at Department of Physics, California State University, Fresno, USA.

^sAlso at Department of Financial and Management Engineering, University of the Aegean, Chios, Greece.

^tAlso at Department of Physics, California State University, East Bay, USA.

^uAlso at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona, Spain.

^vAlso at Department of Physics, University of Michigan, Ann Arbor Michigan, USA.

^wAlso at IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay, France.

^xAlso at Graduate School of Science, Osaka University, Osaka, Japan.

^yAlso at Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg, Germany.

^zAlso at Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan.

^{aa}Also at Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen, Netherlands.

^{bb}Also at CERN, Geneva, Switzerland.

^{cc}Also at Department of Physics, Stanford University, Stanford California, USA.

^{dd}Also at Manhattan College, New York New York, USA.

^{ee}Also at Joint Institute for Nuclear Research, Dubna, Russia.

^{ff}Also at Hellenic Open University, Patras, Greece.

^{gg}Also at The City College of New York, New York New York, USA.

^{hh}Also at Department of Physics, California State University, Sacramento, USA.

ⁱⁱAlso at Moscow Institute of Physics and Technology State University, Dolgoprudny, Russia.

^{jj}Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève, Switzerland.

^{kk}Also at Louisiana Tech University, Ruston Louisiana, USA.

^{ll}Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia, Bulgaria.

^{mmm}Also at Faculty of Physics, M.V. Lomonosov Moscow State University, Moscow, Russia.

ⁿⁿAlso at Department of Applied Physics and Astronomy, University of Sharjah, Sharjah, United Arab Emirates.

^{oo}Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg, Germany.

^{pp}Also at CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France.

^{qq}Also at National Research Nuclear University MEPhI, Moscow, Russia.

^{rr}Also at Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics, Budapest, Hungary.

^{ss}Also at Giresun University, Faculty of Engineering, Giresun, Turkey.

^{tt}Also at Department of Physics and Astronomy, Michigan State University, East Lansing Michigan, USA.