Extended lactation in dairy cows

effects of milking frequency, calving season and nutrition on lactation persistency and milk quality

Sorensen, Annette; Muir, D. Donald; Knight, Christopher Harold

Published in:
Journal of Dairy Research

DOI:
[10.1017/S0022029907002944](https://doi.org/10.1017/S0022029907002944)

Publication date:
2008

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Sorensen, A., Muir, D. D., & Knight, C. H. (2008). Extended lactation in dairy cows: effects of milking frequency, calving season and nutrition on lactation persistency and milk quality. *Journal of Dairy Research*, 75(1), 90-97. <https://doi.org/10.1017/S0022029907002944>

Extended lactation in dairy cows: effects of milking frequency, calving season and nutrition on lactation persistency and milk quality

Annette Sorensen¹, D Donald Muir² and Christopher H Knight^{3*†}

¹ Cancer Research UK Beatson Laboratories, University of Glasgow, Glasgow G61 1BD

² DD Muir Consultants, Irvine KA11 1QQ

³ Faculty of Veterinary Medicine Institute of Comparative Medicine, Division of Cell Sciences, University of Glasgow, Glasgow G61 1QH

Received 17 July 2007 and accepted for publication 23 October 2007

Twelve spring-calving and twelve winter-calving cows were managed for extended lactation cycles of 18-months duration, with the former group then completing a second extended lactation. Half of the cows were fed according to standard management practice for the herd; the other half received supplementary concentrate from week 9 of lactation onwards. Commencing at the same time, half of the udder of each cow was subjected to increased milking frequency (thrice daily rather than twice daily). Lactation persistency (and hence total milk yield) was significantly increased by frequent milking. Winter calving cows and supplemented cows also exhibited better persistency, but this was only evident up until the point of re-breeding, at around lactation week 33. Milk composition was measured in the spring-calving cows in both their first and second extended lactations. Composition altered during the course of the lactation, protein and fat percentages increasing and lactose percentage decreasing, irrespective of treatment. The quality of the milk for processing into cheese, fermented products, heat-treated products and cream liqueurs was assessed by calculation of casein number (casein protein as a proportion of total protein). Processing quality declined across the course of lactation in those groups that showed poor persistency but not in those that maintained a persistent lactation. Milk hygienic quality (somatic cell counts) showed parallel changes. Body condition score increased during the course of lactation but was not affected by supplementation; none of the cows became excessively fat. All cows remained healthy throughout the extended lactations and the majority (33/36) re-bred successfully. By demonstrating that lactation persistency is plastic and can be improved by simple management interventions, the results lend support to the economic arguments in favour of extended lactation cycles. The likely welfare benefits of extended lactation are also discussed.

Keywords: Dairy cattle, extended lactation, persistency, milk quality.

It has been recognized for many years that pregnancy and calving are pre-requisites to bovine lactation. The shape of the lactation curve (a rapid rise to a peak followed by a gradual decline) has hitherto meant that milk production is optimized by re-breeding to achieve a 12-month calving pattern (Holmann et al. 1984). Milk yields have increased

considerably over several decades through genetic selection and improved nutrition, to the point where the highest yielding Holstein herds in the USA have an average annual production of more than 14 000 kg/cow (Kellogg et al. 2001), several-fold greater than those cows for which the 12-month strategy was originally devised. Food intake has risen to a lesser extent (Veerkamp et al. 1995), so the need to mobilize body reserves has increased. Partly as a consequence, fertility has declined (Royal et al. 2000) and health has tended to deteriorate (Pryce et al. 1997). The notion that high-yielding dairy cows are metabolically stressed (Nielsen, 1999) has prompted a desire to explore

*Present address: University of Copenhagen Faculty of Life Sciences, Department of Basic Animal and Veterinary Sciences, Grønnegårdsvej 7 DK 1870 Frederiksberg C, Denmark

†For correspondence; e-mail: chkn@life.ku.dk

alternative strategies that could combine economic viability with improved animal welfare. One such strategy is extended lactation, whereby re-breeding is delayed and management emphasis is given to maximizing lactation persistency (inverse of the rate of decline in yield with time) rather than peak yield. The welfare rationale is evident when one considers lifetime health. Some 65% of health incidents occur in the first 45 d of lactation (Erb et al. 1984), hence, the fewer times that a cow calves the less she is exposed to risk. At its simplest, if the three 12-month lactation cycles typical of the lifetime of intensively managed dairy cattle could be replaced by two extended lactation cycles of 18-months duration, exposure to risk would be reduced by one-third. It seems inevitable that extended lactation would benefit welfare, although this has not yet been rigorously tested.

The economic argument for extended lactation is more debatable. Recent evidence suggests that high-yielding cows perform better in 14-month lactation cycles than in 12-month cycles, but exhibit abnormally long dry periods if managed for 18-month lactation cycles (Ratnayake et al. 1998). In neither of these studies was any attempt made to alter lactation persistency, so the reason for the long dry periods was that yield had declined to an economically non-viable level. Similarly, modelling analysis of 'extended lactations' done retrospectively from national databases (Vargas et al. 2000; Grossman & Koops, 2003) invariably uses data from cows that have failed to re-breed, not cows that have been deliberately managed for extended lactation. There can be little doubt that if lactation persistency could be improved, extended lactations would become profitable. Using the same comparison of three 12-month v. two 18-month lactations referred to previously, it is inevitable that, at some degree of improved persistency, the extended lactation scenario will perform better because it requires only one dry period rather than two.

Whilst it is evident from simple comparison of heifer and cow lactations that persistency is variable, very little systematic effort has been made to manipulate it (McFadden, 1997). We have begun to address this deficiency, by examining the effects of altered nutrition, calving season and milking frequency on lactation persistency in cows managed for 18-month lactation cycles. Since late lactation milk is sometimes associated with poor milk quality (Lucey, 1996) we also determined casein number as a measure of processing quality.

Materials and Methods

Cows

The experiment was conducted at the former Hannah Research Institute. A total of 25 Holstein:Friesian cows were used in a factorial design to study the effects of milking frequency, calving season and nutrition on lactation characteristics during a total of 36, 18-month

extended lactation cycles. Experimental cows were selected from cows calving between 24 March and 23 May 1996 (for the Spring group) or between 14 October and 10 December 1996 (for the Winter group). Parity ranged from 1 to 5 at the outset. All cows were fed a grass silage-based total mixed ration containing 15% crude protein (CP) during winter months and grazed pasture during summer supplemented with sugar-beet pulp during times of poor grass quality. In addition, in-parlour concentrate containing 18% CP was fed according to milk yield or experimental design. Re-breeding was allowed after week 32 post partum. Cows were inseminated when seen in oestrus. Cows were dried-off 8 weeks before expected parturition or when milk yield dropped below 3.5 kg/d for the lowest-yielding udder half, whichever was the earlier.

Experimental design

Initially there were two calving groups, Spring ($n=12$, mean parity 2.4), and Winter ($n=12$, mean parity 2.4). Eleven of the Spring group were then studied during their consecutive extended lactation (Second), calving between September 1997 and January 1998 (i.e. they were now winter-calving), a twelfth previously unstudied cow being added to maintain the size of this group. Commencing in lactation week 9 (9.5 ± 0.64), half of each calving group were fed an additional 3 kg/d of 18% CP in-parlour concentrate (High) whilst the remainder continued to be fed according to milk yield following standard practice for the herd (Low). Cows were blocked onto treatment as they reached lactation week 9. Allocation to High or Low took into account parity, body weight, body condition score and milk yield. Starting at the same time, half of the udder (diagonally opposed quarters) was milked thrice daily (3X) at intervals of 8 h whilst the other half continued on twice-daily milking (2X: 8:16-h intervals). This milking treatment was applied to every cow and continued for the remainder of the lactation. The experimental design thus comprised three calving groups (Spring, Winter, Second $n=12$), two levels of nutrition (High, Low, $n=18$) and two milking frequencies (3X, 2X, $n=36$).

Measurements

Milking was performed in a parlour equipped to collect milk separately from half-udders (diagonally opposed quarters) and to weigh the milk electronically at every milking to a precision of 100 g. Half-udder milk samples were collected at monthly intervals from the Spring cows during both of their extended lactations, i.e. Spring and Second groups, each sample being a proportionally bulked composite sample from all milkings in one day. Milk samples from all cows were analysed for fat and lactose using infra-red technology (MilkoScan, Foss, DK 3400 Hillerød, Denmark) and for somatic cell count (SCC) using flow cytometry (FossoMatic) by Livestock Services (UK) Ltd, Paisley PA3 1TJ. In the first lactation additional

analysis of total protein and detailed analyses of casein protein were performed on samples bulked according to treatment according to methods described in Brown et al. (1995). In the second lactation all protein analyses were done on samples from individual cows. In the earlier parts of the experiment cows were weighed weekly; in the latter parts weighing was performed through an automated weigh-bridge after each milking. Body condition score was measured weekly using a five-point scale where 1 is extremely thin and 5 extremely fat.

Statistical analysis

Statistical analysis was performed using analysis of variance in Genstat 5 Release 4.1, Lawes Agricultural Trust, Rothamsted, UK or Minitab Release11, Minitab Inc, State College, PA16801 USA. Factors included in the model were cow, calving group, nutritional treatment and, for milk yield and composition data only, milking frequency. Lactation persistency was defined as the slope of the decline in milk yield from week 9 of lactation onwards, and calculated using best-fit linear regression of weekly averaged milk yields. Other curve-fits were examined but, with the possible exception of a ninth-grade polynomial, did not produce better fits to the data. Slopes obtained were analysed with peak milk yield included as a co-variate. Effects of milking frequency were analysed within-cow, other factors were analysed between-cow. Effects of pregnancy on lactation persistency were analysed by residual maximum likelihood (REML) owing to an unbalanced design.

Results

Lactation characteristics and milk yield

Half-udder whole-lactation milk yield was significantly increased by milking more frequently (by 33% overall, Table 1) and by nutritional supplementation (by 11%) and differed between calving groups. Yield was greater in winter calving cows than in spring calving (by 15%) and in the second extended lactation of cows that first calved in the spring and then in the winter (by 17%). Cows were dried off in both udder-halves at the same time, hence milking frequency is not a factor in analysis of lactation length or dry period length. Lactation length was not affected by nutrition, but differed between calving groups (Table 1). Second extended lactations were shorter than others, not because of lower yield but because of a shorter calving interval as a result of an improved first-service to conception interval. Dry periods were shorter in Second and were also shorter in High compared with Low, because fewer cows were dried off early as a result of low yield. Body weight and body condition score are cow characteristics and thus independent of the milking frequency treatment, and cows were allocated to nutritional treatments taking into account pretreatment body weight

and condition score so there were no differences at the start of treatment. Body weight and condition score both increased during the course of lactation. Final body weight (i.e. at the end of lactation) was higher in Second than in Spring or Winter, but nutrition had no effect on either final body weight or final body condition score (Table 1).

Annualized milk production was calculated for the eleven cows that completed two consecutive extended lactations (Spring then Second). The time taken to produce the milk was taken to be the length of the two lactations plus one dry period (data in Table 1). On a whole-cow basis (ie 2X+3X udder halves) annual milk production for the Low and High input groups was 6531 ± 243 and 7501 ± 216 kg, respectively. On the basis of production data for the 3X udder half, the most that might have been produced (had both halves of the udder been milked thrice daily and achieved the same level of output) was 7453 ± 327 and 8463 ± 227 kg/year for Low and High, respectively.

Lactation persistency

Persistency was significantly and negatively correlated with peak milk yield ($r^2=0.49$, $P<0.001$) but was not significantly correlated with any measure of body weight or body composition score. When persistency slopes were recalculated by week of recurring pregnancy rather than by week of lactation, it was evident that persistency was reduced significantly in the last third of pregnancy (slopes of -0.1649 , -0.1408 , -0.1038 and -0.3215 for the 10 weeks before conception and the first, second and third 10-week blocks of pregnancy, respectively, SED 0.043 , $P<0.001$, analysis of variance). Accordingly, subsequent analyses included peak yield as a covariate and were run separately for the periods up to week 33 of lactation (when re-breeding commenced) and up to week 20 of recurring pregnancy (i.e. excluding the last 10 weeks) as well as for the whole of the lactation (Table 2). Within lactation groups there were highly significant correlations between the various measures of persistency, and between 2X and 3X udder-halves. Persistency in the first extended lactation of Spring cows was only a modest predictor of persistency in their subsequent lactation (Second). Correlation coefficients of -0.11 (NS), 0.42 ($P<0.05$) and 0.54 ($P=0.01$) were obtained for persistency slopes running from weeks 9 to lactation 33, pregnancy 20 and end of lactation, respectively.

Up until week 33 of lactation, persistency was significantly improved by milking more frequently and Winter cows were more persistent than either Second or Spring (Table 2). Supplementary feeding also tended to improve persistency, although this effect was not significant (Table 2). When analysis was extended to week 20 of pregnancy the only significant effect was a positive one of frequent milking (Table 2) and analysis through to the end of lactation failed to detect any significant differences due to treatments. Slopes for weeks 9–20 were extrapolated to

Table 1. Lactation length, days dry, final body weight and condition score and total lactation milk yield in cows calving in different seasons, fed conventionally (Low) or supplemented (High) and milked by half-udder either twice (2X) or thrice (3X) daily. Milk yields are half-udder. Values are means $\pm$ SE., $n=12$ per group

	Group						P value	
	Nutrition	Spring		Winter		Second		
		2X	3X	2X	3X	2X		3X
Lactation length, d	Low	497±19		556±31		474±18		NS
	High	515±15		575±17		481±18		
Dry period length, d	Low	124±25		84±16		65±6		NS
	High	72±7		56±4		56±2		
Total yield, kg	Low	3563±226	4861±279	4473±346	5960±479	4532±224	5778±292	NS
	High	4259±345	5969±227	4626±531	6433±553	5322±335	6218±208	
Final body weight, kg	Low	672±26		652±16		733±54		NS
	High	647±23		713±36		754±14		
Final BCS, units	Low	4.3±0.4		3.0±0.4		2.9±0.4		NS
	High	3.3±0.28		3.8±0.49		2.8±0.4		

a half-udder yield of 5 kg/d to calculate the theoretical effect of improved persistency on lactation length in the absence of recurring pregnancy. Frequent milking increased theoretical lactation length from 68 ± 3 to 102 ± 8 weeks ($P < 0.001$), whilst other treatments had no significant effect.

Milk composition

Milk compositional analysis was restricted to the Spring and Second groups, i.e. the first and second consecutive extended lactations of the same cows. Results are shown in Fig. 1. For all major components, changes occurring across the course of lactation were of greater magnitude than differences due to treatment effects. Protein (crude protein and casein protein) and fat contents increased as lactation advanced whilst lactose decreased (Fig. 1). Nutritional supplementation slightly but significantly increased the contents of casein protein, fat and lactose and frequent milking increased fat and lactose but not protein. There were few differences between the first and second lactations apart from increased lactose content in the latter. The 2X udder halves of the Low nutrition group consistently showed the greatest decrease in lactose content in later lactation (Fig. 1) and the smallest increase in casein protein content (results not shown).

Milk quality

The processing quality of the milk was assessed by calculating casein number, which is casein protein as a percentage of total protein. The hygienic quality of the milk was assessed by measuring SCC, which is commonly used as an indicator of the health status of the mammary gland. Results are in Fig. 2. Both quality indicators tended to worsen across the course of lactation: casein number fell and SCC increased. These changes were most evident in the unsupplemented cows and 2X udder halves, so were especially evident in Low 2X (Fig. 2). The beneficial effects of frequent milking and nutritional supplementation were both highly significant ($P < 0.001$). There were also significant differences between the two lactations, with better milk quality being recorded in the first lactation. Used as a measure of mastitis, SCC in excess of 200 000 is regarded as being indicative of a disease condition. At no time during either lactation did SCC exceed 200 000 in the 3X udder-halves of supplemented cows (Fig. 2).

Discussion

Extended lactation has the potential to become a welfare-friendly alternative to intensive lactation cycles (Knight, 1984), but for it to achieve economic viability lactation persistency (the slope of decline in milk yield after peak lactation) must be amenable to improvement (Knight & Mainland, 1995). It is evident from the results presented

Table 2. Lactation persistency slopes in cows calving in different seasons, fed conventionally (Low) or supplemented (High) and milked by half-udder either twice (2X) or thrice (3X) daily. Slopes are calculated by best-fit linear regression analysis of weekly yields between week 9 of lactation (L9) and either week 33 of lactation (L33), week 20 of concurrent pregnancy (P20) or the end of lactation. A less negative value indicates greater persistency. All values are adjusted for peak yield as covariate. Values are means, $n=12$ per group. All individual $se \leq 0.01$ or less

		Group						P value			
		Spring		Winter		Second		Frequency	Group	Nutrition	Interaction terms
Nutrition		2X	3X	2X	3X	2X	3X				
L 9 to L33	Low	-0.2942	-0.2977	-0.1958	-0.0790	-0.1633	-0.0650	=0.001	<0.001	NS	NS
	High	-0.2440	-0.2011	-0.1407	-0.0388	-0.1667	-0.1132				
L9 to P20	Low	-0.1754	-0.1757	-0.1621	-0.1514	-0.1932	-0.1271	<0.05	NS	NS	NS
	High	-0.1654	-0.1350	-0.1717	-0.1543	-0.1985	-0.1671				
L9 to end	Low	-0.1728	-0.1783	-0.1564	-0.1521	-0.2068	-0.1414	NS	NS	NS	NS
	High	-0.1567	-0.1311	-0.1675	-0.1527	-0.1992	-0.1682				

here that bovine lactation persistency is indeed plastic and can be improved by simple management procedures. In particular, persistency was increased by milking thrice daily rather than twice daily. Persistency varied between cows in a way that we could not predict from weight or body condition score measured in early lactation, but the effect of milking frequency was seen in all cows. The overall effect would be to increase calculated annual yield from a herd average of around 6000 kg to almost 8500 kg per cow, a considerably greater increase than the 10–15% usually associated with thrice-daily milking. Whilst there have been numerous studies of lactation persistency modelled retrospectively from commercial yield records where the reason for the lactation being long is unknown, to our knowledge this is the first report of a prospective study specifically designed to manipulate persistency under UK conditions. Van Amburgh et al. (1997) have reported on extended lactations using rbST in commercial dairy herds in the USA, and showed that persistency was markedly increased in heifers that were not re-bred. Milking frequency was not reported. Planned calving intervals of 12, 15 and 18 months have been compared experimentally in Swedish cattle, some of which were milked three times daily. Only a preliminary and incomplete report of milk yield responses has been published (Bertilsson et al. 1997), but there was no evidence that persistency was altered by milking frequency. Analysis was apparently done on the whole of the lactation, i.e. including all of the concurrent pregnancy. Comparison of Moshave (family) herds and Kibbutz herds using data from the Israeli national database has also failed to identify differences in persistency, even though at the time of analysis Kibbutz usually milked thrice daily whilst Moshave milked twice daily (CH Knight and E Maltz, unpublished observations). However, the standard management practice in both categories of herd would have included 12-month calving intervals. In the present study persistency was greater in 3X until the last 10 weeks of the recurring

pregnancy, at which time the negative effect of the pregnancy overcame the positive effect of frequent milking. Hence these data are not at variance with our own observations and the overall conclusion appears clear: persistency is responsive to milking frequency, but the effects are negated by recurring pregnancy. The importance of milking frequency is reinforced by the observation that milking infrequently (once daily) decreases lactation persistency (Hickson et al. 2006). The proposed mechanism underlying the effect of increased milking frequency involves the combined action of growth hormone (GH) and prolactin in stimulating the local secretion of IGF1, a mammary secretory cell-survival factor, whilst inhibiting local production of IGFBP5, which inhibits this activity (Flint & Knight, 1997). The net effect is decreased apoptosis, thus increased persistency. The control of lactation persistency has been considered in more detail elsewhere (McFadden, 1997; Knight, 2000; Capuco et al. 2003).

Increased interest in persistency (15 papers listed by PubMed in 2006 compared with an average of 3 a year during the 1990s) reflects a realization that re-breeding at around 60 d post partum may no longer represent optimal management, on either welfare or economic grounds. In a recent theoretical modelling analysis, a 30% improvement in persistency resulted in economically optimal re-breeding times of greater than 450 d for heifers and around 200 d for second lactation cows (De Vries, 2006). For cows on standard nutrition in their second extended lactation we observed a 34% difference in persistency between 2X and 3X udder halves. The overall improvement attributable to 3X milking was around half this figure and we cannot exclude that 3X milking of one udder half caused a decrease in the other half rather than an increase in itself, nevertheless the basic observation of a significant difference in persistency as a result of a management intervention remains valid and our results provide quantitative support for the assumptions made in the model.

Fig. 1. Milk composition of Spring cows managed for extended lactation cycles of 18-months duration, fed conventionally (open symbols) or supplemented with additional concentrate (closed symbols) and milked (on a half-udder basis) twice daily (dashed lines) or thrice daily (continuous lines). For clarity, values are presented as smoothed means (smooth function in Minitab Release 11) with representative SE shown for fat and lactose. Protein was measured in pooled samples. Analysis was performed on unsmoothed means.

The proposed benefits of extended lactation cycles include reduced disease incidence and fewer reproductive problems. Allore & Erb (2000) modelled the effect of extending the voluntary wait period (VWP) by 100 d,

Fig. 2. Milk quality attributes of Spring cows managed for extended lactation cycles of 18-months duration, fed conventionally (open symbols) or supplemented with additional concentrate (closed symbols) and milked (on a half-udder basis) twice daily (dashed lines) or thrice daily (continuous lines). Casein number (top panel) is casein expressed as a percentage of total protein and is a measure of processing quality. SCC is somatic cell count and is conventionally used as a measure of health status (but see text). For clarity, values are presented as smoothed means (smooth function in Minitab Release 11) with representative SE for SCC. Analysis was performed on unsmoothed means.

equivalent to 15-month extended lactation cycles, on the incidence of 10 diseases including mastitis, metabolic and reproductive diseases. All were reduced. The modelled effect of increasing the VWP by 100 d was to increase lactation length by 89 d, as a result of improved fertility later in lactation (Allore & Erb, 2000). Whilst the negative effect of peak yield on fertility is well established, there is little documented evidence to support the expected improvement later in lactation. An improvement in conception to first insemination in cows inseminated at 120–150 d compared with 35–59 d was noted by Schindler et al. (1991) but the same study showed no difference in heifers

and in the Swedish comparison of cows re-bred for 12, 15 or 18-month calving intervals reproductive success did not differ between groups (Ratnayake et al. 1998). The present study was not designed to enable statistical analysis of health or reproductive parameters; the number of cows used was too small. We encountered very few health problems and re-breeding success was similar to the herd average, but more work is needed to properly quantify these variables.

For extended lactation to be profitable it is important that milk quality is not compromised. The increases in total protein and fat content that we observed across the course of the lactation are not unexpected (see Knight et al. 2000 for review) and are likely to represent increased rather than decreased income for the farmer. However, the quality of the milk to the processor is related more to casein content than to protein content, and can be measured as casein number. Casein number typically declines during the course of normal lactations, especially in low-input systems where it can compromise the processing qualities of late-lactation milk (Lucey, 1996). We observed a significant decline that was ameliorated by increased nutrition and by increased frequency, such that 3X milked glands of high input cows maintained a constant or nearly constant casein number throughout. The decline is a consequence of influx of serum proteins and consequential proteolysis of secreted casein; the major proteolytic enzymes are serum-derived and arise in stored milk as a result of paracellular flux through 'leaky' tight junctions (Politis et al. 1988), hence casein number is directly related to epithelial integrity. Integrity was measured in the Spring cows using the ratio of sodium to potassium (Sorensen et al. 2001) and was shown to deteriorate significantly more in 2X than in 3X, hence the observed differences in casein number. From a commercial point of view this is good news; the same treatments that increased persistency also maintained milk processing quality. Interestingly, a 'salty' taste was observed in late lactation milk from extended-lactation cows in the Swedish study (Bertilsson et al. 1997), suggesting that the treatments used had failed to maintain junctional integrity such that the ionic composition of the milk had moved towards the higher sodium, lower potassium content of plasma. The high casein number that we observed is extremely important, associated as it is with higher cheese yield, improved structure in fermented milks, improved stability of evaporated milk and, as a consequence of decreased plasmin activity, improved stability of UHT milk, cream and cream liqueurs.

SCC is often used as an index of udder health, and in the EU a maximum limit for saleable milk is set at 400 000 cells/ml. On this basis, milk from low input, 2X udder-halves would have failed from around week 50 of lactation onwards. It is important to recognize that mastitis only accounts for a quarter to a third of observed variation in SCC (Morris, 1975) and, like proteolytic enzymes, the majority of somatic cells get into milk through tight

junctions that have lost patency (Lin et al. 1995). The gradual increase that we observed in SCC across the course of lactation is related to declining epithelial integrity, not to disease. There were few clinical cases of mastitis and incidence did not increase in late lactation. Importantly, in the same way that frequent milking and increased nutrition prevented the deterioration in processing quality, they also prevented the rise in SCC.

In conclusion, the results provide support for the use of extended 18-month lactation cycles as an alternative to 12-month intensive lactation cycles, provided frequent milking can be practised. Economic and welfare benefits would be expected but have not yet been demonstrated in practice. More research is needed to quantify both aspects.

This research was funded by Department for Environment, Food and Rural Affairs. The Hannah Institute was in receipt of grant-in-aid from Scottish Executive Environment and Rural Affairs Department. We are grateful to John Munro and his staff for care of the animals. The skilled technical support of Stuart Robertson, Jean Thomson, Andrew Law, Eleanor Noble, Sandra Hodge and Julie Currie is gratefully acknowledged. Statistical advice in experimental design and analysis was provided by Tony Hunter and Ian Nevison of BIOSS, for which we are indebted. CHK is grateful to his late father, Brian Knight, for providing a number of the cows used in the work.

References

- Allore HG & Erb HN 2000 Simulated effects on dairy cattle health of extending the voluntary waiting period with recombinant growth hormone. *Preventive Veterinary Medicine* **46** 29–50
- Arbel R, Bigun Y, Ezra E, Sturman H & Hojman D 2001 The effect of extended calving intervals in high lactating cows on milk production and profitability. *Journal of Dairy Science* **84** 600–608
- Bertilsson J, Berglund B, Ratnayake G, Svennersten-Sjaunja K & Wiktorsson H 1997 Optimising lactation cycles for the high-yielding dairy cow. A European perspective. *Livestock Production Science* **50** 5–13
- Brown JR, Law AJR & Knight CH 1995 Changes in casein composition of goats' milk during the course of lactation: physiological inferences and technological implications. *Journal of Dairy Research* **62** 431–439
- Capuco AV, Ellis SE, Hale SA, Long E, Erdman RA, Zhao X & Paape MJ 2003 Lactation persistency: insights from mammary cell proliferation studies. *Journal of Dairy Science* **81** Supplement 3 18–31
- De Vries A 2006 Economic value of pregnancy in dairy cattle. *Journal of Dairy Science* **89** 3876–3885
- Erb HN, Smith RD, Hillman RB, Powers PA, Smith MC, White ME & Pearson EG 1984 Rates of diagnosis of six diseases of Holstein cows during 15-day and 21-day intervals. *American Journal of Veterinary Research* **45** 333–335
- Flint DJ & Knight CH 1997 Interactions of prolactin and growth hormone (GH) in the regulation of mammary gland function and epithelial cell survival. *Journal of Mammary Gland Biology and Neoplasia* **2** 41–48
- Grossman M & Koops WJ 2003 Modeling extended lactation curves of dairy cattle: a biological basis for the multiphasic approach. *Journal of Dairy Science* **86** 988–998
- Hickson RE, Lopez-Villalobos N, Dalley DE, Clark DA & Holmes CW 2006 Yields and persistency of lactation in Friesian and Jersey cows milked once daily. *Journal of Dairy Science* **89** 2017–2024
- Holmann FJ, Shumway CR, Blake RW, Schwart RB & Sudweeks EM 1984 Economic value of days open for holstein cows of alternative milk yields with varying calving intervals. *Journal of Dairy Science* **67** 636–643

- Kellogg DW, Johnson ZB & Pennington JA** 2001 Survey of management practices used for the highest-producing DHI herds in the United States. *Journal of Dairy Science* **84** (E suppl) E120–E127
- Knight CH** 1984 Mammary growth and development: strategies of animals and investigators. *Symposia of the Zoological Society of London* **51** 147–170
- Knight CH** 2000 The importance of cell division in udder development and lactation. *Livestock Production Science* **66** 169–176
- Knight CH & Mainland D** 1995 Physiology of milk production; how can it be manipulated. *Cattle Practice* **3** 169–173
- Knight CH, Sorensen A & Muir DD** 2000 Non-nutritional (novel) techniques for manipulation of milk composition. *British Society of Animal Science Occasional Publication* **25** 223–239
- Lin Y, Xia L, Turner JD & Zhao X** 1995 Morphologic observation of neutrophil diapadesis across bovine mammary gland epithelium *in vitro*. *American Journal of Veterinary Research* **56** 203–207
- Lucey J** 1996 Cheesemaking from grass based seasonal milk and problems associated with late-lactation milk. *Journal of the Society of Dairy Technology* **49** 59–64
- McFadden TB** 1997 Prospects for improving lactation persistency. In *Milk Composition, Production and Technology*, pp. 319–339 (Eds RAS Welch, DJW Burns, SR Davis, AI Popay & CG Prosser). Wallingford, UK: CAB International
- Morris RS** 1975 Criteria for the design and evaluation of bovine mastitis control systems. *International Dairy Federation Bulletin* **85** 356–409
- Nielsen BL** 1999 Perceived welfare issues in dairy cattle, with special emphasis on metabolic stress. *British Society of Animal Science Occasional Publication* **24** 1–7
- Politis I, Lachance E, Block E & Turner JD** 1988 Plasmin/plasminogen in bovine milk: a relationship with involution. *Journal of Dairy Science* **72** 900–906
- Pryce JE, Veerkamp RF, Thompson R, Hill WG & Simm G** 1997 Genetic aspects of common health disorders and measures of fertility in Holstein Friesian dairy cattle. *Animal Science* **65** 353–360
- Ratnayake DRTG, Berglund B, Bertilsson J, Forsberg M & Gustafsson H** 1998 Fertility in cows managed for calving intervals of 12, 15 or 18 months. *Acta Veterinaria Scandinavica* **39** 215–228
- Royal M, Mann GE & Flint AP** 2000 Strategies for reversing the trend towards subfertility in dairy cattle. *Veterinary Journal* **160** 53–60
- Schindler H, Eger S, Davidson M, Ochowski D, Schermerhorn EC & Foote RH** 1991 Factors affecting response of groups of dairy cows managed for different calving-conception intervals. *Theriogenology* **36** 495–503
- Sorensen A, Muir DD & Knight CH** 2001 Thrice-daily milking throughout lactation maintains epithelial integrity and thereby improves milk protein quality. *Journal of Dairy Research* **68** 15–25
- Van Amburgh M, Galton D, Bauman D & Everett RW** 1997 Management and economics of extended calving intervals with use of BST. *Livestock Production Science* **50** 15–28
- Vargas B, Koops WJ, Herrero S & Van Arendonk JAM** 2000 Modeling extended lactations of dairy cows. *Journal of Dairy Science* **83** 1371–1380
- Veerkamp RF, Simm G & Oldham JD** 1995 Genotype by environment interactions: experience from Langhill. *Occasional Publications of the British Society of Animal Science* **19** 59–66