**Three-axis anisotropic exchange coupling in the single-molecule magnets
 $\text{NEt}_4[\text{Mn}^{\text{III}}_2(5\text{-Brsalen})_2(\text{MeOH})_2\text{M}^{\text{III}}(\text{CN})_6]$ ($\text{M}=\text{Ru}$, Os)**

Dreiser, Jan; Pedersen, Kasper Steen; Schnegg, Alexander; Holldack, Karsten; Nehr Korn, Joscha; Waldmann, Oliver; Mironov, Vladimir S.; Tregenna-Piggott, Philip; Sigrist, Marc; Mutka, Hannu; Weihe, Høgni; Bendix, Jesper

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Dreiser, J., Pedersen, K. S., Schnegg, A., Holldack, K., Nehr Korn, J., Waldmann, O., Mironov, V. S., Tregenna-Piggott, P., Sigrist, M., Mutka, H., Weihe, H., & Bendix, J. (2013). *Three-axis anisotropic exchange coupling in the single-molecule magnets $\text{NEt}_4[\text{Mn}^{\text{III}}_2(5\text{-Brsalen})_2(\text{MeOH})_2\text{M}^{\text{III}}(\text{CN})_6]$ ($\text{M}=\text{Ru}$, Os)*. Poster session presented at 4th European Conference on Molecular Magnetism, Karlsruhe, Germany.

Jan Dreiser,^a Kasper S. Pedersen,^b Alexander Schnegg,^c Karsten Holldack,^d Joscha Nehrkorn,^c Oliver Waldmann,^e Vladimir S. Mironov,^f Philip Tregenna-Piggott,^g Marc Sigrist,^h Hannu Mutka,^h Høgni Weihe,^b Jesper Bendix,^b

^a Swiss Light Source, Paul Scherrer Institut, 5232 Villigen, Switzerland and Ecole Polytechnique Fédérale de Lausanne, Inst. de Physique de la Matière Condensée, Station 3, CH-1015 Lausanne, Switzerland; ^b Department of Chemistry, University of Copenhagen, DK-2100 Copenhagen, Denmark; ^c Helmholtz-Zentrum für Materialien und Energie, Institut für Silizium-Photovoltaik, 12489 Berlin, Germany; ^d Helmholtz-Zentrum für Materialien und Energie, Institut für Methoden und Instrumente der Forschung mit Synchrotronstrahlung, 12489 Berlin, Germany;

^e Physikalisches Institut, Universität Freiburg, 79104 Freiburg, Germany; ^f A.V. Schubnikov Institute of Crystallography, 119333 Moscow, Russia; ^g Laboratory for Neutron Scattering, Paul Scherrer Institut, 5232 Villigen PSI, Switzerland; ^h Institut Laue-Langevin, 38042 Grenoble Cedex 9, France.

The introduction of 4d and 5d ions into single-molecule magnets (SMMs) represents an intriguing path to enhance their properties [1]. Strong spin-orbit coupling and the diffuse 4d and 5d orbitals as compared to the 3d case enable large magnetic anisotropies and strong exchange coupling. Here we study isostructural Mn-M-Mn trinuclear SMMs, with M=Ru and Os, by frequency-domain Fourier-transform THz electron paramagnetic resonance (FDFT-EPR), inelastic neutron scattering (INS) and SQUID magnetometry [2,3]. We find that the exchange coupling between Mn^{III} and Os^{III} ions is anisotropic and of three-axis type, i.e., it is antiferromagnetic along x and z axes while it is ferromagnetic along y. A further analysis yields that this is a consequence of orbitally dependent exchange [4,5] providing a profound understanding of the magnetic behavior of the orbitally degenerate Ru^{III} and Os^{III} ions.

(a)

(b)

Figure: (a) Structure of the Mn-Os-Mn SMM from X-ray diffraction. The NEt_4^+ counterion and H atoms have been omitted. Color code: Mn, purple; Os, dark blue; O, red; N, blue; C, grey; Br, brown. (b) FDFT THz-EPR spectra obtained on Mn-Os-Mn.

Literature: [1] X.-Y. Wang, C. Avendano, K. R. Dunbar, *Chem. Soc. Rev.* 2011, **40**, 3213; [2] K. S. Pedersen, J. Dreiser, J. Nehrkorn, M. Gysler, M. Schau-Magnussen, A. Schnegg, K. Holldack, R. Bittl, S. Piligkos, H. Weihe, P. Tregenna-Piggott, O. Waldmann, J. Bendix, *Chem. Commun.* 2011, **47**, 6918; [3] J. Dreiser, K. S. Pedersen, A. Schnegg, K. Holldack, J. Nehrkorn, M. Sigrist, P. Tregenna-Piggott, H. Mutka, H. Weihe, V. S. Mironov, J. Bendix, O. Waldmann, *Chem. Eur. J.* 2013, **19**, 3693; [4] J. J. Borrás-Almenar, J. M. Clemente-Juan, E. Coronado, A. V. Pali, B. S. Tsukerblat, *J. Phys. Chem. A* 1998, **102**, 200; [5] V. S. Mironov, L. F. Chibotaru, A. Ceulemans, *J. Am. Chem. Soc.* 2003, **125**, 9750.