Carbohydrate and amino acid metabolism in rat cerebral cortex in moderate and extreme hypercapnia.

Quistorff, Bjørn; Folbergrová, J.; Norberg, Karin; K. Siesjö, Bo

Published in:
Journal of Neurochemistry

Publication date:
1975

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Quistorff, B., Folbergrová, J., Norberg, K., & K. Siesjö, B. (1975). Carbohydrate and amino acid metabolism in rat cerebral cortex in moderate and extreme hypercapnia. *Journal of Neurochemistry*, 25(4), 457-462.

CARBOHYDRATE AND AMINO ACID METABOLISM IN RAT CEREBRAL CORTEX IN MODERATE AND EXTREME HYPERCAPNIA

J. FOLBERGROVÁ,¹ K. NORBERG, B. QUISTORFF and B. K. SIESJÖ

Brain Research Laboratory, E-Blocket, University Hospital, Lund, Sweden,
and

Department of Biochemistry A, University of Copenhagen, Denmark

(Received 28 February 1975. Accepted 13 March 1975)

Abstract—The time course of changes in glycolytic and citric acid cycle intermediates and in amino acids was studied in acute and steady state hypercapnia. Experiments on unanaesthetized animals exposed to 10% CO₂ for 10, 20 and 60 s showed that there was a transient decrease in glycogen concentration, progressive increases in glucose-6-phosphate and fructose-6-phosphate and decreases in pyruvate and lactate. During this time the levels of amino acids and Krebs cycle intermediates did not change, except for a small fall in malate at 60 s. The results indicate that there was a decrease in glycolytic flux due to an inhibition of the phosphofructokinase reaction. Since the tissue levels of phosphocreatine, ATP, ADP and AMP were unchanged inhibition of phosphofructokinase was probably due to the fall in pH.

Anaesthetized animals were exposed to about 5% CO₂ (for 2, 5, 15, 30 and 60 min) or to about 45% CO₂ (for 5 and 15 min). Except for succinate, which increased, all citric acid cycle metabolites analysed (citrate, α -ketoglutarate, fumarate and malate) decreased with the rise in CO₂-tension. The sum of the amino acids analysed (glutamate, glutamine, aspartate, asparagine, alanine and GABA) decreased at extreme hypercapnia. The results suggest that Krebs cycle intermediates and amino acids are partly used as substrates for energy production when there is reduced pyruvate availability due to hypercapnia.

It is proposed that amino acid carbon is made available for oxidation via transamination (aspartate aminotransferase reaction) and deamination (glutamate dehydrogenase reaction) and that citric acid cycle intermediates are metabolized following a reversal of reactions usually leading to CO₂ fixation.

PREVIOUS results from this laboratory have shown that hypercapnia reduces the tissue concentrations of pyruvate and lactate as well as of the Krebs cycle intermediates citrate, α -ketoglutarate (α -KG) and malate (FOLBERGROVÁ *et al.*, 1972b, 1974a). Although information on changes in succinate and fumarate was not at hand, it was speculated that the increased CO₂ tensions led to a loss of carbon skeletons from the Krebs cycle. The cause of this hypothetical loss was not apparent in view of the fact that hypercapnic conditions may favour anaplerotic reactions due to CO₂ fixation. Since there was a decrease in the sum of the concentrations of glutamate, glutamine and aspartate (FOLBERGROVÁ *et al.*, 1974a), a shift of carbon atoms from the Krebs cycle to the amino acid pool did not seem likely. However, firm conclusions could not be drawn since the tissue concentrations of alanine and GABA were not measured.

The present experiments were undertaken to obtain additional information on changes affecting carbo-

hydrate and amino acid metabolism in the cerebral cortex in hypercapnia, and to allow defining the mechanisms involved. The following three aspects were studied. Firstly, in order to characterize the initial events, unanaesthetized animals were exposed to about 10% CO₂ and cerebral cortex was obtained for analyses after 10, 20 or 60 s of hypercapnia. Secondly, a more complete picture of citric acid cycle changes was obtained by measuring, apart from citrate, α -ketoglutarate and malate, also isocitrate, succinate and fumarate. Since the major part of the citric acid cycle intermediates was measured changes in the size of this pool could be estimated. Thirdly, the previous estimation of amino acid changes, as based on measurements of glutamate, glutamine and aspartate, was extended to include measurements of GABA, asparagine and alanine. Therefore, citric acid cycle intermediates and amino acids were measured in animals exposed to about 5% CO₂ (for 2, 5, 15, 30 and 60 min) or to about 45% CO₂ (for 5 and 15 min). The results obtained allow a tentative interpretation of the mechanisms whereby hypercapnia influences substrate utilization in the brain.

METHODS

The experiments were performed on male Wistar rats that were allowed free access to pellet food and water until

¹ On leave of absence from Institute of Physiology, Czechoslovak Academy of Sciences, Prague.

Abbreviations used: α -KG, α -ketoglutarate; FDH, fructose-1,6-diphosphate; DHAP, dihydroxyacetone phosphate; 3PG, 3-phosphoglycerate; PCr, phosphocreatine; PFK, phosphofructokinase; OAA, oxaloacetate.

operation. There were two main groups: unanaesthetized and anaesthetized. The unanaesthetized animals (205–225 g) were provided with indwelling tail artery catheters in a previous operation performed under halothane anaesthesia. When they had recovered completely from this operation they were introduced into the plastic cylinder of the 'freeze-clamping' apparatus (QUISTORFF, 1975; NORBERG *et al.*, 1975). The cylinder was initially flushed with room air at a rate of 5 l/min. After allowing a steady period of 15–20 min an arterial blood sample was taken for control of blood gases and pH. The 'flushing gas-mixture' was then changed to 20% CO₂ and 20% O₂ in nitrogen, and after about 10 s of hypercapnia changed to 10% CO₂ and 20% O₂ in nitrogen. Brain tissue was sampled for analysis by means of 'freeze-clamping' following 10, 20 and 60 s of hypercapnia. An arterial blood sample was drawn from all animals for determination of P_{O₂} and P_{CO₂} at the moment of sampling.

In the anaesthetized series anaesthesia was induced with 2–3% halothane to allow tracheotomy and immobilization with tubocurarine chloride. The animals were maintained on 70% N₂O and 30% O₂ and ventilated to an arterial P_{CO₂} of 35–40 mm Hg. One femoral artery was cannulated for blood pressure recording and for anaerobic sampling of arterial blood. A skin incision was made over the intact skull bone to accommodate a plastic funnel for freezing of the brain *in situ*.

About 20–30 min after the operative procedure, when the animals were at steady state, the composition of the inspired gas was changed to include either 5 or 45% CO₂. Animals given 5% CO₂ were maintained on 25% O₂ and 70% N₂O. In experiments involving exposure to 45% CO₂ the CO₂ concentration of the gas was increased gradually over a 2 min period (to counteract a fall in blood pressure). The final composition of the inspired gas was 45% CO₂ and 30% O₂ in N₂. Thus, since 45% CO₂ induces anaesthesia nitrous oxide was withdrawn.

Freezing of the tissue *in situ* (PONTÉN *et al.*, 1973) was started 2, 5, 15, 30 and 60 min after the inspired CO₂ tension had been increased in the 5% group and after 5 and 15 min in the 45% CO₂ group. Arterial blood was drawn to determine P_{O₂}, P_{CO₂} and haemoglobin concentration.

The brains of animals from the unanaesthetized and anaesthetized series were dissected at –22°C in a refrigerated glove box. A slice of 1–1.5 mm thickness was cut out from the surface of the freeze-clamped samples by means of a small buzz saw (QUISTORFF, 1975). Cortical tissue from this thin slice was used for analysis.

From the anaesthetized animals the superior-lateral cortical areas of one hemisphere were cut out. All samples were weighed and extracted with HCl-methanol. The HCl-methanol extracts were then brought to 0°C and the tissue was extracted twice with perchloric acid for subsequent enzymatic fluorometric analyses of glucose, glycogen, glucose-6-phosphate (G-6-P), fructose-6-phosphate (F-6-P), fructose-1,6-diphosphate (FDP), dihydroxyacetone phosphate (DHAP), 3-phosphoglycerate (3-PG), lactate, pyruvate, citrate, isocitrate, α -ketoglutarate, succinate, fumarate, malate, aspartate, glutamate, glutamine, asparagine, alanine, GABA, ammonia, phosphocreatine (PCr), ATP, ADP and AMP. The methods used were those of LOWRY & PASSONNEAU (1972). The analytical conditions have been described in previous communications from the laboratory (FOLBERGROVÁ *et al.*, 1972a,b, 1974a,b). The oxaloacetate concentrations was calculated from the aspartate amino-

transferase reaction, using an equilibrium constant of 6.7 (KREBS & KORNBERG, 1957).

The results were statistically evaluated using two-sample *t*-test or Aspin-Welch's test. The following symbols are used: * $P < 0.05$, † $P < 0.01$, ‡ $P < 0.001$.

RESULTS

Unanaesthetized animals

One control group and 3 hypercapnic groups (10, 20 and 60 s, respectively) with 6 animals in each group were studied. The control values for arterial P_{CO₂} and P_{O₂} were 38.3 ± 0.9 and 98.8 ± 2.4 mm Hg, respectively (means $\pm$ S.E.M.). In the 3 hypercapnic groups P_{CO₂} was 77.6 ± 0.7 , 72.2 ± 1.8 and 69.4 ± 2.3 mm Hg, respectively, and arterial P_{O₂} exceeded 95 mm Hg in each animal.

As stated, the unanaesthetized animals were used to evaluate the initial changes in metabolites studied. However, since hypercapnia did not induce significant changes in organic phosphates or amino acids and since the only citric acid cycle intermediate showing a significant change was malate (a decrease from 0.309 ± 0.006 to 0.252 ± 0.006 μ mol/g at 60 s, $P < 0.001$) only the glycolytic metabolites will be dealt with.

Figure 1 illustrates the pattern of changes in glycolytic intermediates at 10 and 20 s (upper panel) and at 60 s (lower panel). At 60 s, there were no significant changes in glycogen or glucose, highly significant

FIG. 1. Changes in the concentrations of glycolytic metabolites in the rat cerebral cortex after 10, 20 and 60 s of hypercapnia (glycogen, glucose, glucose-6-phosphate, fructose-6-phosphate, fructose-1,6-diphosphate, dihydroxyacetone phosphate, 3-phosphoglycerate and pyruvate). The values are means $\pm$ S.E.M. Filled symbols indicate values significantly different from the controls ($P < 0.05$).

changes in G-6-P and F-6-P, and highly significant decreases in 3-PG and pyruvate. The data obtained at shorter exposure periods demonstrate that there was a transient decrease in glycogen concentration, that the concentrations of G-6-P and F-6-P increased during the first 20s, and that the pyruvate concentration was significantly reduced already at 10s. Two sets of results are relevant to the question of the mechanisms of the increases in G-6-P and F-6-P and of the decreases in pyruvate and lactate (not shown), respectively. First, when the hypercapnia was prolonged beyond the 10s period there were further increases in G-6-P and F-6-P whereas the glycogen concentration returned at 60s to normal values. Thus, it appears unlikely that G-6-P and F-6-P accumulated as a result of breakdown of glycogen (see Discussion and FOLBERGROVÁ *et al.*, 1974a). Second, the results obtained are pertinent to the mechanisms of the decrease in pyruvate concentration. Thus, the tissue lactate concentration remained unchanged at 10 and 20s but decreased from a control value of 0.85 ± 0.02 to 0.62 ± 0.03 at 60s. Furthermore, the blood lactate and pyruvate concentrations did not decrease during the course of hypercapnia (not shown). Thus, the decrease in tissue pyruvate concentration could neither be explained in terms of a shift in the lactate dehydrogenase reaction, nor were there concentration gradients favouring transport of lactate or pyruvate from tissue to blood.

In summary, the results obtained on the unanaesthetized animals showed that during the first minute of hypercapnia there is a transient decrease in glycogen concentration, progressive increases in G-6-P and F-6-P, and decreases in pyruvate and lactate. These events precede changes occurring in citric acid cycle intermediates and amino acids, and are unaccompanied by alterations in levels of PCr, ATP, ADP and AMP.

Anaesthetized animals

In all animals the body temperature was close to 37°C, the arterial P_{O_2} exceeded 100mm Hg, and the blood pressure was 140mm Hg, or higher. In the 5% CO_2 series the arterial CO_2 tension increased from a control value of 37.2 ± 1.1 mm Hg to 63.9 ± 1.8 , 69.5 ± 1.2 , 68.7 ± 0.1 , 72.8 ± 0.6 and 76.1 ± 0.9 in the 2, 5, 15, 30 and 60 min groups, respectively. With 45% CO_2 the arterial CO_2 tension was 310 ± 9 and 334 ± 4 at 5 and 15 min, respectively.

Figure 2 illustrates the pattern of changes in citric acid cycle intermediates at 5 and 15 min of hypercapnia (5 and 45% CO_2). With 5% CO_2 the results obtained after 2, 30 and 60 min were very similar to those shown for 5 and 15 min. The results confirm those of our previous reports in showing decreases in citrate, α -KG, malate and oxaloacetate that were proportional to the rise in CO_2 tension (see FOLBERGROVÁ *et al.*, 1972b, 1974a). In addition, they show that there was a reduction in fumarate and a small rise in succinate. The increase in succinate was statis-

FIG. 2. The pattern of citric acid cycle intermediates (pyruvate, citrate, isocitrate, α -ketoglutarate, succinate, fumarate, malate and oxaloacetate) after 5 and 15 min of 5 and 45% hypercapnia as compared to the normocapnic situation. The results are means $\pm$ S.E.M.

tically significant for the groups exposed to 45% CO_2 for 5 and 15 min ($P < 0.01$) but not for the 5% CO_2 groups.

The rise in succinate was not sufficient to balance the reduction in concentration of other cycle intermediates. Thus, in the 45% CO_2 groups the size of the pool was reduced from 1.55 ± 0.04 μ mol/g to 1.29 ± 0.06 ($P < 0.01$) and 1.10 ± 0.02 ($P < 0.01$) μ mol/g at 5 and 15 min, respectively. With 5% CO_2 the pool size appeared to decrease although the changes were not statistically significant. The values were 1.58 ± 0.05 , 1.44 ± 0.07 , 1.35 ± 0.04 , 1.33 ± 0.08 , 1.36 ± 0.05 and 1.37 ± 0.05 μ mol/g in the control and the 2, 5, 15, 30 and 60 min groups, respectively.

The tissue concentrations of amino acids and ammonia are illustrated in Table 1. The changes in glutamate, aspartate, glutamine and ammonia confirm our previous results (FOLBERGROVÁ *et al.*, 1972b, 1974a). Thus, with 5% CO_2 there was a gradual fall in glutamate, a transient increase in aspartate and an increase in glutamine at 60 min. With 45% CO_2 glutamate fell by about 4.5 μ mol/g, aspartate increased by 2.3 μ mol/g and there was a significant rise in ammonia. GABA or asparagine did not change significantly with 5 or 45% CO_2 .

There were no significant changes in alanine concentration. Data obtained with 5% CO_2 suggested that a small fall occurred in hypercapnia. However, since there was no change with 45% CO_2 we conclude that alanine concentration remains essentially unaltered in hypercapnia.

TABLE 1. CONCENTRATIONS OF AMINO ACIDS AND AMMONIA IN THE CEREBRAL CORTEX IN RATS EXPOSED AT 5 AND 45% CO₂ AT DIFFERENT TIMES

Experimental group	Time (min)	Glut	Gln	Asp	Aspn	GABA	Ala	NH ₄ ⁺	Σ Amino acids
5% CO ₂	0	13.67 ±0.33	5.67 ±0.09	3.82 ±0.09	0.087 ±0.004	1.89 ±0.04	0.470 ±0.039	0.281 ±0.043	25.6 ±0.5
	2	12.88 ±0.25	6.11 ±0.28	4.62† ±0.13	0.088 ±0.002	1.83 ±0.03	0.442 ±0.023	0.251 ±0.04	26.0 ±0.3
	5	12.86 ±0.43	5.51 ±0.37	4.85† ±0.17	0.085 ±0.002	1.84 ±0.07	0.422 ±0.005	0.279 ±0.038	25.8 ±0.5
	15	12.55* ±0.26	6.04 ±0.44	4.62* ±0.25	0.085 ±0.003	1.90 ±0.02	0.396 ±0.025	0.244 ±0.029	25.6 ±0.8
	30	12.48 ±0.54	5.78 ±0.30	4.19 ±0.15	0.084 ±0.004	1.89 ±0.07	0.374 ±0.018	0.240 ±0.041	25.0 ±0.7
	60	11.92† ±0.28	6.43* ±0.19	3.74 ±0.13	0.087 ±0.004	1.95 ±0.06	0.397 ±0.035	0.281 ±0.058	24.5 ±1.1
	0	13.19 ±0.23	5.15 ±0.64	4.14 ±0.16	0.078 ±0.003	2.13 ±0.08	0.503 ±0.009	0.253 ±0.017	25.2 ±0.5
	5	10.31‡ ±0.14	4.59 ±0.40	6.19‡ ±0.19	0.086 ±0.006	2.12 ±0.14	0.615 ±0.044	0.330 ±0.037	23.4 ±0.7
	15	8.65‡ ±0.12	5.53 ±0.26	6.41‡ ±0.27	0.092 ±0.005	2.30 ±0.09	0.481 ±0.027	0.469† ±0.033	22.4* ±0.7

The values are means ± S.E.M. The abbreviations are as follows: Glut = glutamate, Gln = glutamine, Asp = aspartate, Aspn = asparagine, GABA = γ-aminobutyric acid, Ala = alanine. The concentrations are expressed in μmol/g wet tissue.

The results of Table 1 indicate that there is a decrease in the amino acid pool (Σ amino acids) during hypercapnic conditions. With 5% CO₂ this decrease was not significant although the results obtained at 30 and 60 min are suggestive. With 45% CO₂ the pool size decreased significantly and at 15 min this decrease amounted to 2.8 μmol/g. At this time there was also an increase in ammonia. The changes in pool size and ammonia indicate that oxidative deamination of amino acids may have occurred (see Discussion).

DISCUSSION

The new metabolites measured in the present anaesthetized series were isocitrate, succinate, fumarate, alanine, GABA and asparagine. The control values for succinate and fumarate are in good agreement with those reported by GOLDBERG *et al.* (1966) but the value for isocitrate was less than half of that given by these authors. In our series the extracts were not treated with Florosil for the isocitrate analyses and the fluorescent blank was too high to allow satisfactory analyses. However, the results were internally consistent and they are reported since they clearly demonstrate a fall in isocitrate levels in hypercapnia. Our values for GABA and alanine are in good agreement with those of other workers (see DUFFY *et al.*, 1972).

The results obtained in the present experiments have greatly facilitated the interpretation of those of the previous series (FOLBERGROVÁ *et al.*, 1972b, 1974a) and the combined results allow a tentative description of the sequence of metabolic events occurring in hypercapnia. This description must remain somewhat speculative and some aspects will require validation with other methods. However, the pattern is sufficiently coherent to be presented as a framework on which subsequent experiments can be founded. Our interpretation is based on the following two facts. (1)

The initial changes affect glycolytic metabolites, and alterations in citric acid cycle intermediates and amino acids represent secondary events. (2) The citric acid cycle pool is decreased. Thus, the balance between reactions that replenish carbon skeletons to the citric acid cycle and those leading to loss of carbon atoms is shifted. Furthermore, estimation of the size of the amino acid pool, and measurement of ammonia content, indicate that amino acids are probably used as substrates during hypercapnia. When discussing the probable mechanisms involved we will make two assumptions. First, the fact that the metabolism of citric acid cycle intermediates and amino acids is compartmented in the brain (BALÁZS & CREMER, 1973) does not influence the conclusions. Second, changes occurring during exposure to 5 and to 45% CO₂ are qualitatively similar and data obtained with 45% CO₂ can therefore be used to demonstrate changes that may escape detection with 5% CO₂. We will discuss in turn glycolytic events, and those affecting citric acid cycle intermediates and amino acids.

Glycolytic events. Of key importance is the fact that pyruvate decreases already after 10 s of hypercapnia. Such a decrease could occur as a result of decreased delivery or increased utilization. In all probability, the second possibility can be excluded. First, the metabolic rate for oxygen remains unchanged at P_{CO₂}, values of 80–90 mm Hg (EKLÖF *et al.*, 1973). Second, there is no indication that pyruvate or phosphoenolpyruvate is removed by accelerated CO₂ fixation. Thus, instead of increasing, the citric acid cycle pool decreased. Furthermore, since the main CO₂-fixing enzyme in the brain is probably pyruvate carboxylase (PATEL, 1974) the rate of CO₂ fixation could vary with the pyruvate concentration, rather than with the CO₂ tension or the bicarbonate concentration (see CHENG, 1971). Third, analyses of the alanine concentration give no support to the possibility that pyruvate is

lost due to a shift in the alanine aminotransferase reaction. Fourth, the results make it less likely that lactate or pyruvate are lost from the tissue by means of transport to the blood.

Accepting that the decrease in pyruvate concentration is due to decreased delivery we must enquire into the possible mechanisms. The present results have shown that there is an accumulation of G-6-P and F-6-P during the first minute of hypercapnia. Since this accumulation does not seem to be secondary to glycogen breakdown (see also FOLBERGROVÁ *et al.*, 1974a) it may signify inhibition of phosphofructokinase (PFK). Unequivocal proof of PFK inhibition would require that the glycolytic flux could be shown to be decreased (KREBS, 1957). Since CMR_{O_2} is unchanged, at least at moderate degrees of hypercapnia (see above) a decreased glycolytic flux should mean that non-carbohydrate substrates are oxidized. Since the amino acid changes indicate that this may be the case (see below) we tentatively conclude that there is a decreased glycolytic flux due to PFK inhibition. This conclusion is entirely in keeping with previous observations on the effect of a decreased pH on PFK activity (DANFORTH, 1965).

Changes affecting citric acid cycle intermediates and amino acids. A decrease in pyruvate concentration due to diminished production implies that a condition of 'substrate depletion' is at hand. Since CMR_{O_2} is unchanged, at least at moderate degrees of hypercapnia, a decreased glycolytic flux should mean that non-carbohydrate substrates are oxidized. In fact, substrate depletion during hypercapnia has some striking similarities with hypoglycaemia (LEWIS *et al.*, 1974a; NORBERG & SIESJÖ, 1975). In both conditions there is a decrease in the size of the TCA pool, a shift in the aspartate aminotransferase reaction towards aspartate formation, a decrease of the amino acid pool, and an increase in ammonia. These events suggest that the non-carbohydrate substrate oxidized during hypercapnia and hypoglycaemia are at least in part tricarboxylic acid intermediates and amino acids. Normally, carbon skeletons are lost from the citric acid cycle via synthetic reactions, e.g. these leading to formation of GABA and glutamine. This loss is probably mainly replenished through CO_2 fixation, and it has been estimated that about 10% of the phosphoenolpyruvate or pyruvate is diverted towards formation of malate or OAA (CHENG, 1971; BALÁZS *et al.*, 1973). Thus, a net flux of carbon atoms is maintained from the glycolytic chain to the citric acid cycle and from the latter to the amino acid pools. However, when less pyruvate is available for oxidation there is probably a relative build-up of OAA that may have two effects. Firstly, by inducing a shift in the aspartate aminotransferase reaction it provides amino acid carbon for oxidation (see DAWSON, 1950). Secondly, by allowing a reversal of the CO_2 fixation reaction it makes the citric acid cycle intermediates available for oxidation. In itself, the citric acid cycle pool is small and would not contribute significantly to the sub-

strate supply. However, whether amino acids are metabolized via the glutamate dehydrogenase reaction (see WEIL-MALHERBE, 1971) or via the reactions of the purine nucleotide cycle (LOWENSTEIN, 1972) their carbon skeletons enter the citric acid cycle before being oxidized. Thus, in hypercapnia and in hypoglycaemia the reduced availability of pyruvate seems to cause a net flux of carbon skeletons from amino acids to the citric acid cycle, and from the latter to the glycolytic chain.

Although the main metabolic events in hypercapnia can be satisfactorily explained there are some dissimilarities between hypercapnia and hypoglycaemia that warrant a discussion, and the mechanisms responsible for some of the changes observed must remain speculative. We will discuss the fate of aspartate, α -KG, alanine, glutamine and GABA.

In hypoglycaemia and in hypercapnia aspartate accumulates at the expense of glutamate, indicating that there is a shift in the aspartate aminotransferase reaction. In severe hypoglycaemia there is a massive accumulation of aspartate whereas in hypercapnia a secondary fall in aspartate to normal or subnormal values occurs (cf. FOLBERGROVÁ *et al.*, 1974a). In both conditions the accumulation of aspartate may be secondary to a relative or absolute increase in OAA. Further metabolism of aspartate would either require a reversal of the aspartate aminotransferase reaction and the subsequent deamination of glutamate, or deamination by the reactions of the purine nucleotide cycle. In hypoglycaemia, transamination could be blocked by the persisting rise in OAA caused by a depletion of NADH and a fall in malate/OAA ratio, and metabolism via the purine nucleotide cycle could be inhibited by lack of GTP (see NORBERG & SIESJÖ, 1975). Apparently, one or both of these pathways operate in hypercapnia.

In both hypercapnia and hypoglycaemia a rise in OAA may thus be one of the mechanisms eliciting a shift in the aspartate aminotransferase reaction. However, since α -KG does not accumulate other reactions must accomplish α -KG removal. In hypercapnia, carboxylation of α -KG to isocitrate could be one of these reactions (see CHENG, 1971). In fact, a reversal of the isocitrate dehydrogenase reaction may, in removing α -KG, by itself induce transamination. There is thus no need to invoke reductive amination of α -KG to explain why its concentration is maintained low. Instead, since ammonia accumulates when there is a decrease in the pool size of amino acids, the conditions seem to favour oxidative deamination of glutamate.

In hypoglycaemia, there are relatively marked decreases in alanine, GABA and glutamine. If carboxylation of α -KG occurs in hypercapnia lack of α -KG could well explain why alanine is not utilized via the alanine aminotransferase reaction, and why GABA is not metabolized. In hypoglycaemia, there appears to be a pronounced redox change in the direction of oxidation (see LEWIS *et al.*, 1974a). Lack of NADH,

and accumulation of NAD^+ , could therefore enhance GABA metabolism by removing succinic semialdehyde. Thus, CO_2 fixation at the α -KG level and absence of a marked redox change could explain why alanine and GABA are not used as substrates in hypercapnia. There is no ready explanation for the delayed increase in glutamine under hypercapnic conditions. However, one factor of importance may be that accumulation of ammonia occurs without depletion of ATP and the conditions may therefore favour amidation of glutamate. In hypoglycaemia, ammonia accumulation is accompanied by energy failure (LEWIS *et al.*, 1974b) and glutamine formation could be blocked.

In summary, although the detailed mechanisms are largely unknown the present results indicate that hypercapnia leads to a relative deficiency of pyruvate and that the changes observed can be at least partly explained by the fact that citric acid cycle intermediates and amino acids are used as alternative substrates.

Acknowledgements—This study was supported by grants from the Swedish Medical Research Council (projects No 14X-263 and 14X-2179), from the Swedish Bank Tercentenary Fund, from U.S. PHS Grant No. R01 SO 7838-05 from NIH, and from the Swedish Board for Technical Development.

REFERENCES

- BALÁZS R. & CREMER J. E. (1973) *Metabolic Compartmentation in the Brain* (BALÁZS R. & CREMER J. E., eds.). Macmillan, London.
- BALÁZS R., MACHİYAMA Y. & PATEL A. J. (1973) in *Metabolic Compartmentation in the Brain* (BALÁZS R. & CREMER J. E., eds.) pp. 55–70. Macmillan, London.
- CHENG S.-C. (1971) *Int. Rev. Neurobiol.* **14**, 125–157.
- DANFORTH W. H. (1965) in *Control of Energy Metabolism* (CHANCE B., ESTABROOK C. W. & WILLIAMSON J. K., eds.) pp. 287–297. Academic Press, New York.
- DAWSON R. M. C. (1950) *Biochem. J.* **47**, 386–395.
- DUFFY T. E., NELSON S. R. & LOWRY O. H. (1972) *J. Neurochem.* **19**, 959–977.
- EKLÖF B., LASSEN N. A., NILSSON L., NORBERG K. & SIESJÖ B. K. (1973) *Acta physiol. scand.* **88**, 587–589.
- FOLBERGROVÁ J., MACMILLAN V. & SIESJÖ B. K. (1972a) *J. Neurochem.* **19**, 2497–2505.
- FOLBERGROVÁ J., MACMILLAN V. & SIESJÖ B. K. (1972b) *J. Neurochem.* **19**, 2507–2517.
- FOLBERGROVÁ J., PONTÉN U. & SIESJÖ B. K. (1974a) *J. Neurochem.* **22**, 1115–1125.
- FOLBERGROVÁ J., LJUNGGREN B., NORBERG K. & SIESJÖ B. K. (1974b) *Brain Res.* **80**, 265–279.
- GOLDBERG N. D., PASSONNEAU J. V. & LOWRY O. H. (1966) *J. biol. Chem.* **241**, 3997–4003.
- KREBS H. A. & KORNBERG H. L. (1957) *Energy Transformations in Living Matter*, pp. 213–285. Springer Verlag, Berlin.
- LEWIS L. D., LJUNGGREN B., NORBERG K. & SIESJÖ B. K. (1974a) *J. Neurochem.* **23**, 659–671.
- LEWIS L. D., LJUNGGREN B., RATCHESON R. A. & SIESJÖ B. K. (1974b) *J. Neurochem.* **23**, 673–679.
- LOWENSTEIN J. M. (1972) *Physiol. Rev.* **52**, 382–414.
- LOWRY O. H. & PASSONNEAU J. V. (1972) *A Flexible System of Enzymatic Analysis*. Academic Press, New York.
- NORBERG K. & SIESJÖ B. K. (1975) *J. Neurochem.* in press.
- NORBERG K., QUISTORFF B. & SIESJÖ B. K. (1975) *Acta Physiol. scand.* Submitted for publication.
- PATEL M. S. (1974) *J. Neurochem.* **22**, 717–724.
- PONTÉN U., RATCHESON R. A., SALFORD L.-G. & SIESJÖ B. K. (1973) *J. Neurochem.* **21**, 1127–1138.
- QUISTORFF B. (1975) *Analyt. Biochem.* In press.
- WEIL-MALHERBE H. & GORDON J. (1971) *J. Neurochem.* **18**, 1659–1672.