The burden of domestication

a representative study of welfare in privately owned cats in Denmark

Sandøe, Peter; Nørspang, Annika Patursson; Forkman, Björn; Bjørnvad, Charlotte Reinhard; Kondrup, Sara Vincentzen; Lund, Thomas Bøker

Published in:
Animal Welfare

DOI:
[10.7120/09627286.26.1.001](https://doi.org/10.7120/09627286.26.1.001)

Publication date:
2017

Document version
Peer reviewed version

Citation for published version (APA):

Sandøe, P., Nørspang, A. P., Forkman, B., Bjørnvad, C. R., Kondrup, S. V., & Lund, T. B. (2017). The burden of domestication: a representative study of welfare in privately owned cats in Denmark. *Animal Welfare*, 26(1), 1-10. <https://doi.org/10.7120/09627286.26.1.001>

The burden of domestication – A representative study of welfare in privately owned cats in Denmark[§]

Peter Sandøe^{*1,2}, Annika Patursson Nørspang¹, Björn Forkman¹, Charlotte Reinhard Bjørnvad³, Sara Vincentzen Kondrup² & Thomas Bøker Lund²

*Corresponding author: pes@sund.ku.dk, +4535333059. ¹University of Copenhagen, Department of Large Animal Sciences, Grønnegårdsvej 8, 1870 Frederiksberg C., Denmark; ²University of Copenhagen, Department of Food and Resource Economics, Rolighedsvej 25, 1958 Frederiksberg C., Denmark; ³University of Copenhagen, Department of Veterinary Clinical and Animal Sciences, Dyrslægevej 16, 1870 Frederiksberg C., Denmark

Abstract

The way in which domestic cats are kept and bred has changed dramatically over the last two centuries. Notably a significant number of cats are kept indoors, most of them are neutered and many are selectively bred. This likely has consequences for their welfare. A few studies link housing, neuter status and breeding in cats to risks of welfare problems. However, the study presented here is the first to quantify the risks and document the prevalence of risk factors. It builds on results from a questionnaire sent to a representative sample of the Danish population. Using the responses from cat owners who keep cats in the home (N=378) the paper aims to investigate how indoor confinement, neutering and selective breeding affect health, behaviour and other factors relating to cat welfare. The paper reports that confined cats had significantly more behavioural problems than free-roaming cats; that a smaller proportion of the free-roaming cats suffered from the behavioural problems investigated; and that entire cats had significantly more behavioural problems than neutered cats. Finally, significantly more purebred cats than domestic shorthair cats were found to have diseases. Being confined, living as an intact female and being purebred are therefore significant risk factors for behavioural or health problems associated with reduced welfare in privately owned cats.

Keywords: animal welfare, behavioural problems, confinement, health issues, neuter status, purebred cats

Introduction

In the last fifty years there has been a dramatic increase in the number of people who keep cats as companion animals. For example, in the UK between 1965 and 2004 the number of cats kept per hundred inhabitants increased from fewer than 8 to more than 16 (Sandøe et al 2016a Ch 1). Today in most Western societies cats enjoy popularity as companions that is comparable to dogs, and in

[§] The reference of the printed version is:

Sandøe, P., Nørspang, A. P., Forkman, B., Bjørnvad, C. R., Kondrup, S. V., & Lund, T. B. (2017). The burden of domestication: a representative study of welfare in privately owned cats in Denmark. *Animal Welfare*, 26(1), 1-10. The definitive version is available at <https://doi.org/10.7120/09627286.26.1.001>

Europe there are more households with a cat than households with a dog (FEDIAF 2014). In Denmark, although there are fewer households with cats than with dogs, the total number of domestic cats kept is higher than that of dogs (Danmarks Statistik 2000), as is the case in the United States (AVMA 2012), whereas in Australia domestic cats are reported to be the second most common companion animal, with numbers just below those of dogs (Richmond 2013). Over this period of rapidly rising popularity of the cat as a companion animal, dramatic changes in the way cats are bred, kept and cared for have taken place. However, little is known about how these changes affect the welfare of the cats.

One important shift concerns the housing of cats. Today many cats are confined indoors; seemingly this happens more commonly in the US than in Europe (Rochlitz 2005; Bayer 2013). Confinement protects the cat from road accidents, injuries from fights and other dangers, but it may at the same time prevent the cat from performing important natural behaviours (Palmer & Sandøe 2014). It appears that it also puts the cat in higher risk of developing certain diseases (Robertson 1999; Rand et al 2004; Slingerland et al 2009). The main cause of this is believed to be an inactive lifestyle, which can put individuals at greater risk of developing certain lifestyle related diseases. Similarly, behavioural problems have been linked to confinement, and specifically cats' inactivity and their uniform, unchanging life and environment (Heidenberger 1997; Amat et al 2009; Bain & Stelow 2014). A bored or stressed cat might also perform unwanted behaviour, such as excessive vocalization, aggressiveness or house soiling.

A second shift is that the majority of domestic cats are now neutered (Chu et al 2009; Sandøe et al 2016b). The surgical removal of reproductive organs permanently to prevent cats from breeding has an impact on welfare for various reasons. The neutered cat needs to undergo surgery and recovery, and complications may develop from anaesthesia or surgical trauma. In the longer term neutering increases the risk of obesity, which can lead to diabetes and other health-related diseases (Robertson 1999; Rand et al 2004; Colliard et al 2009). On the other side, neutering also protects cats from disease; and in both males and females it seems to reduce aggressiveness (Finkler & Terkel 2010). In males, neutering significantly reduces roaming and fighting activity, thereby indirectly reducing the risk of traffic injuries and bite wounds with associated complications such as abscesses, transmission of feline immunodeficiency virus (FIV) and feline leukemia virus (FeLV) (Hart & Barrett 1973). Neutering in female cats reduces the risk of developing estrogen responsive mammary tumours (Overley et al 2005) and it indirectly decreases malnutrition and disease in kittens, by reducing the population density in a given area.

The third major shift of note concerns organized cat breeding for specific traits. Although feline domestication has existed for long period, selective breeding of purebred cats is relatively recent (O'Brien & Johnson 2007). Despite the good intentions of breeders pure-breeding can have a negative effect on animal welfare. Sandøe and others (2016a Ch 7) have divided these negative effects into three groups: breeding of extreme phenotypes which in themselves create health and welfare problems; increased prevalence of diseases caused by a lack of genetic diversity; and increased prevalence of behavioural problems.

In sum, then, domestic cats may suffer from a number of serious welfare problems as a result of the way they are housed, taken care of, and bred. However, to the authors' knowledge no representative studies which link different factors in the breeding and lifestyle of companion cats to the prevalence of welfare problems in the domestic cat population have been published. The purpose of the current study is to make a start on filling this research gap.

The paper is based on results from a questionnaire sent to a representative sample of the Danish population. Based on the responses of cat-owning respondents (n=415) we examine how indoor confinement, neuter status and selective breeding affect cat health and behaviour.

The paper does not look into effects selective breeding on specific breeds but only looks at how purebred cats as a group are affected. To look at specific breeds would not have been feasible given the limited number of owners of purebred cats participating in the questionnaire and the wide diversity of breeds owned. However, the study does seek to clarify whether purebred cats as a group have a higher frequency of behavioural and health problems than domestic shorthair cats and/or mixed breed cats.

The study uses results relating to Danish domestic cats, but the correlations it identifies can most likely be generalized beyond Denmark, and its findings should therefore be of interest to authorities, private organizations, veterinarians and other professionals with an interest in the welfare of privately owned cats worldwide.

Material and methods

Survey design

The data are based on a survey containing 45 questions (Questionnaire 1; 2). All participants were asked a number of demographic questions as well as questions relating to their attitudes to cats in general, to roaming and to stray cats. Additional questions were given to cat owners about their cat, such as age, breed, gender, neuter status, regarding the number of cats in the household, the cat's behaviour, health status, the type of activities provided by the owner, environmental enrichment, feeding, veterinary care, housing of the cat and potential problems related to housing. Data collection was carried out by a Danish survey company (Norstat) in October 2015. The respondents belonged to Norstat's pre-recruited panel. A gross sample from this panel (N=6120) was invited to participate in the survey. Sampling had quotas on age, gender and geography (NUTS2 regional level) according to Danish census data.

A combined online and telephonic design was performed. Respondents aged between 18-64 years responded online and respondents over the age of 65 years were interviewed by phone. This mixed mode design was chosen with a view to obtaining a high degree of representativeness while holding data collection costs down. Specifically, while Internet data collection is less expensive, it is known that Danes in the +65 age segment use the Internet less frequently and are best reached by

telephone. The final, net sample was N=2003, resulting in a response rate of 33%. To account for non-response bias the cases were weighted according to official statistics on gender x age x region. Of the 2003 people who responded, 415 (weighted frequency) were cat owners. Cat owners were instructed to complete the survey for their oldest cat, and those responses form the basis of the results presented here.

Statistical analysis

Descriptive analyses were conducted to display the prevalence of cats in Denmark, breed type, neutering status, how the cats were kept, behavioural problems, and health issues (i.e. the cat being overweight or suffering from one of the following diseases: arthritis, oral disease, kidney disease, urinary disease, diabetes, metabolic disease). After this the data were analysed to discover whether the main explanatory variables under investigation, i.e. breed type, neutering status, and how cats are kept, were associated with behavioural problems. This was done by reporting the unadjusted prevalence of behavioural problems across the different categories of the explanatory variables. Following this, odds ratio results were reported (with 95% CI) from multivariate logistic regression models for each main explanatory variable after adjustment for the age of the cats (cat age was inserted as a categorical variable with 5 brackets (“0-1 year” to “more than 7 years”) and the two other main explanatory variables. The data were also analysed to find whether breed type and neutering status were associated with health issues. Results from this were again reported with prevalence and odds ratio results (with 95% CI) from logistic regression models after adjustment for the age of the cats and the two other main explanatory variables.

Confined cats were defined as cats that are either indoor cats with no access to the outdoors or indoor cats with limited access to the outdoors part of the year (e.g. in a summer house). Garden cats were defined as cats with access to a closed garden, and free-roaming cats were defined as indoor cats with the opportunity to roam freely outside. A final category was outdoor cats that rarely or never came inside the house. Apart from featuring in the demographic characteristics of the cat population (Table 1) these outdoor cats were excluded from the study, as welfare problems are difficult to study in them.

All statistical analyses were performed using IBM SPSS Statistics Version 21. In all analyses, statistical significant difference was set at the 95% level.

Results

Size of the cat owner population

In all 2003 people were asked if they had cats (i.e. at least one cat) in their household or had previously done so. The majority of people, 1327 (66.2%), answered negatively, 261 (13.0%) people had previously had a cat in the household, and 415 (20.7%) persons currently had at least one cat. Thus 33.7% of the sampled Danish households keep or have previously kept at least one cat.

Demographic characteristics of the cat population

Table 1 displays the distribution of the three variables that are linked in this paper to behavioural and health problems in privately owned cats: breed of the cat, neuter status, and housing type.

Table 1. Prevalence of cat breeds, sex/neutering status, and housing in Danish households with cats	
Breed (N=415)	
Domestic shorthair	61.5%
Mixed breed	20.5%
Purebred	15%
Don't know	3%
Sex/neutering status (N=415)	
Intact male	3.9%
Intact female	9.2%
Neutered male	47%
Neutered female	39.1%
Don't know	0.9%
How cats are kept (N=415)	
No outdoor access ^a	16.8%
Only outdoor access part of the time ^a	3.6%
Access to a closed garden ^b	7.8%
Outdoor access through cat flap ^c	25%
Outdoor access when owner lets the cat out ^c	38%
Outdoor cats that rarely or seldom come inside	8.8%
How cats are kept (reduced sample: N=378)	
Confined	22.4%
Garden access	8.5%
Free-roaming	69.1%
^a Defined as confined in the remainder of the analysis;	
^b Defined as garden-cat in the remainder of the analysis;	
^c Defined as free-roaming in the remainder of the analysis	

As can be seen 15% of domestic cats in Denmark are purebred. The prevalence of the reported breeds were as follows: Abyssinian 0.7%, British Shorthair 0.5%, Burmese 1.2%, European shorthair 0.5%, Birman 0.7%, Maine Coon 3.9%, Norwegian Forest cat 2.9%, Persian 1.2%, Ragdoll 0.7%, Russian Blue 0.5%, Siamese/Oriental shorthair 0.5%, other breeds 1.5%; 0.2% of

respondents did not know which breed their purebred cat was. Since most of the breeds involve small numbers of individuals we decided to treat purebred cats as a single undifferentiated group.

Regarding neuter status it can be seen from the survey that 86% of the cats are neutered while 4% of the male cats and 9% of the female cats are intact. Owners house their cats in a wide variety of ways. Around 9% of cats live more or less permanently outdoors, and since it is likely that the owners have limited knowledge of their cats' welfare and behaviour they are excluded from the remainder of the study. Among the remaining cats, a significant minority, 22%, are classified as 'confined' in that they never have access to outdoors or are only allowed out on specific occasions (e.g. when the family is in a summer house), and a small fraction, 9%, have access to an enclosed garden ('garden access'), but the large majority, 69%, are allowed to roam freely outdoors either by using a cat flap or by being let out by the owner ('free-roaming').

Prevalence of behavioural problems and health issues

When asked about behavioural problems 21.7% of owners report that their cat damages furniture or things, e.g. by scratching, 15.1% report fear of other cats, dogs or people, 12.4% report problems with house soiling, 11.0% report that their cat displays signs of boredom, e.g. excessive vocalization, 4.4% report aggressive behaviour towards owner, 4.0% towards guests, and 3.8% towards other pets in the household. Furthermore 5.4% report that they have "other behaviour problems". About half of the owners report that they have none of the problems listed. In sum it can, based on our study, be said that half of all cats in Denmark shows one or more behaviours that the owner views as a problem.

When asked about their cats' health, owners reported the following: 9.5% of the cats are overweight, 4.9% have arthritis, 4.0% have oral disease, 2.1% have kidney disease, 2.1% have urinary disease, 1.3% have diabetes, and 0.6% have metabolic disease. A large proportion, 75.9%, of the cats were reported by their owners as having none of the mentioned diseases, and just 2.9% of the owners did not know if their cat had any of the mentioned diseases. Except for being overweight, the health problems at issue were reported quite infrequently. In the remainder of the analysis diabetes, arthritis, oral disease, kidney disease, urinary disease and metabolic disease were therefore collapsed into one variable: disease. After this, as can be seen from Table 5 (Total column), which displays the prevalence of health issues, 13.8% of the cats were found to have one or more of these diseases, 10.3% of the cats were overweight, whilst 75.2% had none of these conditions.

What explains behavioural problems?

Table 2 reports prevalence of the behavioural problem for confined, garden and free-roaming cats.

Table 2. Prevalence of behavioural problems in confined, garden and free-roaming cats (and total) as reported by the owners in a representative study of Danish cats. N = 378									
	Prevalence (in %)			Total	Adjusted OR (95% CI)*				
	Confined	Garden	Free- roaming		Confined		Garden		
					OR	(95% CI)	OR	(95% CI)	
House soiling	18.2%	18.4%	9.8%	12.4%	2.38	(1.12 5.02)	4.17	(1.41 12.32)	
Damage furniture or things	35.5%	25.1%	16.8%	21.7%	2.44	(1.35 4.42)	2.10	(0.85 5.21)	
Aggressive behaviour towards owner	5.4%	11.4%	3.2%	4.4%	1.08	(0.28 4.13)	2.56	(0.61 10.69)	
Aggressive behaviour towards guests	6.6%	4.0%	3.1%	4.0%	1.99	(0.58 6.87)	1.57	(0.21 11.54)	
Aggressive behaviour towards other pets in the household	5.1%	3.6%	3.4%	3.8%	0.70	(0.15 3.29)	0.78	(0.10 6.14)	
Displays signs of boredom	19.1%	6.1%	9.0%	11.0%	2.63	(1.28 5.40)	0.65	(0.14 3.10)	
Fears other cats, dogs or people	13.1%	8.6%	16.5%	15.1%	0.93	(0.44 1.94)	0.71	(0.19 2.65)	
Other problems	9.5%	3.3%	4.4%	5.4%	3.03	(1.14 8.09)	1.21	(0.15 9.62)	
None of the problems listed	37.1%	44.7%	54.7%	49.9%	0.51	(0.30 0.86)	0.59	(0.27 1.29)	
*Results from binary logistic regression (N=359), where 'free-roaming' cat was set as reference value. Adjusted for age of the cat, breed type, and neutering status.									

Significantly more owners of free-roaming cats than those of confined cats report that their cats have none of the studied behavioural problems. Specifically, confinement increases the probability of house soiling and that the cat damages furniture or other things and displays signs of boredom.

On the other hand, few differences in behavioural problems were seen between purebreds, domestic shorthair and mixed breed cats (**Table 3**).

Table 3. Prevalence of behavioural problems in purebred, domestic shorthair and mixed breed cats (and total) as reported by the owners in a representative study of Danish cats. N = 367									
	Prevalence (in %)			Total	Adjusted OR (95% CI)*				
	Purebred	Mixed breed	Domestic shorthair		Purebred		Mixed breed		
					OR	(95% CI)	OR	(95% CI)	
House soiling	1.4%	12.2%	14.5%	12.4%	0.06	(0.01 0.55)	0.93	(0.41 2.13)	
Damage furniture or things	18.3%	26.7%	19.8%	21.7%	0.74	(0.34 1.58)	1.59	(0.84 3.01)	
Aggressive behaviour towards owner	9.1%	3.0%	3.3%	4.4%	2.23	(0.65 7.67)	0.81	(0.17 3.90)	
Aggressive behaviour towards guests	4.1%	5.7%	3.1%	4.0%	1.11	(0.24 5.25)	2.07	(0.57 7.52)	
Aggressive behaviour towards other pets in the household	6.8%	1.5%	3.0%	3.8%	2.36	(0.64 8.71)	0.53	(0.07 4.14)	
Displays signs of boredom	13.1%	17.0%	9.2%	11.0%	1.39	(0.56 3.48)	2.15	(0.98 4.71)	
Fears other cats, dogs or people	8.1%	25.6%	14.4%	15.1%	0.58	(0.21 1.60)	2.21	(1.13 4.32)	
Other problems	1.8%	7.3%	6.0%	5.4%	0.24	(0.03 1.81)	1.30	(0.43 3.87)	
None of the problems listed	47.9%	46.5%	52.2%	49.9%	0.98	(0.54 1.79)	0.76	(0.44 1.32)	
*Results from binary logistic regression (N=359), where 'domestic shorthair' cat was set as reference value. Adjusted for age of the cat, how the cat is kept, and neutering status									

However, significantly fewer purebred cats than domestic shorthair cats have house soiling problems; and significantly more mixed breed cats than domestic shorthair cats fear other cats, dogs or people.

In the analysis of associations between neutering status and behavioural problems (**Table 4**) non-neutered female and male cats were collapsed (non-neutered cat; N=41). Likewise neutered female and male cats were collapsed (N=335).

Table 4. Prevalence of behavioural problems in neutered and non-neutered cats (and total) as reported by the owners in a representative study of Danish cats. N = 376

	Prevalence (in %)			Adjusted OR (95% CI)*		
	Not neutered	Neutered	Total	Not neutered		
				OR	(95% CI)	
House soiling	14.6%	11.9%	12.4%	1.55	(0.56	4.28)
Damage furniture or things	11.9%	22.7%	21.7%	0.52	(0.19	1.45)
Aggressive behaviour towards owner	2.6%	4.6%	4.4%	0.41	(0.05	3.19)
Aggressive behaviour towards guests	10.7%	3.2%	4.0%	3.73	(1.06	13.16)
Aggressive behaviour towards other pets in the household	2.6%	4.0%	3.8%	0.80	(0.10	6.29)
Displays signs of boredom	17.8%	10.2%	11.0%	1.68	(0.65	4.35)
Fears other cats, dogs or people	17.5%	14.9%	15.1%	1.64	(0.64	4.19)
Other problems	12.9%	4.5%	5.4%	4.13	(1.32	12.97)
None of the problems listed	28.6%	52.5%	49.9%	0.32	(0.15	0.68)

*Results from binary logistic regression (N=359), where ‘neutered’ cat was set as reference value. Adjusted for age of the cat, how the cat is kept, and breed type

Significantly fewer owners of non-neutered cats report that none of the behavioural problems listed are present. Among specific problems, non-neutered cats have significantly more aggressive behaviour towards guests and “other problems”.

What explains health issues?

Table 5 displays the prevalence of health issues for domestic shorthair, purebred and mixed breed cats.

Table 5. Prevalence of overweight, disease or none of the diseases listed in purebred, domestic shorthair and mixed breed cats (and total) as reported by the owners in a representative study of Danish cats. N = 367

	Prevalence (in %)				Adjusted OR (95% CI)*					
	Purebred	Mixed breed	Domestic shorthair	Total	Purebred			Mixed breed		
					OR	(95% CI)		OR	(95% CI)	
Overweight	3.3%	15.5%	9.4%	10.3%	0.34	(0.08	1.49)	1.78	(0.80	3.96)
Disease**	36.1%	14.1%	7.2%	13.8%	7.11	(3.30	15.29)	1.95	(0.84	4.52)
None of the diseases listed	57.4%	69.0%	82.6%	75.2%	0.29	(0.15	0.55)	0.47	(0.26	0.88)

*Results from binary logistic regression (N=359), where ‘domestic shorthair’ cat was set as reference value. Adjusted for age of the cat, how the cat is kept, and neutering status
** Diseases prompted for: arthritis, oral disease, kidney disease, urinary disease, diabetes, metabolic disease.
Totals across type of cat breed do not sum to 100%, because a subset of respondents responded ‘don’t know’

Significantly more purebreds than domestic shorthair cats had one or more of the previously mentioned diseases (arthritis, oral disease, kidney disease, urinary disease, diabetes, or metabolic disease); and significantly more domestic shorthair cats had none of the studied health issues.

Also when owners were asked their opinion on their cat's general health there was a significant difference between purebreds and the two other breed types (χ^2 29.456, df.2; $p < 0.000$; $N = 368$). Thus, 62.3 % of owners of purebred cats responded that their cat was "generally healthy" compared to 85.9% and 90.6% of owners of mixed breed and domestic shorthaired cats, respectively.

An additional analysis shows that there is a clear difference in the prevalence with which owners of purebred (75%), mixed breed (35%), or domestic shorthaired cats (48%) have their cat vaccinated (Pearson's χ^2 20.88; df 2; $p < 0.000$). To check whether the differences found in observed disease occurrences between purebred and other cats were confounded by the difference in vaccination levels – as a proxy for concern about the health of the cats – we ran the analysis of the relationship between cat breed and disease (cf. Table 5) with additional explanatory variables inserted indicating whether the cat is vaccinated. However, the significant differences laid out in Table 5 were retained also after controlling for vaccination.

Table 6 displays the prevalence of health issues for non-neutered and neutered cats.

Table 6. Prevalence of overweight, disease or none of the diseases listed in neutered and non-neutered cats as reported by the owners in a representative study of Danish cats. $N = 376$						
	Prevalence (in %)			Adjusted OR (95% CI)*		
	Not neutered	Neutered	Total	Not neutered		
				OR	(95% CI)	
Overweight	7.3%	10.4%	10.3%	0.82	(0.23	2.86)
Disease**	17.1%	13.7%	13.8%	1.51	(0.57	4.01)
None of the diseases listed	75.6%	75.2%	75.2%	0.93	(0.41	2.10)
*Results from binary logistic regression ($N = 359$, where 'neutered' cat was set as reference value. Adjusted for age of the cat, how the cat is kept, and breed type						
** Diseases prompted for: arthritis, oral disease, kidney disease, urinary disease, diabetes, metabolic disease.						
Totals across type of cat breed do not sum to 100%, because a subset of respondents responded 'don't know'						

As can be seen, no significant differences were found here; neuter status did not affect the probability of health issues.

Discussion

Behavioural problems

The current study found a correlation between confinement and behavioural problems. This is important since there are strong voices arguing in favour of more confined cats, particularly in the US and in Australia. Still, as we saw, above 69.1% of cats in Denmark are allowed to roam freely

outside, and only 22.4% are confined indoors. Across the world the proportion of cats confined indoors varies. In the UK, like Denmark, the majority of the cats are allowed to roam outdoors (Rochlitz 2005). In the US, by contrast, it is estimated that 50–60% of cats are confined indoors (Patronek et al 1997; Bernstein 2007). In recent years some studies have investigated the benefits and other consequences of keeping cats confined as opposed to allowing roaming; both ways of living seem to be associated with risks and benefits.

The current study found confined cats to have a higher prevalence than free-roaming cats of nearly all of the studied behavioural problems. Confined cats also had a higher frequency of behavioural problems than garden cats in every respect assessed except aggressiveness towards the owner and house soiling. Our results are supported by other work, such as the study of Amat and others (2009), which identified five risk factors for the development of behavioural problems, with no outdoor access being one of them.

The reasons for higher levels of behavioural problems in confined cats are numerous and vary from one problem to another. In general, behavioural problems are likely to be due to increased stress, insufficient mental stimulation and lack of physical activity (Bain & Stelow 2014). Confinement reduces space and the variety and forms of potential activity available to most cats, and it locates cats in places designed around human convenience and comfort (Palmer & Sandøe 2014). The current study confirms that there is an association between the way the cat is kept and behavioural problems, and that being confined increases the likelihood of behavioural problems. Although confinement does provide some advantages, not all cats adapt to an indoor environment equally well (Jongman 2007).

We did not get information relating to the owners previous experience and knowledge about cat behaviour. One study showed a reduction in behavioural problems in kittens where owners were given advice on feline behaviour and on the appropriate education of their kitten by veterinary behaviourists during initial vaccination visits (Gazzano et al 2015). It is likely that behavioural problems can be reduced if owner awareness of feline behaviour and education is increased and veterinarians show a responsibility to support this awareness.

We also found a correlation between neuter status and behavioural problems. Neutered cats had significantly fewer behaviour problems than intact cats. Among specific behaviour problems, the amount of aggressive behaviour towards guests was significantly higher in intact cats than it was in neutered cats. This supports the widespread belief that neutering not only prevents reproduction but also curbs problems humans encounter with the behaviour of cats (Knol & Egberink-Alink 1989; Scarlett et al 2002; Fatjó et al 2006).

The link between behavioural problems and welfare is not simple. Some behavioural problems may just be a sign of the cat enjoying natural behaviour, such as scratching, and chewing, which is only a problem for the cat when it is deprived of appropriate environmental outlets for these behaviours (Herron & Buffington 2010). Other motivations or emotions causing behavioural problems, e.g.

increased anxiety, may be a sign that the cat actually has a welfare problem (Levine 2008). Some may be more difficult to interpret. Inappropriate elimination, for example, may both be a perfectly natural behaviour (e.g. marking behaviour or preference for an alternative substrate for the elimination) or a consequence of increased anxiety (Neilson 2004a). Furthermore, there may be an indirect link here with welfare in that cats with increased levels of welfare problems may have a more difficult time with their owner, making it more likely that the cat is relinquished or euthanized (Salman et al 2000; Kass et al 2001). Although it is true that behavioural problems are only to some extent direct signs of compromised cat welfare, they may indirectly affect welfare and longevity through owner reactions. Thus problem behaviour may need to be redirected to avoid owner frustrations (Jongman 2007).

The most frequently reported behavioural problem with cats in this study was the display of destructive behaviour. This behaviour is not a direct problem for the cat, but it will typically be a problem for the owner. Cats can damage furniture and other things in the home in several ways, but scratching probably accounts for most of the reported problems in our study. Scratching is a natural marking behaviour for the cat. It causes scent marks to be deposited from the inter-digital glands, leaving olfactory and visual signs, and it helps to maintain the cat's claws (Rochlitz 2007; Herron & Buffington 2010). Surfaces for scratching, such as scratching posts, should therefore be provided in attractive places in order to avoid unwanted scratching on furniture (Rochlitz 2005). Indoor cats may be short of suitable places to perform their scratching behaviour, and the display of destructive behaviour can therefore be a sign of boredom and lack of stimulation. Damage to household objects will often draw the owners' attention, and even if the consequence is scolding the cat will still learn to link the destructive behaviour and owner-attention. In time some cats will develop the habit of scratching as a means of attracting the owner's attention.

Significantly more confined and garden cats than free-roaming cats had house soiling problems and eliminated in places other than their litter box or outside. A cat's house soiling can be a cause of considerable frustration to the owner, and cats that show inappropriate elimination behaviour have a higher risk of relinquishment (Sung & Crowell-Davis 2006). The problems here can be categorized under three main categories: medical problems, marking, and toileting problems (Neilson 2004). They may have underlying motivations that owners find hard to understand. It is important to find the underlying motivation for the house soiling problem and to rule out or remedy any medical problem before wider adjustments are made. Urine marking is a natural behaviour that has a variety of communicative functions, including identification, and laying down emotional, temporal, and spatial information (Crowell-Davis et al 2004), but it is problematic when performed indoors. The motivation for indoor urine marking can be anxiety/stress, whereas toileting problems are often caused by medical issues, anxiety, aversions or preferences over litter boxes (Neilson 2004). Marking behaviour is primarily performed by fertile cats, but it can also occur in neutered cats.

Amat and others (2009) report an increased prevalence of behavioural problems in purebred cats. This finding is not supported by this study, since purebred cats did not have a significantly higher

frequency of any of the studied behavioural problems as compared with domestic shorthair and mixed breed cats.

It should be noted that other factors than those we study may have an effect on the prevalence of behavioural problems, e.g. the age of the cat when adopted, its provenance, how long the person has been the owner of the cat and how long the cat has lived in the same environment. It is a limitation of our study that we have not looked at these factors.

Health issues

Recent studies of confinement and free-roaming in cats have found that both ways of living are associated with risks and benefits. It has been found, for example, that indoor cats are at greater risk of developing such diseases as feline urologic syndrome, hyperthyroidism, dental disease (Buffington 2002; Buffington et al 2006; Rochlitz 2007), of suffering from diabetes mellitus (Rand et al 2004; Slingerland et al 2009), and of being ‘skinny fat’ (Bjornvad et al 2011). None of these diseases can be confirmed by this study, which found that confined cats did not have more health issues than garden cats or free-roaming cats. Part of the explanation for this may be that owners of confined animals spend more time stimulating their cats and give them extra resources indoors to compensate for missing behavioural opportunities.

Several studies have described a relation between confinement and obesity in cats (e.g. Sloth 1992; Robertson 1999). It is worth mentioning that not all studies confirm this association (Colliard et al 2009; Courcier et al 2010), and neither did this study. The proportion of overweight cats found in our study is lower than that found in other studies (Allan et al 2000; Lund et al 2005; Colliard et al 2009). This may be because the owners reported in this study were not given any tool to estimate their cats’ body condition and made subjective assessments. Also, studies show that owners underestimate their cats body condition and are unable to recognise that their pet is overweight (Colliard et al 2009; Courcier et al 2010). Furthermore, a recent Danish study of confined, adult neutered cats found that body condition score (BCS) underestimates the level of body fat percent (BF%) in these cats (Bjornvad et al 2011). The study suggests that confined cats have higher BF% as a result of low activity level, resulting in less muscle mass, and thus a higher BF%, as compared with control cats. The cats are skinny fat or suffer from sarcopenic obesity. The current study did not isolate sarcopenic obesity as a health issue, but it may be reasonable to expect that some of the confined cats in it were skinny fat.

Owing to a lack of genetic diversity, purebred cats are predisposed to various diseases (Sandøe et al 2016 Ch 7). Since only 62 of 415 (15.0%) of the cats in this survey were purebred cats, no breed specific analyses were performed. However, when purebred cats as a group were compared with domestic shorthair and mixed breed cats, the result was evident: a significantly higher prevalence of disease was reported by owners in the purebred group.

Selective breeding has led to purebred cat breeds that are fancied by many cat owners, but it has also compromised the health of purebred cats as a result of limited genetic diversity (Sandøe et al

2014). More than 20 diseases linked to inbreeding are seen within purebred cats, although not all breeds are affected (Lipinski et al 2008). According to our study, the two most popular breeds in Denmark are the Maine Coon and Norwegian Forest cat. Both breeds suffer from chronic gingivostomatitis (Kortegaard et al 2006; Fødevareministeriet 2013) and approximately 6.3% of Maine Coons suffer from hypertrophic cardiomyopathy (Godiksen et al 2011).

It may be suspected that the finding that purebred cats suffer from higher levels of disease compared to the other groups of cats is a reflection of the fact that the owners of these cats in addition to the higher monetary investment at the time of procurement also in general invest more and that they are more engaged in their cats' well-being than owners of non-purebred cats. Thus these owners could be more likely to notice and/or report health problems than owners of non-purebred cats; and the reported higher levels of disease could be a reflection of this rather than of a level of disease that is actually higher.

It is indeed the case that owners of purebred cats show a higher level of engagement, in that, for example, they more often have their cats vaccinated. However, even when we control for level of vaccination we still find a higher level of disease among purebred cats. Thus, the findings of this study leave little doubt that, viewed as a group, purebred cats suffer more from health issues than mixed breeds and domestic shorthairs. However, it should be stressed that it is compatible with the findings in our study that some breeds of purebred cat may not be particularly prone to diseases.

Animal welfare implications and conclusion

The aim of this study was to investigate how the ways in which cats are bred, live and are taken care of affects their welfare. The study focused on three factors, which were linked to risks of welfare problems. These are confinement, being intact and being purebred. The link with reduced welfare in purebred cats proceeds through increased disease load. By contrast, in the case of confinement and being intact there is a more indirect link to welfare via behaviour problems. Only some of the latter problems will directly affect welfare, but they all have an indirect effect via negative owner reactions.

For purebred cats there is clearly a need for more research into the disease problems linked to different breeds, and for greater focus on health in the breeding of cats. On both counts, research into purebred cats lags behind the canine sector by some distance. As regards the way cats are housed, there is a need for better information to be provided for prospective cat owners on the need to accommodate the behavioural needs of their cats if they are confined indoors; and clearly our findings are also of relevance to the often polarized indoor-outdoor debate. Finally our findings support the already widespread view that there are good reasons for neutering cats that are not kept for breeding purposes.

Acknowledgements

We would like to thank the Danish Knowledge Centre for Animal Welfare for economic support, Merete Fredholm for advice on breeding issues and Paul Robinson and Clare Palmer for checking the paper's English.

Conflicts of Interest

None

References

- Allan FJ, Pfeiffer DU, Jones BR, Esslemont DHB and Wiseman MS** 2000 A cross-sectional study of risk factors for obesity in cats in New Zealand. *Preventive veterinary medicine* 46: 183-196
- Amat M, de la Torre JLR, Fatjó J, Mariotti VM, Van Wijk S and Manteca X** 2009 Potential risk factors associated with feline behaviour problems. *Applied Animal Behaviour Science* 121: 134-139
- AVMA** 2012 *U.S. Pet Ownership & Demographics Sourcebook*. American Veterinary Medical Association. 2012 Edition. AVMA: Schaumburg, Illinois, United States
- Bain M and Stelow E** 2014 Feline aggression toward family members: a guide for practitioners. *Veterinary Clinics of North America: Small Animal Practice* 44: 581-597
- Bayer** 2013 News Release. Bayer-AAFP Study Reveals Half of America's 74 Million Cats Are Not Receiving Regular Veterinary Care. <http://www.catvets.com/public/PDFs/PressReleases-Media/AVMA-BVCUS3PressRelease.pdf>. Accessed on March 31. 2016
- Bernstein PL** 2007 The human-cat relationship. In: Rochlitz I (ed) *The welfare of cats* pp 47-89. Springer: Dordrecht, The Netherlands
- Bjornvad CR, Nielsen DH, Armstrong PJ, McEvoy F, Hoelmkjaer KM, Jensen KS, Pedersen GF and Kristensen AT** 2011 Evaluation of a nine-point body condition scoring system in physically inactive pet cats. *American Journal of Veterinary Research* 72: 433-437
- Buffington CAT** 2002 External and internal influences on disease risk in cats. *Journal of the American Veterinary Medical Association* 220: 994-1002
- Buffington CAT, Westropp JL, Chew DJ and Bolus RR** 2006 Risk factors associated with clinical signs of lower urinary tract disease in indoor-housed cats. *Journal of the American Veterinary Medical Association* 228: 722-725
- Colliard L, Paragon B, Lemuet B, Bénet J and Blanchard G** 2009 Prevalence and risk factors of obesity in an urban population of healthy cats. *Journal of Feline Medicine & Surgery* 11: 135-140
- Courcier EA, O'Higgins R, Mellor DJ and Yam PS** 2010 Prevalence and risk factors for feline obesity in a first opinion practice in Glasgow, Scotland. *Journal of Feline Medicine & Surgery* 12: 746-753
- Crowell-Davis SL, Curtis TM and Knowles RJ** 2004 Social organization in the cat: A modern understanding. *Journal of Feline Medicine and Surgery* 6: 19-28
- Danmarks Statistisk** 2000 Familiernes kæledyr. 40 pct. af familierne holder kæledyr. *Nyt fra Danmarks Statistik* Nr. 499

- Fatjó J, Ruiz-de-la-Torre JL and Manteca X** 2006 The epidemiology of behavioural problems in dogs and cats: A survey of veterinary practitioners. *Animal Welfare* 15: 179-185
- FEDIAF** 2014 *Facts & Figures 2014*. FEDIAF. European Pet Food Industry Federation: Brussels, Belgium
- Finkler H and Terkel J** 2010 Cortisol levels and aggression in neutered and intact free-roaming female cats living in urban social groups. *Physiology & Behavior* 99: 343-347
- Fødevareministeriet** 2013 *Rapport fra arbejdsgruppen vedrørende avl af dyr*. Fødevareministeriet: Copenhagen, Denmark
- Gazzano A, Bianchi L, Campa S, and Mariti C** 2015 The prevention of undesirable behaviors in cats: Effectiveness of veterinary behaviorists' advice given to kitten owners. *Journal of Veterinary Behavior: Clinical Applications and Research* 10: 535-542
- Godiksen MT, Granstrøm S, Koch J and Christiansen M** 2011 Hypertrophic cardiomyopathy in young Maine Coon cats caused by the p. A31P cMyBP-C mutation -the clinical significance of having the mutation. *Acta Veterinaria Scandinavica* 53: 1-11
- Hart BL and Barrett RE** 1973 Effects of castration on fighting, roaming, and urine spraying in adult male cats. *Journal of the American Veterinary Medical Association* 163: 290-292
- Heidenberger E** 1997 Housing conditions and behavioural problems of indoor cats as assessed by their owners. *Applied Animal Behaviour Science* 52: 345-364
- Herron ME and Buffington CAT** 2010 Environmental enrichment for indoor cats. *Compendium on Continuing Education for the Practising Veterinarian* 32
- Jongman EC** 2007 Adaptation of domestic cats to confinement. *Journal of Veterinary Behavior: Clinical Applications and Research* 6: 193-196
- Kass PH, New Jr JC, Scarlett JM and Salman MD** 2001 Understanding animal companion surplus in the United States: Relinquishment of nonadoptables to animal shelters for euthanasia. *Journal of Applied Animal Welfare Science* 4: 237-248
- Knol BW and Egberink-Alink ST** 1989 Treatment of problem behaviour in dogs and cats by castration and progestagen administration: A review. *Veterinary Quarterly* 11: 102-107
- Kortegaard HE, Venø T, Koch J and Eriksen T** 2006 Kronisk gingivo-stomatitis hos kat. *Dansk Veterinærtidsskrift* 89: 18-21
- Levine ED** 2008 Feline fear and anxiety. *Veterinary Clinics of North America: Small Animal Practice* 38: 1065-1079
- Lipinski MJ, Froenicke L, Baysac KC, Billings NC, Leutenegger CM, Levy AM, Longeri M, Niini T, Ozpinar H and Slater MR** 2008 The ascent of cat breeds: genetic evaluations of breeds and worldwide random-bred populations. *Genomics* 91: 12-21
- Lund EM, Armstrong P, Kirk CA and Klausner J** 2005 Prevalence and risk factors for obesity in adult cats from private US veterinary practices. *International Journal of Applied Research in Veterinary Medicine* 3: 88-96
- Neilson J** 2004 Thinking outside the box: Feline elimination. *Journal of Feline Medicine and Surgery* 6: 5-11
- Neilson J** 2004a Feline House Soiling: Elimination and Marking Behaviors. *Clinical Techniques in Small Animal Practice* 19: 216-224.
- O'Brien SJ and Johnson WE** 2007 The Evolution of CATS. *Scientific American* 297: 68-75

- Overley B, Shofer FS, Goldschmidt MH, Sherer D and Sorenmo KU** 2005 Association between ovariectomy and feline mammary carcinoma. *Journal of Veterinary Internal Medicine* 19: 560-563
- Palmer C and Sandøe P** 2014 For their own good: captive cats and routine confinement. In: Gruen L (ed) *The ethics of captivity* pp 135-155. Oxford University Press: Oxford and New York
- Patronek GJ, Beck AM and Glickman LT** 1997 Dynamics of dog and cat populations in a community. *Journal of the American Veterinary Medical Association* 210: 637-642
- Questionnaire 1** Questionnaire translated into English. Available at: http://animaletics.ku.dk/documents/Questionnaire_English.pdf
- Questionnaire 2** Original questionnaire in Danish. Available at: http://animaletics.ku.dk/documents/Questionnaire_Danish.pdf
- Rand JS, Fleeman LM, Farrow HA, Appleton DJ and Lederer R** 2004 Canine and feline diabetes mellitus: nature or nurture? *The Journal of nutrition* 134: 2072S-2080S
- Richmond R** 2013 Pet ownership in Australia. *Australian Veterinary Journal* 91
- Robertson I** 1999 The influence of diet and other factors on owner-perceived obesity in privately owned cats from metropolitan Perth, Western Australia. *Preventive veterinary medicine* 40: 75-85
- Rochlitz I** 2005 A review of the housing requirements of domestic cats (*Felis silvestris catus*) kept in the home. *Applied Animal Behaviour Science* 93: 97-109
- Rochlitz I** 2007 Housing and welfare. In: Rochlitz I (ed) *The welfare of cats* pp 177-203. Springer: Dordrecht, The Netherlands
- Salman MD, Hutchison J, Ruch-Gallie R, Kogan L, New Jr JC, Kass PH and Scarlett JM** 2000 Behavioral reasons for relinquishment of dogs and cats to 12 shelters. *Journal of Applied Animal Welfare Science* 3: 93-106
- Sandøe P, Corr S and Palmer C** 2016a *Companion animal ethics*. Universities Federation for Animal Welfare, Wiley Blackwell: Chichester, UK
- Sandøe P, Bjørnvad CR, Forkman B, Nørspang AP and Lund TB** 2016b Danskere og katte - resultater fra en repræsentativ spørgeskemaundersøgelse. *Dansk Veterinærtidsskrift*, issue 5 10 April 2016: 10-15.
- Sandøe P, Meyer I, Fredholm M and Proschowsky HF** 2014 Selektiv avl af racehunde og -katte: kan den ske på en velfærdsmæssigt forsvarlig måde? In: *Dyrevelværd i Danmark 2014* pp 36-49. Fødevarestyrelsen: Copenhagen, Denmark
- Scarlett JM, Salman MD, New JG and Kass PH** 2002 The role of veterinary practitioners in reducing dog and cat relinquishments and euthanasias. *Journal of the American Veterinary Medical Association* 22: 306-311
- Slingerland L, Fazilova V, Plantinga E, Kooistra H and Beynen A** 2009 Indoor confinement and physical inactivity rather than the proportion of dry food are risk factors in the development of feline type 2 diabetes mellitus. *The Veterinary Journal* 179: 247-253
- Sloth C** 1992 Practical management of obesity in dogs and cats. *Journal of Small Animal Practice* 33: 178-182
- Sung W and Crowell-Davis SL** 2006 Elimination behavior patterns of domestic cats (*Felis catus*) with and without elimination behavior problems. *American Journal of Veterinary Research* 67: 1500-1504