UNIVERSITY OF COPENHAGEN

Københavns Universitet

Assessment and Analysis of two Pinus kesiya provenance trials at Cashel, Zimbabwe (PV 144)

Trial No. 3 in the Pinus kesiya series

Hansen, Christian Pilegaard; Nyoka, Isaac; Mhongwe, Caleb H.

Publication date:
2003

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Hansen, C. P., Nyoka, I., & Mhongwe, C. H. (2003). *Assessment and Analysis of two Pinus kesiya provenance trials at Cashel, Zimbabwe (PV 144): Trial No. 3 in the Pinus kesiya series*. Danida Forest Seed Centre. Results and Documentation No. 18

Download date: 17. Nov. 2026

Assessment and analysis of a *Pinus kesiya* provenance trial at Cashel, Zimbabwe (PV 144)

Trial No. 3 in the *Pinus kesiya* series

by

Christian Pilegaard Hansen[2], Isaac Nyoka[1] and Caleb H. Mhongwe[1]

Forest Research Centre[1], Forestry Commission, Zimbabwe

Danida Forest Seed Centre[2], Denmark

Results and Documentation No. 18

Danida Forest Seed Centre

August 2003

Citation:
Hansen. C.P., I. Nyoka and C.H. Mhongwe. 2003. Assessment and analysis of a *Pinus kesiya* provenance trial at Cashel, Zimbabwe. Results and Documentation No. 18. Danida Forest Seed Centre, Humlebaek, Denmark.

Reproduction is allowed with citation

ISSN 0902-3224

Cover photo:
Stem defects as basal sweep, butt sweep and sinuosity are common for *P. kesiya*. Here Cashel trial site, Zimbabwe.
Photo: Christian Pilegaard Hansen

This publication can be requested from:

Danida Forest Seed Centre
Krogerupvej 21. DK-3050 Humlebaek, Denmark
Phone: +45-49190500
Fax: +45-49190258
Email: dfsc@sns.dk
Web Site: www.dfsc.dk

and/or be downloaded from the DFSC homepage:
www.dfsc.dk/publications/

Technical Editor: Melita Jørgensen

Print:
Toptryk A/S, Graasten

Results and documentations are publications of analyses of e.g. provenance trials, carried out between DFSC and other institutions.
DFSC publications are distributed free of charge

Danida Forest Seed Centre (DFSC) is a Danish non-profit institute which has been working with development and transfer of know-how in management of tree genetic resources since 1969. The development objective of DFSC is to contribute to improve the benefits of growing trees for the well-being of people in developing countries. DFSC's programme is financed by the Danish International Development Assistance (Danida).

Preface

This report presents the results of a joint assessment of the *Pinus kesiya* provenance trial established at Cashel, Zimbabwe. The trial was established by the Forest Research Centre, Forestry Commission, Zimbabwe in December 1992 as part of an international series of provenance trials of the species.

The joint FRC/DFSC field assessment took place in August 1998 by a team lead by Dr. Isaac Nyoka and Mr. Caleb H. Mhongwe of FRC and Christian Pilegaard Hansen of DFSC.

The Forest Research Centre is thanked for their kind assistance in the planning and arrangements for the field assessment.

Useful comments and assistance in the statistical analysis and interpretation of results were received from Anders Ræbild and Erik D. Kjaer, DFSC.

Contents

Preface i
Contents iii
Acronyms and abbreviations iv

1 Background 1

2 P. kesiya provenance trials in Zimbabwe 2
2.1 Earlier trials 2
2.2 Present trial - trial PV144 at Cashel 2

3 Field assessment and data management 4

4 Statistical analysis 5
4.1 Plot of raw data 6
4.2 Statistical model 6
4.3 Co-variates 6
4.4 Check of model assumptions 7
4.5 What to do when model assumptions are not fulfilled? 8
4.6 Fixed or random effects 9
4.7 Test of differences between provenances 9
4.8 Lsmeans (estimates from the fixed model) 10
4.9 Best Linear Unbiased Predictors (BLUPs - estimates from the random model) 10

5 Results of statistical analysis of individual traits 11
5.1 Survival 12
5.2 Height growth 14
5.3 Diameter growth 16
5.4 Mean volume of tree 18
5.5 Total volume per hectare 20
5.6 Stemform 22
5.7 Wood density 24
5.8 Foxtailing 26
5.9 Cone setting 28
5.10 Aphid attack 30
5.11 Frequency of leaning trees 32

6 Discussion and conclusions 34
6.1 Growth 34
6.2 Adaptation 34
6.3 Quality 34
6.4 Conclusion 35

7 References 36

Annexes

Annex 1 Trial design map
Annex 2 Trial description
Annex 3 Site description
Annex 4 Plot data set

Acronyms and abbreviations

BSO	Breeding seed orchard
CAMCORE	Central America and Mexico Coniferous Resources Co-operative, USA
CIEF	Centro de Investigaciones y Experiencias Forestales, Argentina
CSIRO	Commonwealth Scientific and Industrial Research Organisation, Australia
CSO	Clonal seed orchard
Danida	Danish International Development Assistance
DFSC	Danida Forest Seed Centre, Humlebæk, Denmark
EMBRAPA	Empresa Brasileira de Pesquisa Agropecuária, Petrolona, Pernambuca, Brazil
FAO	Food and Agriculture Organization of the United Nations, Rome, Italy
FRC	Forest Research Centre, Forestry Commision, Harare, Zimbabwe
ICFRE	Indian Council of Forestry Research and Education, Dehra Dun, India
IF	Instituto Florestal, São Paulo, Brazil
IPEF	Instituto de Pesquiasa e Estudos Florestais, Piracicaba, Brazil
OFI	Oxford Forestry Institute, United Kingdom
RCFTI	Research Centre for Forest Tree Improvement, Forest Science Institute, Vietnam
RFD	Royal Forest Department, Thailand

1. Background

The Cashel trial forms part of an international series of provenance trials of *P. kesiya*.

The objective of the international series is to explore and analyse the genetic variation in growth, quality and adaptive traits among provenances of *P. kesiya* throughout the range of the species. The results will facilitate an informed choice of seed source in planting programmes. Furthermore, the results will be useful when planning conservation activities of the species.

Below the background of the international series is briefly described.

Initial research on inter-population differences in *P. kesiya* was undertaken in Zambia in the 1950s. The test material included provenances from the Philippines, Vietnam and Assam. A comprehensive review of these studies is provided by Armitage and Burley (1980).

During 1969, FAO and the Forest Research Institute of Australia sponsored seed collections of 19 seed sources of *P. kesiya* from the Philippines (17 provenance collections and 2 commercial seedlots). The material was complemented by two Zambian land races (of Philippine and Vietnamese origin, respectively). These collections were used for provenance trials in a large number of countries for which the then Commonwealth Forestry Institute supplied advice and assisted in data processing and interpretation (Burley and Wood 1976). Results from individual trials were reviewed by Gibson and Barnes (1984). They concluded that neither provenance representation nor test site representation warranted an international evaluation. It was recommended that a more comprehensive exploration and analysis of the genetic variation of *P. kesiya* should be undertaken. Recommendations in this regard was also put forward by the Sixth Session of the FAO Panel of Experts on Forest Gene Resources (FAO 1988).

Exploration of provenance variation and collection of seed for field trials took place in the late 1980s in collaboration between national institutions in Brazil, Myanmar, China, Madagascar, Philippines, Thailand, Vietnam, Zambia, Zimbabwe, Oxford Forestry Institute (OFI) and Danida Forest Seed Centre (DFSC). In 1988, seed collections were complete and distribution to collaborating countries could begin (Barnes et al. 1989). Distribution of seed was co-ordinated by OFI and handled by DFSC. During 1989-93, seed of 42 provenances and land races from the above 9 countries were distributed to 20 institutions in 19 countries. Some seedlots were separated by family to allow testing of individual progenies.

A status on seed distribution and established field trials is found in DFSC (1996) and DFSC (1997). Some 30 trials have been established in 17 countries. Trials in Argentina, Brazil, Colombia, Indonesia, South Africa, Swaziland, Vietnam and Zimbabwe are reported with high survival and are in general in good conditions. Status of trials in Burundi, India, Rwanda and Sri Lanka is unknown, as no information has been received from these countries. Trials established in Fiji, Kenya, Nepal, the Philippines and Thailand have been abandoned because of fire damage, drought and browsing.

In a circular letter sent out by OFI and DFSC in 1996, host institutes were asked about their interest in undertaking a joint evaluation and were at the same time asked about the status of the trials (DFSC 1996). Positive responses in regard to the proposal of undertaking a joint assessment and analysis of trials have been received from all countries where existence of trials has been confirmed. The number, distribution and representation of provenances in these trials were considered sufficient to justify an assessment and analysis of the international series. Of special interest is the possibility of an in-depth analysis of provenance x site interactions, thanks to the representation of the same set of provenances at many trial sites.

A manual was elaborated during 1997-98 with a proposal for a set of characters to be assessed in all trials (DFSC 1998). Field assessment of trials commenced in April 1998 in Vietnam.

2. *P. kesiya* provenance trials in Zimbabwe

2.1 Earlier trials

P. kesiya was first introduced into Zimbabwe, probably from the Philippines in 1935 (Mullin *et al.* 1984).

A first provenance trial was established in 1968 at John Meikle Forest Research Station with four seedlots of Philippine, Vietnam, Assam and Burma origin (Mullin et al. 1984). In 1970/71 this was followed by trials at nine sites with seed lots of Philippine origin. These trials form part of the first international series of trials as described in Chapter 1. In the early 80's a third series of trials was established at four sites with provenances from Thailand plus four local controls.

Analysis results of the earlier trials are summarized by Mullin *et al.* (1984), and will be briefly discussed in Chapter 6 in connection with the results of the analysis of the present trial.

2.2 Present trial - trial PV144 at Cashel

FRC has established a provenance trial of *P. kesiya* in December 1992 at Cashel. The trial has the identification number PV144.

Information on trial design is in Annex 1 and establishment and site information is in Annex 2 and 3.

The trial forms part of the international series. The 21 seedlots represented in the trial are displayed in the below table together with summary information.

Most of the seedlots in the trial originate from natural forests (collections within the natural distribution range). Morarano (Madagascar) is an example of a land race. The origin of the source is most likely Vietnam. Unfortunately, no seed lot description is available. Two seed lots are from clonal seed orchards, in Zambia and Zimbabwe respectively. The seedlot from the Zimbabwean CSO (1776/88) originates from a single clone according to available information. Seed from all the ramets of this clone in the orchard has been bulked and thus represent half-sibs where the male percentage is (theoretically) a mixture of all the other clones in the seed orchard. The results for this seedlot should be interpreted with great care as it does not provide a fair representation of the seed orchard as such.

Seedlots 1624 (Baoshan), 1632 (Ceheng) and 1633 (Shangsi) are *P. yunnanensis*, whereas all others are *P. kesiya*.

Prov. no.	Acc. no. (DFSC)	Provenance	Country	Latitude	Longitude	Altitude m.a.s.l.	Rainfall mm/year	Temp. °C	No. Coll.
1	01447/84	Mt. Province	Philippines	17 15 N	120 30 E	2300	.	.	83
2	01448/84	Benquet	Philippines	16 35 N	120 30 E	1600	.	.	64
3	01515/8500	Dathien	Vietnam	11 58 N	108 27 E	1550	1769	17.9	45
4	01517/8500	Ho Tien	Vietnam	11 51 N	108 32 E	1500	1769	17.9	11
5	01519/8500	Lang Hanh	Vietnam	11 37 N	108 16 E	950	2059	21.5	22
6	01521/8500	Nong Krating	Thailand	18 05 N	98 35 E	1080	1332	22.2	17
7	01523/8500	Doi Inthanon	Thailand	18 32 N	98 35 E	1000	2084	20	30
8	01525/8500	Nam Now	Thailand	16 40 N	101 33 E	800	1316	.	35
9	01624/86	Baoshan (1)	China	24 51 N	99 12 E	1750	.	.	26
10	01632/86	Ceheng	China	24 24 N	105 34 E	800	.	.	25
11	01633/86	Shangsi	China	21 37 N	107 57 E	530	.	.	25
12	0163686	Jingdung (2)	China	24 28 N	100 51 E	1350	.	.	34
13	01637/86	Jinghong	China	22 25 N	101 10 E	1250	.	.	24
14	01638/86	Lancang	China	22 40 N	104 03 E	1620	2059	21.5	22
15	01772/8800	Zokhua	Burma	22 25 N	93 40 E	1600	2335	15.0	20
16	01773/8800	Aungban	Burma	20 41 N	96 37 E	1350	1303	20.5	61
17	01776/88	John Meikle CSO	Zimbabwe	18 43 S	32 51 E	1268	1725	17.9	1
18	01778/8800	Clonal Seed Orchard	Zambia	13 00 S	28 00 E	1300	.	.	10
19	01786/8800	Morarano	Madagascar	18 40	47 02 E	900	.	.	32
20	01572/8500	Coto Mines	Philippines	15 32 N	120 05 E	800	1000	25.1	9
21	01680/86	Dakha	Vietnam	14 48 N	107 56 E	1200	2684	21.6	40

Note: 'No. coll.' is the number of mother trees represented in the seed lot.

3. Field assessment and data management

The assessment in general followed the methodology described in DFSC (1998) which involves the characters:

1. Survival;
2. Health;
3. Social status (Kraft);
4. Height;
5. Diameter (DBH);
6. Straightness;
7. No. of whorls;
8. No. of branches in whorl;
9. Branch diameter;
10. Crown length;
11. No. of forks;
12. Position of first fork;
13. Foxtail;
14. Flowering and fruiting;
15. Wood density (Pilodyn);

For a detailed description of the assessment methodology, please refer to DFSC (1998).

The assessment of the Cashel trial did not include branch diameter and crown length, because the trial had been recently pruned, making assessment of these traits impossible. Under assessment of health, special attention was paid to assessment of woolly aphid attack and and 'leaning trees' (see later description)

The assessment was a full assessment, i.e. all trees in all plots were included.

Relative wood density was measured with a Pilodyn wood tester with pin diameter 2.0 mm.

Assessment data was recorded in the field on assessment sheets, see example in DFSC (1998). The data was immediately after the assessment entered to a lap-top computer in spreadsheet format. From the shread-sheet, data was later imported to a SAS-dataset for further analysis.

4. Statistical analysis

Overview of steps involved in the statistical analysis



The objectives of the statistical analysis are:

- to examine statistically significant differences between seedlots (provenances) in adaptability, growth and quality traits. A list of analysed traits is provided in Chapter 5;
- to conclude and recommend on the practical application of the results (species and provenance recommendations);
- to investigate patterns of genetic variation;
- to provide data for an overall analysis of the international series of provenance trials of *Pinus kesiya*, i.e. analysis across sites. This step will await completion of the analysis of individual trials.

Statistical analysis is done on plot values, e.g. plot averages or plot sums. Calculation of plot values is described in Annex 4.

The SAS analytical package has been used for the analysis (SAS, 1990).

The statistical analysis of each trait follows a sequence of steps. They are:

1. Plot of raw data;
2. Formulation of statistical model;
3. Test of co-variates;
4. Check of model assumptions;
5. When model assumptions are not fulfilled: (a) transformation of data; (b) deletion of outliers; (c) weight statement;
6. Test of differences between species and provenances;
7. Calculation of lsmeans (estimated from a model with fixed effects);
8. Calculation of BLUPs (estimated from a model with random effects).

The statistical analysis is illustrated in the above figure and the steps are further described in the below text.

Generally speaking, two different approaches are applied in the statistical analysis: a fixed effect approach and a random effect approach. The fixed effects approach is concerned with the genetic entries (seedlots) actually in the trial, whereas the random effects approach concerns what would happen if the experiment was to be repeated. Following the fixed effect approach, the estimates are calculated as least square means (lsmeans), whereas the random effect approach gives the best linear unbiased predictors (BLUPs). See further explanation below.

4.1 Plot of raw data

The main purpose of the plots is to indicate the scale of the variable along with a first impression of the variation within the trial. Often the visual inspection of the data reveals clear differences between the provenances, or gives hints regarding proper transformations of the data. Obvious outliers (extreme values) may also be identified already at this stage.

The most useful single plot is probably a plot of the variable against the provenances, marking the values with values identifying the blocks. However, other plots may also be relevant, e.g. plotting the values by block or by the distance along the axis of the trial.

4.2 Statistical model

The test of differences between seedlots (provenances) is based on the model:

$$X_{jk} = \mu + provenance_j + block_k + \varepsilon_{jk}$$

where X_{jk} is the value of the trait in question (e.g. height) in plot *jk*,
μ is the grand mean,
$provenance_j$ is the effect of seedlot number *j* and is assumed to be either a fixed or a random effect, according to which approach is used (see later),
$block_k$ is the effect of block *k* in the trial, assumed to be a random effect, and
ε_{jk} is the residual of plot *jk* and is assumed to follow a normal distribution $N(0, \sigma_e^2)$.

Please note that the controls (seedlots not *Pinus kesiya*/*P.yunnanensis*) are considered (analysed) together with these sources not considering that they actually are sources of different species.

4.3 Co-variates

In order to reduce the residual variation in trials with heterogeneous trial conditions (e.g. variation in soil, elevation, slope and exposure within trial), a number of co-variates are included in the model. As a standard routine the following four co-variates are tested:

plotx: Horizontal position of plot within trial (see map of trial);
ploty: Vertical position of plot within trial (see map of trial);

To catch non-linear patterns of site variation vertically and horizontally, plotx2 and ploty2 are applied:

plotx2: $plotx2=(plotx - mean(plotx))^2$
ploty2: $ploty2=(ploty - mean(ploty))^2$

In addition to the above four co-variates, additional co-variates are considered in some of the trials:

level: Level of plot in relation to a reference plot within the trial (0);
plotxy $plotxy=plotx \times ploty$

In testing the effect of co-variates, we start with a model with all co-variates included. Co-variates

that are not significant are removed successively by removing the least significant co-variate and running the model again until all remaining co-variates in the model are significant ($P<0.10$).

4.4 Check of model assumptions

The statistical model rests on a number of standard assumptions. Key assumptions are (see e.g. Box *et al.* 1978):

(i) that the residuals are independent;
(ii) that the residuals follow a normal distribution;
(iii) that there is variance homogeneity in effects included in the model.

The model assumptions are checked graphically by producing a number of plots:

1. Student's residuals versus predicted values;
2. Cooks distance versus predicted values;
3. Student's residuals versus provenance;
4. Frequency chart of residuals;
5. Student's residuals versus block;
6. Student's residuals versus plotx;
7. Student's residuals versus ploty;
8. Student's residuals versus level (if level is among the considered co-variates).

The residuals represent variation that can not be accounted for by the model. For each observation, the model calculates a predicted value, taking into account the effects of the model (provenance, block and co-variates). The residual variation is then the difference between the observed value and the predicted value.

Student's residuals (also called 'standardised residuals') are calculated as the residual divided by its standard error. If the assumption of normal distributed residuals is valid, the Student's residuals have the property of a normal distribution with mean 0 and variance 1, meaning that 95% of the values should lie within ± 1.96. In cases of trials with imbalance, the Student's residuals correct for imprecision due to low sample numbers, and in models with co-variates they compensate for large deviations at extreme values.

The Student's residual e_t for observation *ij* is given by

$$e_t = \frac{e_{ij}}{s_{ij}} = \frac{X_{ij} - P_{i\bullet} - B_{\bullet j}}{s_{ij}}$$

where e_{ij} is the residual, X_{ij} is the value for observation *ij*, $P_{i\bullet}$ is the effect of provenance *i*, $B_{\bullet j}$ is the effect of block *j*, and s_{ij} is the standard deviation (standard error) of observation *ij*.

Cooks distance gives a measure of the influence of a single observation (plot) on the model, and gives an indication of possible 'outliers' (see below) (Afifi & Clark 1996). A high value indicates an observation with a large influence on the outcome of the model.

In the following, a description of the check of the model assumptions is given.

Independence

The assumption of independence means that the residual of one observation is not dependent on the residual of another. This assumption is typically violated when using pseudo-replicates, e.g. when doing more observations on the same experimental unit and treating them as different experimental units. Another example is when two or more plots of the same provenance within the same block are treated as independent observations. In such cases, an average of the values should be used as the block value for the provenance in question.

The graphical check of the residuals does not reveal possible problems with observations dependent upon each other, and there is no easy method to ensure that the condition of independence is fulfilled. Proper design and planning of the experiment result and application of a correct statistical model is the best insurance to obtain independent observations.

The assumption of independence may also be violated if there is a time- or site-dependency in the data. To check for such dependency, residuals are plotted against the horizontal and vertical axis of the trial (plotx and ploty) and where applicable, also the level of plot, to investigate any systematic environmental variation. Usually there is none, as the co-variates (plotx and ploty) account for this.

Normality

The assumption of normality may be checked in various ways, graphically as well as by statistical tests. In this analysis, we use the frequency chart of residuals as a graphical check. In the frequency chart, the frequencies should be more or less bell-shaped with no large tails at the ends. A formal statistical test, the Shapiro-Wilk statistic, is given in the SAS-procedure UNIVARIATE with the option NORMAL (SAS 1988a). This procedure also offers different kinds of plots of the residuals. However, since the test is usually considered to be conservative, rejecting only severe deviations from normality, the test results should be considered with caution (Brockhoff, pers. comm.).

When the number of observations is low, it becomes increasingly difficult to check the assumption of normality. Even though the frequency chart may show a rather odd and irregular distribution, this need not be a sign of non-normality. At small sample sizes it is not unlikely that odd distributions may result from random variation, and unless the test for normality demonstrates that the assumption is violated, there is no need to reject the model. On the other hand, when the number of samples is very large, the test for normality may

become rejected even though the frequency chart of residuals appears to be normal. This is because the power of the test increases with the number of observations, and even small deviations from normality may result in rejection of the hypothesis of normal distributed residuals. In such cases it should be considered whether the frequency chart indicate that the assumption is fulfilled, or the deviations are so large that transformation of data (see later) is required.

Deviations from the assumption of normality may also be interpreted as a distribution with a large number of outliers (see later).

Variance inhomogeneity

Variance inhomogeneity occurs when different experimental units (blocks and provenances) have different variance. A typical example is when the residuals of some provenances appear very clustered in the diagram of Student's residuals versus provenances, whereas the residuals of other provenances are spread out, often with values of Student's residuals exceeding ±2. This may result from a simple scale effect (larger provenances have larger variance), in which case the plots of Student's residuals and Cook's distance versus predicted values appear funnel-shaped. It may also be related to the provenance itself (some provenances are more variable than others). In this case, the variance inhomogeneity will be displayed in the plot of student's residuals versus provenance.

Outliers

Outliers are extreme observations that do not follow the trends of the remaining data. Such observations may have a large influence on the estimates and statistical tests of the model and should therefore be considered carefully.

Outliers are detected by inspection of the plots of Student's residuals and of Cook's distance. Observations that have values of Student's residuals exceeding ±2.5 (rule of thumb), and observations with large values of Cook's distance, are possible outliers and should be investigated further. Outliers may be due to errors in the recording or typing of data, or due to mislabelling of the seedlots in the nursery or in the field. Poor survival in the plot, leaving only a few trees to use in the calculation of plot means is another source of outliers. However, it also happens that the outliers are due to some unexplained variation, perhaps in soil conditions or other environmental variation. Finally it should be mentioned that a large number of outliers might indicate that the distribution of residuals is not normal, and hence that a transformation of data is required (see later).

When outliers occur as a result of errors, the dataset should of course be corrected, which will solve the problem. It is less obvious what to do in the cases where there are no easy explanations. Outliers should only be excluded if it can be justified, i.e. an explanation can be given. In a few cases, however, explanations were not found, and observations were excluded alone on basis of the extreme nature of the value. Great care is required in the decision to exclude plot values, as it will have great importance for the result of the analysis, especially with few blocks (replications). Running the analysis again without the outlier(s) gives an indication of the sensitivity of the analysis in regard to the outliers, and assist in deciding whether to keep or delete the extreme observation(s).

In the interpretation of the statistical analysis in this report, it is always mentioned if one or more extreme values have been considered as outliers and omitted from the analysis, and on what grounds.

4.5 What to do when model assumptions are not fulfilled?

In many cases one or more of the model assumptions are not fulfilled. In the below, procedures for correction are described.

Independence

Apart from making sure that the statistical design and the model is correct there is not much to do about dependence between observations. If some clear variation can be observed in the residuals, other co-variates than the ones mentioned above could be considered.

Deviations from normality

Usually deviations from normality are handled by transformation of data. Snedecor & Cochran (1980) and Afifi & Clark (1996) provide guidance on data transformations:

1. Counts (of rare events) often follow a Poisson distribution and are transformed with the square root.
2. Variables having a binomial character (e.g. dead or alive) summarised in a proportion (e.g. living trees in a plot) may be transformed with the arc sine transformation.
3. If the standard deviation varies directly with the mean, a logarithmic transformation may stabilise the variance.

There are theoretical reasons for choosing the above transformations (Snedecor & Cochran 1980), but it follows from Afifi & Clark (1996) that the range of transformations may be seen as a continuum and that various other transformations are available.

None of the variables included in the present assessment have the character of a Poisson distribution, but the square root transformation has nevertheless in some cases been applied.

In many cases the analysis of survival data results in skewed distributions of the residuals, with tails at either the lower or upper end (many trees dead or many trees alive). In such cases an arc sine transformation of data will often prove useful.

The arc sine transformation is given by

$$\arcsin(proportion) = \sin^{-1}\sqrt{proportion}$$

where proportion is a figure between zero and one (e.g. the surviving fraction of trees). An important property of the transformation is that the variance near zero or one is stretched out, thus facilitating the analysis of variance (Snedecor & Cochran 1980).

For many growth variables, the variance increases directly with tree size, and the proper transformation is thus the logarithm. In most cases, the natural logarithm (ln) has been applied to achieve a normal distribution of residuals.

Variance inhomogeneity

In the cases where the variance varies with the size of the variable, a transformation of data is the proper way to solve the problem (see above). However, in some cases the provenances simply have different variances irrespective of size, and it is necessary to weight the observations with weights proportional to the reciprocals of the error variances to ensure variance homogeneity (SAS 1988b, cf. Afifi & Clark 1996). There may also be cases where different blocks have different variances, but this has not been observed in the present trial(s).

Weighting occurs in the following sequence: An ordinary analysis of variance of the variable is performed. The residuals from this analysis are grouped according to provenance, and the variance of the residuals for each provenance is calculated. The inverse of these variances is then used as weights in an analysis of variance. When calculating the sums of squares in the model, the weights are multiplied with the squared value of the deviance of each observation from the predicted value (SAS 1988b). This has the effect that provenances with small variances have a larger influence on the model than provenances with larger variances. In other words, the more stable the provenance, the more it counts in the analysis. Provenances with large uncertainty on the other hand have less influence.

4.6 Fixed or random effects

A special problem relates to the choice between considering the effects in the statistical model as fixed or random. Statistically speaking, fixed effects are considered as *parameters* (unknown constants). Random effects are considered *stochastic variables* with an expected value of zero and a variance (Skovgård 1994). Fixed effects are used when the individual groups (seedlots) are of interest. Models with random effects are used when interest is in the size of the variation between the groups (described by the variance), including groups that are not represented in the trial. In analysis of random effects it is important that the groups are representative of a larger population of groups, and they should preferably be chosen by randomisation (Skovgård 1994). In the words of Stonecypher (1992), 'fixed models address estimating and testing to infer the existence of true differences among means, whereas the random models address estimating and testing to infer the existence of components of variance'.

To choose between a fixed or a random effects model is a choice with no simple answer. Stonecypher (1992) has formulated the following two questions to facilitate a choice:

1. 'Are the conclusion confined to the things actually studied; to the immediate sources of these things; or extended to apply to more general population?'

2. 'In complete repetitions of the experiment would the same things be studied again; would new samples be drawn from the same sources; or would new samples be drawn from the general population?'

When the objective is to estimate components of variance, the effects should be considered as random. If the objective is to estimate differences among means, the effects should be considered as fixed. In some cases fixed and random effects may be combined in the model (mixed models). This is the case when special designs are applied, such as split-plot or nested designs.

In our model with only provenances and blocks, it is necessary to choose between considering the provenance effect as random or as fixed. If the aim is to compare the specific provenances and the actual production on the site, it is natural to consider the provenance effect as fixed. If, on the other hand, (i) the provenances are assumed to be representatives of a population of provenances; (ii) the aim is to expand the conclusions to this population; (iii) to estimate the production and (iv) should the experiment be repeated, then the provenance effect should be considered as random.

The results of the statistical tests are irrespective of whether the provenance effect is considered a fixed or a random effect. However, there are major differences in the estimates resulting from the two approaches (see below). Since it may be argued that both the fixed and the random approaches are relevant in this analysis, both sets of estimates have been calculated.

4.7 Test of differences between provenances

In our statistical model, differences between provenances for a given trait are tested by an F-test comparing the mean square of provenances with the residual mean square. The hypothesis tested is that there is no difference between the provenances. If the F-test is significant, we reject the hypothesis and conclude that there are significant differences between the provenances.

The testing is done using the GLM procedure in SAS (SAS 1990). Since the testing of random vari-

ables may involve combinations of different mean squares (Skovgård 1994), an approximation called Satterthwaites approximation is used in the calculation of degrees of freedom (SAS 1988).

4.8 Lsmeans (estimates from the fixed model)

In the fixed model approach, the estimates for the provenances are calculated as the least square means (lsmeans). The main difference between raw means and lsmeans is that lsmeans account for missing values and imbalanced designs. Thus, in completely balanced designs there are no differences between lsmeans and the raw means. It follows that the lsmeans are the best estimates for the given provenance in the trial.

The confidence intervals and limits are calculated from the formula (Skovgård 1994)

$$X \pm t_{1-\alpha/2,(a-1)(b-1)} \sqrt{s^2 / b}$$

where X is the least square mean, α is the confidence level (in this case 0.05, giving a 95% confidence interval), a is the number of provenances and b is the number of replicates (blocks) of each provenance. s^2 is the mean square of the error (MS_e). The confidence limits are calculated directly by SAS in the LSMEANS statement with the CL option.

Since the estimates are calculated individually, different provenances may have different lengths of the confidence intervals (due to different variances). In the cases where the data have been weighted, the confidence intervals are adjusted according to the variance of each provenance and thus are of different lengths.

Special problems arise when the data has been transformed. If the least square means and the confidence limits are calculated on basis of the transformed values, the back-transformed estimates will be geometric means rather than arithmetic means. This implies that the estimates become biased towards lower values, and compared to the real values actually are under-estimates. If on the other hand the estimates are calculated using raw data, the lsmeans will be arithmetic means (comparable to the real mean values), but the confidence limits are based on a faulty distribution and will be wrong. In this analysis we have calculated estimates on the transformed values in order to get a fair representation of the differences between provenances. Usually the figures are presented together with a raw mean to circumvent the problem with under-estimation.

4.9 Best Linear Unbiased Predictors (BLUPs - estimates from the random model)

In the random approach, the provenance effects are seen as coming from a normal distribution with an expected value and a variance. This is in opposition to the fixed effect approach, where the provenance effects are seen as constants. Estimating provenance effects in random models is more complicated than in fixed models, because the observed variation between provenances is contemplated as a mixture between true provenance effects and random error variation (*cf.* White & Hodge 1989). The variation between the provenances is therefore always larger than the true 'genetic' variation, except in cases where the error variation is negligible.

In order to predict the effect of a given provenance, it is necessary to correct the estimates for the part of the variance that is due to random error variation. This is done by calculating the best linear unbiased predictors (BLUPs, White & Hodge 1989). The calculation of BLUPs is cumbersome and only feasible with a suitable software package. In this case, the SAS procedure MIXED has been used. It follows from the above that the predicted values for the provenances fall within a smaller range than the least square means. Often the results are presented as deviations from the mean value to allow for easier comparison between different experiments. The deviations are expressed either in real values (m, cm^2 etc.) or in % deviation from the mean value. Here deviations are presented as % deviations from the mean values.

The problems with transformed values are the same as described for the least square means above. A further complication arises when calculating the deviations from the mean value in percent. If the mean value is calculated on the base of transformed values, and the deviations are calculated on the basis of this back-transformed mean, the deviations from the mean will not sum to zero. In this analysis, we have therefore chosen to base the deviations from the mean value on values calculated after transformation.

The BLUPs are presented with *t*-type confidence intervals. However, these should be interpreted with caution since it is probably wrong to assume that the underlying distribution of the estimates is normal because of the limited sample size (Littell et al. 1996). Confidence intervals are presented to give an impression of the variation between the provenances and should not be interpreted with respect to differences between provenances.

5. Results of statistical analysis of individual traits

The below table displays the traits selected for analysis, grouped into growth traits, adaptive traits and quality traits. For a full description of the traits and their calculation, please refer to Annex 4.

Group	**Trait description**	**Analysed trait**
Growth	Height growth	Height of tree with diameter corresponding to mean basal area (H_G)
	Diameter growth	Diameter of tree corresponding to mean basal area (D_G)
	Mean volume of tree	Average of volumes of trees in plot
	Standing volume per hectare	Volume per hectare
Adaptation	Survival	Survival percentage
	Cone setting	Average score of old, open cones
	Foxtailing	Foxtailing percentage
	Woolly aphid	Frequency of attack (%)
	'Leaning trees'	Frequency of 'leaning trees' (%)
Quality	Stemform	Stemform score (1-9)
	Relative wood density (Pilodyn)	Diameter adjusted pilodyn readings

5.1 Survival

OVERVIEW OF ANALYSIS	
Co-variates	None
Data transformation required	Yes. Arc sin transformation
Weight statement	Yes
Outliers	None

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	9.97	9.85 (***)	0.0001
Block	5	0.12	0.11	0.99
Error	100	1.01		

Data has been transformed before analysis using an arc-sin transformation. This has been done to better fulfil the model assumptions as described in Chapter 4.

The survival percentage is high – 100 per cent or close to 100 per cent – for most provenances. The only provenance with a somewhat lower survival (92%) is the Aungban provenance from Myanmar. If this source is left out of the analysis there are not statistically significant differences between provenances.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Survival (%)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Survival. Best linear un-biased predictors (BLUPs)



5.2 Height growth

OVERVIEW OF ANALYSIS	
Co-variates	PLOTY
Data transformation required	No
Weight statement	Yes
Outliers	Yes. Shangsi (Block 1)

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	89.95	88.68(***)	<0.0001
Block	5	7.17	7.07	<0.0001
PLOTY	1	7.45	7.34	0.0079
Error	98	1.01		

The height of the Shangsi provenance in Block 1 is very different from the performance of the provenance in the other 5 blocks (much taller in Block 1). An investigation of the assessment data did not reveal any mistakes or data errors which could explain this. It was decided to exclude the plot from the analysis because of its extreme nature.

There are highly significant differences among the provenances in regard to height growth.

The Philippine sources are ranked at the top, with Coto Mines at the very top, and Benquet and Mt. Province ranked three and four. Also the Vietnamese sources are ranked high in height growth with Da Thien ranked second and Da Kha ranked 5, Ho Thien 7, and Lang Hanh 9. The offsprings from the Zambia CSO, and John Meikle CSO (Zimbabwe) are also doing relatively well, ranked 6 and 8 respectitively. There are only small, and statistically not significant differences among many of the top-ranking sources.

The poorest height performers are the *P. yunnanensis* sources Shangsi and Ceheng, the Burmese sources (Zok Hua and Aungban). Thai and Chinese *P. kesiya* sources are intermediate.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Height (m)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Height. Best linear un-biased predictors (BLUPs)



5.3 Diameter growth

OVERVIEW OF ANALYSIS	
Co-variates	PLOTY
Data transformation required	No
Weight statement	Yes
Outliers	Yes. Shangsi (Block 1)

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	29.00	27.78(***)	<0.0001
Block	5	12.23	11.72	<0.0001
PLOTY	1	61.21	58.64	<0.0001
Error	98	1.04		

As was the case with the analysis of height growth, Shangsi (Block 1) is identified as 'outlier', and has been excluded from the analysis.

There are highly significant differences between seed lots in regard to diameter growth.

The three *P. yunnanensis* sources, Shangsi, Baoshan and Ceheng, have the poorest diameter growth, as was the case with height growth.

The ranking at the top, however, is quite different to what was seen for height growth. Most notably, the Philippine sources, which were top-ranking in height growth, are ranked low in diameter growth, especially the Coto Mines and Mt. Province. Dathien (Vietnam), which was also ranked high in height growth, has the highest diameter. It is followed by Lancang (China), Doi Inthanon (Thailand) and Jinghong (China), which were intermediate in height growth. The three other Vietnamese sources, Lang Hanh, Ho Thien, Da Kha, and the Zambia CSO are also ranked at the top as was the case with height growth. The Burmese sources Zok Hua and Aungban are among the poorest sources.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Diameter (cm)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Diameter. Best linear un-biased predictors (BLUPs)



5.4 Mean volume of tree

OVERVIEW OF ANALYSIS	
Co-variates	PLOTY
Data transformation required	No
Weight statement	Yes
Outliers	Yes. Shangsi (Block 1)

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	76.51	73.86(***)	<0.0001
Block	5	17.61	17.00	<0.0001
PLOTY	1	91.11	87.95	<0.0001
Error	98	1.04		

Mean volume of tree is calculated as the average of the volumes of individual trees. As both height and diameter are included in the volume formula (see Annex 4), the trait thus illustrates a combined effect of height and diameter.

As for the other growth traits, the Shangsi sources in Block 1 has been omitted from the analysis as it differs very much from the other plots (much taller and bigger trees in this plot than in the other plots). No explanation has been found and the plot is thus omitted from the analysis on the basis of the extremeness of the observation. The omission has little practical importance, as the Shangsi provenance as seen in the previous analysis, is among the poorest growth performers.

The *P. yunnanensis* sources, Baoshan, Shangsi and Ceheng, have very low volumes and are ranked at the very bottom. Of the *P. kesiya* sources, the Burmese sources Aungban and Zok Hua have the lowest volumes. Slightly better is the sources Jindung (China), Nong Krating and Nam Now (Thailand) and Mt. Province (Philippines).

At the top we find the Vietnamese sources, Dathien, Da Kha, Ho Thien and Lang Hanh. All 4 tested Vietnamese seed sources are among the best five performers, together with the Zambia source (ranked second). Coto Mines of the Philippines is also among the best due to a good height growth. There are only small, and not statistically significant differences, among the top-ranking provenances.

Intermediate growth performers are the Chinese sources Jinghong and Lancang, Benquet (Philippines), Doi Inthanon (Thailand), Morarano (Madagascar), and John Meikle CSO (Zimbabwe).

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Volume of mean tree (m3)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Volume of mean tree. Best linear un-biased predictors (BLUPs)



5.5 Total volume per hectare

OVERVIEW OF ANALYSIS	
Co-variates	PLOTY
Data transformation required	No
Weight statement	Yes
Outliers	No

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	50.72	48.25(***)	<0.0001
Block	5	10.71	10.19	<0.0001
PLOTY	1	29.61	28.17	<0.0001
Error	99	1.05		

The analysis of total volume production can be seen as an analysis summarising survival, height growth and diameter growth in one analysis as all three traits are included in the calculation of volume per hectare.

The analysis is performed on the full dataset, i.e. no outliers. Shangsi (Block 1) that has been excluded in the other growth trait analysis is included in the present analysis as the value – even though close to being an outlier – is less extreme than in the other analysis.

The ranking of the provenances is almost identical to what was seen for volume of mean tree; the Vietnamese sources – Dathien, Da Kha, Lang Hanh and Ho Thien are top-ranked together with the seed lot from Zambia. They are followed by Coto Mines (Philippines), Lancang and Jinghong (China), Benquet (Philippines), Doi Inthanon (Thailand) and Morarano (Madagascar). There are no statistical significant differences among the top-ranking provenances.

In the bottom end of the ranking, we find – again – the sources of *P. yunnanensis* and the *P. kesiya* sources Aungban and Zok Hua from Myanmar.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Volume per hectare (m3/ha)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Volume. Best linear un-biased predictors (BLUPs)



5.6 Stemform

OVERVIEW OF ANALYSIS	
Co-variates	PLOTY
Data transformation required	No
Weight statement	Yes
Outliers	None

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	25.86	24.55 (***)	<0.0001
Block	5	8.01	7.60	<0.0001
PLOTY	1	15.52	14.74	0.0002
Error	99	1.05		

There are highly significant differences between provenances in regard to stemform.

The Zambian seed orchard offspring has the highest stemform score. Lang Hanh and Dathien from Vietnam are next. Then we have John Meikle seed orchard (Zimbabwe) and the Morarano landrace from Madagascar. The seed orchard sources, and especially the Zambia CSO, are ranked high. Improvement of stemform has been a key issue in the selection of plus trees for the orchard, and the results illustrate that this selection has been effective.

The best 'un-improved' sources are the Vietnamese sources Lang Hanh, Dathien and Ho Thien and the Morarano landrace from Madagascar. The latter source probably originates also from Vietnam. The Da Kha source from Vietnam is ranked considerably lower than the other Vietnamese sources.

The Philippine sources are of intermediate stemform together with Nam Now (Thailand), Zok Hua (Myanmar) and Shangsi, Jinghong and Lancang from China.

Poorest stemform has Aungban (Burma), Jingdung (China) and Doi Inthanon (Thailand).

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Stemform (1-9 score)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Stemform. Best linear un-biased predictors (BLUPs)



5.7 Wood density (Pilodyn)

OVERVIEW OF ANALYSIS	
Co-variates	PLOTX, PLOTX2 and PLOTY2
Data transformation required	No
Weight statement	Yes
Outliers	Da Kha (Block 1)

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	14.46	13.79 (***)	<0.0001
Block	5	22.53	21.48	<0.0001
PLOTX	1	11.02	10.50	0.0017
PLOTX2	1	4.16	3.97	0.0493
PLOTY2	1	12.50	11.92	0.0008
Error	92	1.05		

Diameter adjusted pilodyn readings (ref. Annex 4) are used in the analysis.

The Da Kha source in Block 1 is omitted from the analysis because of the extreme nature of this plot (very high pilodyn reading) compared to other plots. An investigation of assessment data did not reveal anything which could explain the extreme nature of the plot value.

The analysis of variance reveals highly significant differences between provenances.

The Philippine sources, and especially Coto Mines, have the lowest pilodyn readings, corresponding to highest wood densities. The Philippine sources have a relatively slow diameter growth compared to many other sources, and the high wood density may be related to this. Also the Zok Hua source from Myanmar (but not the Aungban source) and John Meikle CSO and Zambia CSO sources have high densities.

The Vietnamese sources have lower wood densities than the top-ranking sources, especially when compared to the Philippine sources Coto Mines and Benquet.

The *P. yunnanensis* sources – Ceheng, Baoshan and Shangsi, have the lowest wood densities. Low densities have also Jingdung and Lancang from China.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Pilodyn



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Pilodyn. Best linear un-biased predictors (BLUPs)



5.8 Foxtailing

OVERVIEW OF ANALYSIS	
Co-variates	None
Data transformation required	No
Weight statement	Yes
Outliers	None

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	9.30	9.67 (***)	<0.0001
Block	5	7.22	7.51	<0.0001
Error	100	0.96		

Analysis of frequency of foxtails gives an indication of the adaptability of the source to site (lower frequency means in general a better adaptation). Moreover, and of great practical importance, foxtails is related to the quality of the tree. Trees with foxtails often have ramicorns, i.e. thick branches from a lower whorl growing (co-evolving) vertically along with the main stem. They are able to develop as there are no whorls on the foxtail to suppress vertical growth of the branches. Trees with coarse branching are thus often a result of foxtails. This means lower quality, especially if timber is the final product. In addition, harvesting operations become more difficult. Apart from leading to coarse branching, foxtails will often result in stem breaks as the wood of the foxtail is of low density.

There are significant differences among seedlots in regard to foxtail percentage.

The Vietnamese sources (Hothien, Dathien, Da Kha and Lang Hanh) have the highest frequencies of foxtails. The Philippine sources (Benquet, Mt. Province and Coto Mines) also have high levels of foxtails, as has the Morarano landrace from Madagascar.

The lowest levels of foxtails are found among the Burmese sources (Aungban and Zok Hua), the *P. yunnanensis* sources (Ceheng, Shangsi and Baoshan), and the Thai sources (Nam Now, Nong Krating and Doi Inthanon). Chinese sources (Jinghong, Jingdung and Lancang) are intermediate, as are the seedlots from John Meikle and Zambia CSOs.

It is important to note the large size of the confidence intervals of many of the seed lots, indicating that there are only significant differences between top and low ranked provenances, whereas many of the differences within these groups are not statistically significant.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Foxtail percentage (%)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Foxtailing. Best linear un-biased predictors (BLUPs)



5.9 Cone setting

OVERVIEW OF ANALYSIS	
Co-variates	None
Data transformation required	No
Weight statement	Yes
Outliers	None

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	22.88	22.81 (***)	<0.0001
Block	5	5.98	5.96	<0.0001
Error	100	1.00		

Data on flowering and fruiting provides an indication of the adaptation of the sources to site. Abundant flowering and fruiting is generally interpreted as a sign of good adaptation to the site, and vice versa.

As the trial is still young, only about 6 years of age at the time of assessment, flowering and fruiting may only have just commenced and the tested seedlots do not express their full potential for flowering and fruiting. Consequently, the analysis should be interpreted with care.

The analysed trait is average score of old, open cones. Assessments have been done using a 1-9 scale where score 1 refers to no flowers or cones. More details in DFSC (1998).

The analysis of variance shows highly significant differences among provenances in regard to cone setting.

The most abundant cone setter is the *P. yunnanensis* source of Shangsi, closely followed by the *P. kesiya* sources Zok Hua and Aungban from Myanmar. Also the Chinese sources Jindung, Lancang and Jinghong set more cones than the other sources.

The John Meikle CSO and Coto Mines (Philippines) sources have the lowest cone setting among the tested. Low cone setting is also expressed by the Thai sources (Nam Now, Nong Krating and Doi Inthanon), the Vietnamese sources Ho Thien and Lang Hanh and the two other tested sources from the Philippines (Benquet and Mt. Province).

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Old, open cones (1-9 score)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Old, open cones. Best linear un-biased predictors (BLUPs)



5.10 Aphid attack

OVERVIEW OF ANALYSIS	
Co-variates	PLOTY
Data transformation required	Yes. Arc sin transformation
Weight statement	Yes
Outliers	Aungban, John Meikle, Shangsi and Zokhua in Block 1

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	3.82	3.69 (***)	<0.0001
Block	5	18.63	18.00	<0.0001
PLOTY	1	4.89	4.72	0.0323
Error	99	1.03		

Attack of the pine woolly aphid, *Pineus pini* (L), has been assessed on all trees. The analyzed trait is the proportion of affected trees to total number of trees in plot.

An arc-sin transformation of data was required to obtain a normal distribution of residuals. Four 'outliers' were identified, all in Block 1. An examination of data did not give any explanation or reveal any errors in data recording or calculations. It was, however, decided to omit them from the analysis because of the extreme nature of the plots.

The analysis of variance reveals statistical significant differences between the tested sources.

The two seed lots originating from seed orchards (John Meikle CSO, Zimbabwe and Zambia CSO) have the lowest frequencies of aphid attack. Resistance to aphid attack has probably been one of the traits considered when selecting for the orchards, and the analysis results suggest that this selection has been efficient.

In the opposite end of the ranking, we find Zok Hua (Myanmar) having the highest frequency of aphid attack. Also the Philippine sources, especially Coto Mines and Mt. Province show high levels of aphid attack.

There are no clear patterns of provenance variation, as we have seen for some of the other analyzed traits. As an example, the Vietnamese provenances are spread throughout the ranking, not grouped together as we have seen elsewhere. It is also important to stress that the confidence intervals are extremely large. This is maybe indicating that assessment of woolly aphid has not been consistent. Results should therefore be interpreted with care.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Aphid attack (%)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Aphid attack. Best linear un-biased predictors (BLUPs)



5.11 Frequency of 'leaning trees'

OVERVIEW OF ANALYSIS	
Co-variates	PLOTX
Data transformation required	No
Weight statement	Yes
Outliers	No

RESULTS FROM ANALYSIS OF VARIANCE				
Effect	**DF**	**MS**	**F-value**	**P-value**
Provenance	20	3.39	3.28 (***)	<0.0001
Block	5	4.41	4.27	<0.0015
PLOTX	1	7.37	7.13	0.0089
Error	99	1.03		

A prominent feature of the Cashel trial is a large number of trees with a very peculiar growth habit. They bend to the ground with the crown almost touching the ground. Other trees have collapsed completely and are fallen over. This growth habit has not been observed in any of the other trials and the reason behind it is not known. One explanation could be that the trees are growing in over-luxuriant conditions (former agricultural land) and that this may have caused the strange growth habit of many trees. It has been experienced elsewhere that too favourable site conditions can lead to tree collapse (flop) (Armitage and Burley 1980).

As part of the field assessment, all 'leaning' trees were recorded, and the proportion of leaning trees to total number of trees in plot has been analyzed to check for differences among provenances.

The analysis of variance reveals significant differences among provenances.

Ceheng (*P. yunnanensis*) has the lowest percentage of leaning trees, followed by Jinghong (China), Baoshan (*P. yunnanensis*) and Zok Hua (Myanmar). Doi Inthanon (Thailand), John Meikle CSO (Zimbabwe) and Da Kha (Vietnam) have the highest percentages.

There is a tendency of slow growing sources having fewer problems with 'leaning' trees, e.g. the *P. yunnanensis* sources, and fast growing sources having higher frequencies. The picture, however, is not clear, and there are many exceptions. The confidence intervals are also quite wide, and there are not statistically significant differences between many of the provenances, only between the very top and low-ranked sources.

Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Percentage of leaning trees (%)



Pinus kesiya provenance trial, Cashel, Zimbabwe

International Series of Pinus kesiya provenance trials, Trial No. 3

Established December 1992. Assessed August 1998

Leaning trees (%). Best linear un-biased predictors (BLUPs)



6. Discussion and conclusions

6.1 Growth

Growth traits should be given key importance in the interpretation of trial results, as wood for pulp has high priority.

The analysis of growth traits (height, diameter, volume of mean tree, total volume per hectare) all indicate that the Vietnamese sources are performing well, especially the Dathien source. The results thus indicate that the Vietnamese sources represent an interesting genetic resource, and that introductions and further exploration of the provenance variation of material from within that region could be an interesting route to follow in the future.

Also the Zambia CSO source has a high volume production at the same level as the Vietnamese sources, and is a good choice of seed source for immediate supply.

Philippine sources (esp. Coto Mines) are also performing relatively well in regard to growth. They have a fast height growth (Coto Mines is the top-ranking provenance in height growth), but a slower diameter growth, making the volume production lower than would be expected based on the height alone.

The Zimbabwean seedlot from John Meikle CSO is not among the top-ranking sources, which is somewhat surprising. As earlier mentioned, the seedlot originates from only one clone in the orchard, and this might explain the somewhat disappointing performance. A 'normal' seedlot, sampling all clones in the orchard, would probably perform better.

The analysis indicates very clearly that *P. yunnanensis* has no potential on sites similar to the trial site. It is performing very poorly both in height and diameter. The same with the Myanmar sources Aungban and Zokhua.

The growth analysis results are in accordance with the results of Mullin *et al.* (1984). They reported, based on results from trial 31 at John Meikle Experimental Station, the Vietnamese source from Dalat and the Philippine source from Benquet to perform considerably better in height growth than Burmese and Assam sources. The Dalat source is close to the seed collection sites of the Hothien, Dathien and Lang Hanh sources in the present series.

Production potential, based on the best performing sources and based on growth in the first 6 years, is 15-17 m^3/ha/year.

6.2 Adaptation

Adaptive traits include survival percentage, foxtail frequency, flowering and health (aphid attack and 'leaning' trees).

The survival percentage is high for all provenances, and there are no statistically significant differences among the top-ranking provenances.

Vietnamese and Philippine sources, esp. the Vietnamese sources, have more foxtails than Thai, Myanmar and *P. yunnanensis* sources. Foxtails may also influence the quality of the wood as discussed in section 5.8. The Zambian and Zimbabwean sources are intermediate in regard to foxtails.

The results on cone-setting should not be given too much importance because of the early stage of assessment. The results at this early development stage indicate that the Myanmar and *P. yunnanensis* sources have higher cone-setting than the other sources, and that the Vietnamese, Philippine and Thai sources set fewer cones.

In regard to woolly aphid, John Meikle CSO and Zambia CSO express the lowest levels of infestion. Apart from that, the picture is rather confusing with no clear pattern of geographical variation.

The analysis of 'leaning trees' should be interpreted with care. First of all, the reason for the strange growth habit expressed by many trees in the trial is not known, and it has not been experienced in any other trial in the series. There are significant differences among the tested sources, but no clear pattern of provenance variation. Results may indicate that sources with faster growth have more problems than slower growing sources, but the picture is not evident, and there are exceptions.

6.3 Quality

Analyzed quality parameters are stemform, and wood density (pilodyn). Because of pruning, branching habit (branch diameter) could unfortunately not be assessed and analyzed.

The seedlots from the clonal seed orchards in Zambia and Zimbabwe have the best stemform, suggesting that selection has been effective. The Vietnamese sources also have good stemform, as have the landrace from Madagascar. The Philippine sources have a slightly poorer form, but are still above average.

In regard to wood density, the Philippine sources, esp. Coto Mines, are outstanding. They compare favorable to the seed orchard offspring from Zambia and Zimbabwe. The Vietnamese sources have lower basic density than the Philippine and seed orchard material, and this – together with the relatively high levels of foxtails – may be a constraint for a wider use of these sources.

There is good accordance between the results (straightness and wood density) in the present trial and the results of Mullin *et al.* (1984). They also found the Vietnamese source to have better

straightness than other sources, and with a lower basic density that the Philippine source. They further found a significant negative correlation between basic density and altitude of provenance collection site. The results of the present trial confirms this negative correlation for the Philippine sources (only three sources, though), whereas no such correlation is present in the material as such.

6.4 Conclusion

The trial is only 6 years of age, and it may thus be premature to draw very specific and detailed conclusions based on the present assessment. It is thus recommended that the trial is maintained, and re-assessed later on.

Based on the results of the present assessment, the Vietnamese sources show promise. They are fast-growing and they have a – for *P. kesiya* – good stemform. They are less favorable in regard to wood density, especially compared to the Philippine sources, and they set many foxtails. It is concluded, however, that Vietnamese material could be an important genetic resource with a potential in Zimbabwe. The feasibility and possibilities to include this material in seed production and further breeding activities of the species should be further investigated.

In regard to practical recommendations on seed sources for immediate use in plantation establishment, John Meikle CSO would be an obvious choice. The trial results do unfortunately not provide reliable information on the performance of this source, as the seedlot only represent one clone in the orchard. Other seed sources may be available within Zimbabwe, e.g. the BSOs established with material from Zambia and Madagascar in the early 90's, which may soon come into production.

If imports are to be considered, the results from the present trials indicate that material from the CSO in Zambia would be a good choice. It is a good volume producer, and it has an excellent stemform.

7. References

Afifi, A.A. & V. Clark. 1996. Computer-aided multivariate analysis. Chapman & Hall, London (3rd Edition)

Armitage, F.B. and J. Burley. 1980. *Pinus kesiya*. Tropical Forestry Papers No. 9. Department of Forestry, Commonwealth Forestry Institute, University of Oxford, Oxford, U.K. 199p

Barnes, R.D., H. Keiding and G.L. Gibson. 1989. Progress on seed collection for the second stage international provenance trials of *Pinus kesiya*. Forest Genetic Resources Information No. 17.

Box, G.E.P., W.G. Hunter and J.S. Hunter. 1978. Statistics for experimenters. John Wiley and Sons Inc., New York.

Burley, J. and P.J. Wood 1976 Manual on Species and Provenance Research with particular reference to the tropics. Tropical Forestry Papers No. 10. Department of Forestry, Commonwealth Forestry Institute, University of Oxford, Oxford, U.K. 226 p.

DFSC 1996. Proposal for a common evaluation of an international series of *Pinus kesiya* trials. DFSC Working Paper. 8 p.

DFSC 1997. Status on *Pinus kesiya* international provenance trials - Proposed Work Program 1996-99. DFSC Working Paper. Danida Forest Seed Centre, Humlebæk, Denmark. 10 p.

DFSC 1998. International Series of Provenance Trials of *Pinus kesiya*. Field Assessment manual. Danida Forest Seed Centre, Humlebæk, Denmark. 19 p.

FAO 1988. Report of the Sixth Session of the FAO Panel of Experts on Forest Gene Resources. Food and Agriculture Organization of the United Nations. Rome, Italy. 79 p.

Gibson, G.L. and Barnes, R.D. 1984 Status of the international provenance trials of *Pinus kesiya* and alternatives for future development. In: Barnes, R.D. and Gibson, G.L. (Eds.): Provenance and genetic improvement strategies in tropical forest trees. Proceedings of the IUFRO Conference, Mutare, Zimbabwe, April 1984. Commonwealth Forest Institute and Forest Research Centre, Harare.

Keiding, H., H. Wellendorf and E.B. Lauridsen.1986. Evaluation of an international series of teak provenance trials. Danida Forest Seed Centre, Humlebæk, Denmark. 81 p.

Littell, R.C., Milliken, G.A., Stroup, W.W., Wolfinger, R.D. 1996. SAS® System for mixed models. SAS Institute Inc., Cary, NC, 633 pp.

Mullin, L.J., D.R. Quaile and W.R. Mills. 1984. *Pinus kesiya* provenance trials in Zimbabwe. In 'Provenance and genetic improvement strategies in tropical forest trees'. Mutare, Zimbabwe, April 1984. Eds: Barnes, R.D. and Gibson, G.L. Commonwealth Forestry Institute, Oxford and Forest Research Institute, Harare

SAS 1988a. SAS® Procedures Guide, Release 6.03 Edition. SAS Institute Inc., Cary, NC, 441 pp.

SAS 1988b. SAS/STAT® Users Guide, Release 6.03 Edition. SAS Institute Inc., Cary, NC, 1928 pp.

SAS 1991. SAS® System for Statistical Graphics, First Edition. SAS Institute Inc., Cary, NC, 697 pp.

Skovgård, I.M. 1994. Statistisk Forsøgsplanlægning (*in Danish*). Jordbrugsforlaget, Copenhagen, Denmark, 318 pp.

Skovgaard, I.M. & P. Brockhoff. 1998. Multivariate analysis and variance components. Lecture notes, Dept. of Mathematics and Physics, The Royal Veterinary and Agricultural University, Copenhagen, 41 pp.

Snedecor, G.W. and W.G. Cochran. 1980. Statistical methods. Iowa State University Press, 7th ed., 507 pp.

Stonecypher, R.W. 1992. Computational methods. *In*: Fins, L., Friedman, S.T. & Brotschol, J.V. (eds.): Handbook of quantitative forest genetics. Kluwer Academic Publishers, Dordrecht, pp. 195-228.

White, T.L. G.R. Hodge. 1989. Predicting breeding values with applications in forest tree improvement. Kluwer Academic Publishers, Dordrecht, 367 pp.

Personal communication

Brockhoff, P. Lecturer in statistics. The Royal Veterinary and Agricultural University, Copenhagen.

Annexes

Annex 1. Trial design map

															Plot y
			Block 2												
		21	20	19	18		Block 3			Block 4			Block 5		11
		14	15	16	17	1	2	3	1	2	3	1	2	3	10
		13	12	11	10	6	5	4	6	5	4	6	5	4	9
			7	8	9	7	8	9	7	8	9	7	8	9	8
			6	5	4	12	11	10	13	14	15	13	14	15	7
			1	2	3	13	14	15	13	14	15	13	14	15	6
			19	20	21	18	17	16	18	17	16	18	17	16	5
			18	17	16	19	20	21	19	20	21	19	20	21	4
	Block 1		13	14	15	15	16	17	18	19	20	21			3
12	11	10	9	8	7	14	13	12	11	10	9	8			2
1	2	3	4	5	6	1	2	3	4	5	6	7			1
									Block 6						
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	

----------Plot x----------

Layout of trial at Cashel. Plot positions are defined by two coordinates (PLOTX, PLOTY).

Randomization

Plot no.	Block 1	Block 2	Block 3	Block 4	Block 5	Block 6
1	01680/86	01632/86	01523/85	01517/85	01773/88	01777/88
2	01517/85	01572/85	01447/84	01772/88	01517/85	01572/85
3	01515/85	01680/86	01777/88	01680/86	01572/85	01517/85
4	01638/86	01773/88	01773/88	01636/86	01515/85	01778/88
5	01773/88	01448/84	01680/86	01572/85	01637/86	01525/85
6	01523/85	01523/85	01448/84	01521/85	01777/88	01773/88
7	01777/88	01624/86	01638/86	01637/86	01448/84	01633/86
8	01624/86	01633/86	01624/86	01523/85	01624/86	01786/88
9	01772/88	01778/88	01519/85	01519/85	01638/86	01680/86
10	01633/86	01521/85	01786/88	01773/88	01633/86	01519/85
11	01572/85	01636/86	01778/88	01624/86	01632/86	01632/86
12	01637/86	01772/88	01772/88	01448/84	01636/86	01638/86
13	01519/85	01637/86	01525/85	01786/88	01778/88	01772/88
14	01632/86	01519/85	01633/86	01515/85	01786/88	01447/84
15	01786/88	01515/85	01517/85	01525/85	01525/85	01636/86
16	01525/85	01447/84	01515/85	01777/88	01519/85	01515/85
17	01448/84	01638/86	01521/85	01633/86	01680/86	01521/85
18	01778/88	01777/88	01636/86	01638/86	01523/85	01523/85
19	01636/86	01517/85	01572/85	01632/86	01447/84	01637/86
20	01521/85	01525/85	01632/86	01447/84	01521/85	01624/86
21	01447/84	01786/88	01637/86	01778/88	01772/88	01448/84

Acc. no. (DFSC)	Provenance	Country	Latitude	Longitude	Altitude m.a.s.l.	Rainfall mm/year	Temp. °C	No. Coll.
01447/84	Mt. Province	Philippines	17 15 N	120 30 E	2300	.	.	83
01448/84	Benquet	Philippines	16 35 N	120 30 E	1600	.	.	64
01515/8500	Dathien	Vietnam	11 58 N	108 27 E	1550	1769	17.9	45
01517/8500	Ho Thien	Vietnam	11 51 N	108 32 E	1500	1769	17.9	11
01519/8500	Lang Hanh	Vietnam	11 37 N	108 16 E	950	2059	21.5	22
01521/8500	Nong Krating	Thailand	18 05 N	98 35 E	1080	1332	22.2	17
01523/8500	Doi Inthanon	Thailand	18 32 N	98 35 E	1000	2084	20	30
01525/8500	Nam Now	Thailand	16 40 N	101 33 E	800	1316	.	35
01624/86	Baoshan (1)	China	24 51 N	99 12 E	1750	.	.	26
01632/86	Ceheng	China	24 24 N	105 34 E	800	.	.	25
01633/86	Shangsi	China	21 37 N	107 57 E	530	.	.	25
0163686	Jingdung (2)	China	24 28 N	100 51 E	1350	.	.	34
01637/86	Jinghong	China	22 25 N	101 10 E	1250	.	.	24
01638/86	Lancang	China	22 40 N	104 03 E	1620	2059	21.5	22
01772/8800	Zokhua	Burma	22 25 N	93 40 E	1600	2335	15.0	20
01773/8800	Aungban	Burma	20 41 N	96 37 E	1350	1303	20.5	61
01776/88	John Meikle CSO	Zimbabwe	18 43 S	32 51 E	1268	1725	17.9	1
01778/8800	Clonal Seed Orchard	Zambia	13 00 S	28 00 E	1300	.	.	10
01786//8800	Morараho	Madagascar	18 40	47 02 E	900	.	.	32
01572/8500	Coto Mines	Philippines	15 32 N	120 05 E	800	1000	25.1	9
01680/86	Dakha	Vietnam	14 48 N	107 56 E	1200	2684	21.6	40

Trees are assessed starting in lower left corner of plot, going up and down the rows, ending in upper right corner.

Annex 2. Trial description Cashel, Zimbabwe

TRIAL ESTABLISHMENT AND MANAGEMENT

Year and month of establishment: *December 1992*

Area (ha): *1.0 ha*

Initial spacing (m x m): *2.7 m x 2.7 m*

Soil preparation (time, method/intensity):
Planting method(age of seedlings, type):

Beating up (time, %):

Irrigation (time, amount): *None*

Fertilization (time, type, amount):

Weeding (time, intensity):

Thinning (time, intensity): *None, but pruned in 1998*

Firelines:

TRIAL DESIGN

Statistical design: *Randomized complete block design*

No. of replications (blocks): *6 replications*

No. of treatments (provenances): *21 provenances (see list in Annex 1)*

Plot size (No. of trees in plot): *16 (4 x 4)*

Demarcation (blocks, plots): *Labels, poles in plot corners. Good demarcation*

PROTECTION STATUS

Status (describe any disturbances/damages):

Guarding (permanent, regular, none):

Annex 3. Trial site description

Site description – Cashel, Zimbabwe

LOCATION

Province:
District:
Latitude (degrees and minutes):
Longitude (degrees and minutes):
Altitude (m above sea level): *1450 m above sea level*

Managing office/institution: *Forest Research Centre, Forestry Commission*
Owner: *Forest Research Centre, Forestry Commission*
User(s): *Forest Research Centre, Forestry Commission*

Distance to nearest office responsible for management of the trial (km):
Distance to nearest villages/towns (km):
Number of inhabitants in the nearest villages/towns:

Type of area (e.g. research station, managed forest, etc.): *Research Station*

CLIMATE

Nearest weather station:
Name of the station: *Cashel*
Latitude (degrees and minutes): *19°33'N*
Longitude (degrees and minutes): *32°47'E*
Altitude (m a.s.l.): *1190*

Climatic data[1]	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.	Year
Rainfall (mm)	193.7	165.0	108.7	39.2	17.7	10.4	7.0	12.4	10.8	36.3	94.5	195.0	890.7
Temp. mean (°C)	.	.	.	.	.	.	.	.	.	.	.	.	.
Temp. mean max.[2] (°C)	.	.	.	.	.	.	.	.	.	.	.	.	.
Temp. mean min.[3] (°C)	.	.	.	.	.	.	.	.	.	.	.	.	.
Evapotranspiration[4] (mm)	.	.	.	.	.	.	.	.	.	.	.	.	.

[1] Period of observations Unknown [2] Average of daily maximum temperatures
[3] Average of daily minimum temperatures [4] Potential evapotranspiration (ETP) - Penman's formula

Rainy season:

Number/type of seasons: [x] one [ ] two [ ] even/irregular

Period(s): *Nov. – April* (specify months)

Length of rainy season:
No. of intermediate days: *60* (pre- and posthumid period of the growing season)
No. of wet days: *180* (growing season)

Number of dry months per year (< 50 mm rain/month): *4 months*
Frost (number of days/year):
Prevailing winds (direction, period, speed): *NE (W)*

TOPOGRAPHY (slope) of trial site

☐ Flat/gentle (0-8%) ☐ Intermediate (9-30%) ☐ Steep (>30%)

GENERAL SOIL DESCRIPTION

Soil texture		Soil depth		Soil drainage/ n		Gravel content, topsoil	
1. Light/sandy		1. Shallow (< 50 cm)		1. Well drained	X	1. None (< 15 %)	
2. Medium/loamy	X	2. Deep (50-100 cm)		2. Seasonal		2. Gravelly (15-35 %)	
3. Heavy/clayey		3. Very deep (> 100 cm)	X	3. Permanent		3. Stony (> 35 %)	

Organic matter content		Reaction (pH)		Soil salinity		Groundwater	
1. Poor (< 2 % DM)	X	1. Acid (pH < 6.5)	X	1. None	X	1. Shallow (< 50 cm)	
2. Medium (2-5 % DM)		2. Neutral (6.5-7.5)		2. Moderate		2. Deep (50 - 150 cm)	
3. Rich (> 5 %)		3. Alkaline (pH> 7.5)		3. High		3. Very deep (>150 cm)	X

Specify soil unit, soil association and phases (subdivisions of soil units) according to the Soil map of the world (FAO-Unesco 1971-1979), if known:

VEGETATION

Natural (original) vegetation type:.

Dominant natural (original) genera/species:

Land use history: *Agricultural land (peas)* → *Abandoned for six years* → *Widdringtonia hodflora* → *trial site*

RESULTS OF SOIL SAMPLE

Sample 1: Block 2

Description	Unit	Result
Depth of sample	m	1.2
Clay (<2 µm)	%	52.5
Fine silt (2-20 µm)	%	16.6
Coarse silt (20-63 µm)	%	5.9
Fine sand (63-125 µm)	%	4.9
Fine medium sand (125-250 µm)	%	6.1
Coarse medium sand (250-500 µm)	%	5.7
Coarse sand (500-2000 µm)	%	8.5
Org. Mat.	%	1.2
Lime	%	0.0
pH-H_2O	-	4.91
P		- 1

Results noted as - 1: Amount not detectable

Sample 2: Block 6

Description	Unit	Result
Depth of sample	m	1.2
Clay (<2 µm)	%	24.1
Fine silt (2-20 µm)	%	22.1
Coarse silt (20-63 µm)	%	16.3
Fine sand (63-125 µm)	%	23.7
Fine medium sand (125-250 µm)	%	10.1
Coarse medium sand (250-500 µm)	%	2.4
Coarse sand (500-2000 µm)	%	1.2
Org. Mat.	%	0.6
Lime	%	0.0
pH-H_2O	-	5.34
P		- 1

Results noted as - 1: Amount not detectable

Annex 4. Plot data set

Aggregated data set at plot level. This annex describes the variables in the plot dataset, and displays the data. The plot dataset has been prepared from the individual tree dataset and holds the following parameters.

PARAMETER NAME	EXPLANATION
BLOCK	Block No.
SEEDLOT	Seedlot No.
SITE	Name of site
DATEEST	Establishment data of trial (MM/YY)
DATEASS	Date of assessment (MM/YY)
PLOTX	Co-variate
PLOTY	Co-variate
PLOTX2	Co-variate
PLOTY2	Co-variate
PLOTXY	Co-variate
PLOTYX	Co-variate
DBH1	Average of diameters (arithmetic mean)
HEIGHT	Average height (arithmetic mean)
SURV	Survival percentage (%)
GMEAN	Mean basal area per tree (m2)
GSUM	Total basal area in plot (m2)
MEANPILO	Mean pilodyn for plot
WHORLS	Average number of whorls
BRANCH	Average number of branches in whorl
DIABRA	Diameter of largest branch (cm)
FORK	Frequency of trees with one or more forks (%)
FOXTAIL	Frequency of foxtails in plot (%)
NO_TOTAL	Number of trees in plot
NO_G	Number of trees in basal area calculation
NOHEIGHT	Number of trees in height-diameter regression
KRAFT1.. KRAFT5	Frequency of individual Kraft scores 1 to 5 (%)
KRAFT	Average Kraft index
STEM	Average stemform
STEM1..STEM9	Frequency of individual stemform scores 1 to 9 (%)
BRARATIO	Branch ratio (not calculated in this trial)
FO_INDEX	Forking index (not calculated in this trial)
INTNODE	Average distance between whorls (m)
CROWNPCT	Crown percentage (not calculated in this trial)
FO_POS	Average position of lower fork (m above ground)
TOP	Percentage of trees with broken top
VTREE	Average volume of tree in plot
MF	Average score of male flowers
FF	Average score of female flowers
CL	Average score of conelets
CC	Average score of closed cones
OC	Average score of old, open cones
LOTH	Percentage of 'leaning trees'
APHID	Percentage of trees with woolly aphid
DG	Diameter corresponding to mean basal area at breast height (cm)
HG	Height for tree with diameter corresponding to mean basal area (m)
PILOCORR	Mean pilodyn reading adjusted for diameter effect
VHA	Volume per ha

DG- Diameter corresponding to mean basal area at breast height (1.3 m)

DG is calculated using the following formula:

$$DG = \left(\sqrt{\frac{\sum_{i=1}^{n} \pi \times \left(Di/2\right)^2}{n}} \over \pi \right) \times 2$$

where
D_i is the diameter at breast height of tree No. i (in cm);
n is the total number of trees in plot.

GHA - Basal area per hectare

Basal area in m^2 per hectare is calculated using the formula:

$$GHA = \frac{\sum_{i=1}^{n} \pi \times \left(Di/2 \times 100\right)^2}{n \times sp^2} \times 10000$$

where
D_i is the diameter at breast height of tree no. i (in cm);
n is the total number of trees in plot; and
sp is the spacing in m;

HG- Height of tree with diameter corresponding to mean basal area

A linear regression per plot is prepared using the model:

$$Height = \beta + \alpha \times \ln(DBH1)$$

where
DBH1 is the diameter of stem (first stem if more than one stem) in cm;
α is the slope of the regression line;
β is the intercept with y-axis;

For each plot, α and β are estimated using PROC REG (SAS 1990).

HG for the plot is then calculated using the linear regression estimates (α and β) and plot DG (as previously calculated).

PILOCORRR - Correction of pilodyn readings

Tree diameter (ring width) influences the pilodyn reading, i.e. trees with larger diameter (rings) will normally have larger pilodyn readings (deeper penetration of the pilodyn pin) than trees with smaller diameter, all other factors equal (ceteris paribus).

In order to adjust pilodyn readings for the variation in diameter, a correction factor has been introduced. By doing so, we are reducing the variance due to differences in individual tree size, and the provenances are in the subsequent analysis compared assuming that they have the same average tree size. In other words, we compare the level of the trait rather than the actual value.

The adjustment has been made using the GLM procedure in SAS (SAS 1990). The following model is applied:

. . .

PROC GLM;
CLASS plot;
MODEL pilo = plot DBH1;
LSMEANS plot OUT=A;

. . .

Forking index (not used in the analysis of the present trial)

The forking index is calculated using the formula:

$$FO_INDEX = \frac{\sum_{i=1}^{n} FORK_i / FO_POS_i}{n}$$

where
$_{FORKi}$ is the number of forks observed on tree i
FO_POS_i is position above ground of first fork (in m) on tree i; and
n is the total number of trees with forking data in the plot

Branching index (not used in the analysis of the present trial)

The branching index is calculated using the formula:

$$BR_INDEX = \frac{\sum_{i=1}^{n} BRANCH_i \times DIABRA_i}{n \times 10}$$

where
$BRANCH_i$ is the number of branches on tree i;
$DIABRA_i$ is the branch diameter on tree i; and
n is the total number of trees with branching data in the plot.

INTNODE - Average distance between whorls

The INTNODE parameter is calculated using the formula:

$$INTNODE = \frac{\sum_{i=1}^{n} (HEIGHT_i - 0.5) / WHORLS_i}{n}$$

where

$HEIGHT_i$ is the height of tree i
$WHORLS_i$ is the number of whorls on tree i; and
n is the total number of trees with observations on whorls in the plot.

Plot Data Set - Cashel

BLOCK	SEED	PROVNAME	SITE	DATEEST	DATEASS	PLOTX	PLOTY	PLOTX2	PLOTY2	PLOTXY	PLOTYX	_TYPE_	_FREQ_	DBH1	HEIGHT	SUREV	GMEAN	GSUM	MEANPILO	WHORLS	BRANCH	DIABRA	FORK	FOXTAIL	NO_TOTAL	NO_NOG	NOHEIGHT	KRAFT1	KRAFT2	KRAFT3	KRAFT4	KRAFT5	KRAFTM	STEM1	STEM2
1	01447/84	MT.PROVINCE	Cashel	1292	898	6	5	0.25	6.25	30	66	0	16	13.3125	9.24	100	0.014411	0.230579	20.03125	12.875	5.0625	0.875	6.25	16	16	15	0	6.25	75	18.75	0	3.125	6.4	0	0
1	01448/84	BENQUET	Cashel	1292	898	5	4	0.25	12.25	20	60	0	16	14.1	8.58125	100	0.015874	0.253984	21	12.6875	4.4375	0.5	12.5	16	16	16	0	6.25	62.5	31.25	0	3.25	6.0625	0	6.25	0
1	01515/85	DATHIEN	Cashel	1292	898	3	1	6.25	42.25	3	45	0	16	14.275	9.530769	100	0.016789	0.268629	24.607143	9.375			0.25	18.75	16	16	13	12.5	18.75	31.25	37.5	0	2.9375	6.642857	0	7.14286
1	01517/85	HOTHIEN	Cashel	1292	898	2	1	12.25	42.25	2	30	0	16	15.34	9.346667	93.75	0.019362	0.290437	24.892857	9.6			0.142857	28.571429	16	15	15	6.666667	20	26.666667	46.666667	0	3.133333	5.666667	0	6.66667
1	01519/85	LANGHANH	Cashel	1292	898	4	3	2.25	20.25	12	52	0	16	14.45625	8.16	100	0.016592	0.265472	23.1	10.6875	4.25	0.375	25	16	16	15		6.666667	0	60	33.333333	0	3.2	7.733333	0	0
1	01521/85	NONGKRATING	Cashel	1292	898	5	5	0.25	6.25	25	55	0	16	13.94	7.869231	93.75	0.017205	0.258069	23.153846	10.33333	5.466667		0.266667	13.333333	16	15	13	0	13.33333	80	6.6666667	0	2.933333	5.615385	0	0
1	01523/85	DOIINTHANON	Cashel	1292	898	6	1	0.25	42.25	6	90	0	16	15.5933333	7.833333	100	0.019371	0.290567	24.433333	11.2	4.466667		0.125	12.5	16	15	15	0	6.666667	73.333333	20	0	3.133333	5.066667	0	6.66667
1	01525/85	NAMNOW	Cashel	1292	898	6	4	0.25	12.25	24	72	0	16	14.63125	6.953333	100	0.017049	0.272787	23.966667	13.8125	5.9375	0.1875	6.25	16	16	15	0	6.25	87.5	6.25	0	3		6.6	0	0
1	01572/85	COTOMINES	Cashel	1292	898	2	2	12.25	30.25	4	28	0	16	13.54	9.627273	93.75	0.015288	0.229325	24.076923	12.61538	4.538462		0.571429	26.666667	16	15	11	0	0	92.857143	7.1428571	0	3.071429	6.75	0	0
1	01624/86	BAOSHAN(1)	Cashel	1292	898	5	2	0.25	30.25	10	70	0	16	9.7125	5.725	100	0.011398	0.182366	25.59375	4.6875	5.0625	1.125	12.5	16	16	16	0	25	18.75	50	6.25	3.375	5.875	0	0	6.25
1	01632/86	CEHENG	Cashel	1292	898	5	3	0.25	20.25	15	65	0	16	7.93333333	3.13125	100	0.006173	0.092592	28.136364	3.6875	4.6875	0.8125	0	16	15	16	0	0	81.25	12.5	6.25	3.25	6.0625	0	0	0
1	01633/86	SHANGSI	Cashel	1292	898	3	2	6.25	30.25	6	42	0	16	13.2357143	7.371429	87.5	0.017021	0.2383	27.821429	8.071429	5.571429		0.857143	21.428571	16	14	14	0	21.42857	35.714286	35.714286	7.142857	3.285714	5.5	0	0
1	01636/86	JINGDUNG(2)	Cashel	1292	898	4	5	2.25	6.25	20	44	0	16	13.7125	7.133333	100	0.015423	0.246774	24.033333	8.6875	5.866667		0.4375	0	16	16	15	0	6.25	37.5	50	6.25	3.5625	4.733333	0	0
1	01637/86	JINGHONG	Cashel	1292	898	1	2	20.25	30.25	2	14	0	16	13.7875	7.735714	100	0.016345	0.261512	25.5	9.6875	5	0.375	6.25	16	16	14	0	6.25	50	43.75	0	3.375		5.785714	0	0
1	01638/86	LANCANG	Cashel	1292	898	4	1	2.25	42.25	4	60	0	16	15.3375	7.9375	100	0.019206	0.3073	24.3125	9.5	0.3125	31.25	16	16	16	0	0	75	18.75	6.25	3.3125	5.9375	0	0	0	6.25
1	01680/86	DAKHA	Cashel	1292	898	1	1	20.25	42.25	1	15	0	16	14.01875	8.275	100	0.015986	0.255774	33.307692	9.933333			0.2	33.333333	16	16	12	0	23.07692	53.846154	23.076923	0	3	4.916667	0	16.6667
1	01772/88	ZOKHUA	Cashel	1292	898	4	2	2.25	30.25	8	56	0	16	12.96875	7.2375	100	0.018153	0.290442	24.666667	10.0625	5.625	0.9375	6.25	16	16	16	6.25	0	68.75	25	0	3.125	5.75	6.25	0	0
1	01773/88	AUNGBAN	Cashel	1292	898	5	1	0.25	42.25	5	75	0	16	14.22	6.828571	93.75	0.016813	0.252189	23.566667	9			0.266667	0	16	15	14	0	6.666667	66.666667	26.666667	0	3.2	5.214286	0	0
1	01777/88	JOHNMEIKLECSO	Cashel	1292	898	6	2	0.25	30.25	12	84	0	16	14.00625	8.528571	100	0.016088	0.257401	23.066667	11.1875	5.8125	0.3125	12.5	16	16	14	0		14.28571	50	28.571429	7.142857	3.285714	6.230769	0	0
1	01778/88	ZAMBIACSO	Cashel	1292	898	4	4	2.25	12.25	16	48	0	16	14.59375	8.5	100	0.017015	0.272241	21.65625	13.5	4.75	0.3125	18.75	16	16	14	0	25	31.25	37.5	6.25	3.25		7.214286	0	0
1	01786/88	MORARANO	Cashel	1292	898	6	3	0.25	20.25	18	78	0	16	14.2466667	7.426667	93.75	0.016441	0.246622	24.333333	12	6.466667		0.666667	13.333333	16	15	15	0	0	73.333333	26.666667	0	3.266667	5.666667	0	0
2	01447/84	MT.PROVINCE	Cashel	1292	898	5	10	0.25	6.25	50	30	0	16	12.8125	8.107143	100	0.013075	0.209207	22.6875	12.125	4.1875	0.75	12.5	16	16	14	0	18.75	62.5	18.75	0	3		6.214286	0	0
2	01448/84	BENQUET	Cashel	1292	898	5	7	0.25	0.25	35	45	0	16	13.3625	8.228571	100	0.014942	0.239079	21.107143	11.3125	4	0.3125	18.75	16	16	14	0	0		66.666667	33.333333	0	3.333333	7	0	0
2	01515/85	DATHIEN	Cashel	1292	898	4	10	2.25	6.25	40	24	0	16	13.9	7.530769	87.5	0.015363	0.215083	23.75	11	4.642857		0.571429	14.285714	16	14	13	0	0	57.142857	42.857143	0	3.428571	7	0	0
2	01517/85	HOTHIEN	Cashel	1292	898	5	11	0.25	12.25	55	25	0	16	14	7.825	100	0.015514	0.248228	23.233333	10.1875	4.125	0.625	12.5	16	16	12	0	0	43.75	50	6.25	3.625		6.615385	0	0
2	01519/85	LANGHANH	Cashel	1292	898	3	10	6.25	6.25	30	18	0	16	13.2133333	7.758333	93.75	0.014065	0.210971	23.5	12.6	4.466667		0.466667	0	16	15	12	0	6.666667	80	13.333333	0	3.066667	7.416667	0	0
2	01521/85	NONGKRATING	Cashel	1292	898	6	9	0.25	2.25	54	42	0	16	12.50625	5.821429	100	0.012539	0.200619	23.785714	11.25	5.125	0.375	0	16	16	14	0	12.5	87.5	0	0	2.875		6.214286	0	0
2	01523/85	DOIINTHANON	Cashel	1292	898	4	7	2.25	0.25	28	36	0	16	14.4	7.29	93.75	0.017389	0.260835	21.6	12.73333	4.333333		0.6	0	16	15	10	0	0	60	40	0	3.4	5.4	0	0
2	01525/85	NAMNOW	Cashel	1292	898	4	11	2.25	12.25	44	20	0	16	12.925	6.493333	100	0.013359	0.213746	22.833333	11.9375	5.625	0.3125	0	16	16	15	0	12.5	62.5	25	0	3.125	6.375	0	6.25	0
2	01572/85	COTOMINES	Cashel	1292	898	5	6	0.25	2.25	30	50	0	16	12.8428571	8.65	87.5	0.013349	0.18689	20.181818	12.92857	4.071429		0.714286	0	16	14	8	0	0	92.307692	7.6923077	0	3.076923	6.125	0	0
2	01624/86	BAOSHAN(1)	Cashel	1292	898	4	8	2.25	0.25	32	32	0	16	7.8125	3.433333	100	0.006158	0.09853	23.5	4.3125	5.5	0.8125	0	16	16	15	0	0	93.75	0	6.25	3.125		5.6	6.666667	0
2	01632/86	CEHENG	Cashel	1292	898	4	6	2.25	2.25	24	40	0	16	7.07692308	2.7125	100	0.004388	0.057048	23	4.6875	4.9375	0.5	6.25	16	16	13	16	0	0		92.307692	7.6923077	0	3.076923	5.307692	0	15.3846
2	01633/86	SHANGSI	Cashel	1292	898	5	8	0.25	0.25	40	40	0	16	10.24	4.457143	93.75	0.009455	0.141829	22.55	6.4	5.866667		0.4	6.6666667	16	15	14	0	0	73.333333	20	6.666667	3.333333	6.428571	0	0
2	01636/86	JINGDUNG(2)	Cashel	1292	898	5	9	0.25	2.25	45	35	0	16	12.4875	6.253333	100	0.012961	0.20738	24.433333	9.75	5.5625	0.75	0	16	16	15	0	0	37.5	56.25	6.25	3.6875		5.8	0	0
2	01637/86	JINGHONG	Cashel	1292	898	3	9	6.25	2.25	27	21	0	16	13.3	7.086667	100	0.013968	0.223482	23.1875	10.0625	5.3125	0.5	0	16	16	15	0	0	31.25	68.75	0	3.6875		6.214286	0	0
2	01638/86	LANCANG	Cashel	1292	898	6	10	0.25	6.25	60	36	0	16	12.61875	6.685714	100	0.01427	0.228316	22.714286	10.3125	4.875	0.5625	0	16	16	14	0	6.25	37.5	50	6.25	3.5625	6.5	0	0	0
2	01680/86	DAKHA	Cashel	1292	898	6	6	0.25	2.25	36	60	0	16	14.16875	8.32	100	0.016085	0.257355	21.115385	10.625	5.625	0.4375	12.5	16	16	10	0	6.25	43.75	50	0	3.4375		6.2	0	0
2	01772/88	ZOKHUA	Cashel	1292	898	4	9	2.25	2.25	36	28	0	16	10.23125	5.11875	100	0.010502	0.168038	21.5	9.5625	5.0625	0.625	0	16	16	16	0	6.25	81.25	12.5	0	3.0625	5.9375	0	0	0
2	01773/88	AUNGBAN	Cashel	1292	898	6	7	0.25	0.25	42	54	0	16	11.9533333	5.566667	93.75	0.013008	0.195124	22.125	9.2	4.666667		0.533333	0	16	15	15	0	6.666667	53.333333	40	0	3.333333	4.666667	13.33333	0
2	01777/88	JOHNMEIKLECSO	Cashel	1292	898	6	11	0.25	12.25	66	30	0	16	12.9733333	7.457143	93.75	0.01344	0.201594	21.214286	11.6	4.133333		0.266667	13.333333	16	15	7	0	12.5	37.5	50	0	3.375	6.777778	0	0
2	01778/88	ZAMBIACSO	Cashel	1292	898	6	8	0.25	0.25	48	48	0	16	12.5066667	7.228571	93.75	0.012476	0.187137	22.5	12.2	4.466667		0.2	6.6666667	16	15	14	0	6.666667	40	53.333333	0	3.466667	7.785714	0	0
2	01786/88	MORARANO	Cashel	1292	898	3	11	6.25	12.25	33	15	0	16	12.28125	6.784615	100	0.012133	0.194129	23.692308	11.8125	5.4375	0.5625	0	16	16	13	6.25	6.25	56.25	31.25	0	3.125		7.076923	0	0
3	01447/84	MT.PROVINCE	Cashel	1292	898	8	10	6.25	6.25	80	48	0	16	12.0375	7.676923	100	0.011702	0.187236	21.233333	12.125	4.25	0.5625	12.5	16	16	13	0		14.28571	78.571429	7.1428571	0	2.928571	6.428571	0	0
3	01448/84	BENQUET	Cashel	1292	898	7	9	2.25	2.25	63	49	0	16	12.4933333	8.078571	93.75	0.012715	0.190719	20.392857	10.8	4.8		0.6	6.6666667	16	15	14	0	0	71.428571	28.571429	0	3.533333	7.214286	0	0
3	01515/85	DATHIEN	Cashel	1292	898	9	5	12.25	6.25	45	99	0	16	15.3933333	9.718182	93.75	0.018903	0.283551	21.9	11.27273	4.454545		0.333333	20	16	15	11	0	0	90.909091	0.9090091	0	3.090909	7.714286	0	0
3	01517/85	HOTHIEN	Cashel	1292	898	9	6	12.25	2.25	54	90	0	16	14.8125	8.178571	100	0.017387	0.278192	21.34375	11	5	0.375	18.75	16	16	14	0	0	7.142857	35.714286	57.142857	0	3.5	7.1875	0	0
3	01519/85	LANGHANH	Cashel	1292	898	9	8	12.25	0.25	72	72	0	16	13.64375	7.330769	100	0.014978	0.239645	21.5	12.5	5.6875	0.375	6.25	16	16	13	0	0		53.333333	46.666667	0	3.466667	7.692308	0	0
3	01521/85	NONGKRATING	Cashel	1292	898	8	5	6.25	6.25	40	88	0	16	13.625	6.25	100	0.014668	0.23468	22.21875	11.5625	4.875	0.25	0	16	16	16	0	6.25	81.25	12.5	0	3.0625	6.25	0	0	0
3	01523/85	DOIINTHANON	Cashel	1292	898	7	10	2.25	6.25	70	42	0	16	13.9625	7.253333	100	0.015426	0.246813	22.1875	12.3125	4.75	0.4375	0	16	16	15	0		6.666667	53.333333	40	0	3.333333	5.933333	0	0
3	01525/85	NAMNOW	Cashel	1292	898	7	6	2.25	2.25	42	70	0	16	13.9875	6.493333	100	0.015987	0.25579	20.84375	11.9375	5.25	0.1875	0	16	16	15	0	6.25	75	18.75	0	3.125		6.466667	0	0
3	01572/85	COTOMINES	Cashel	1292	898	7	4	2.25	12.25	28	84	0	16	14.01875	8.628571	100	0.015825	0.253207	18.46875	13.5	4.4375	0.5625	6.25	16	16	14	0		13.33333	86.666667	0	0	2.866667	6.428571	0	0	0
3	01624/86	BAOSHAN(1)	Cashel	1292	898	8	8	6.25	0.25	64	64	0	16	7.49333333	3.325	100	0.005458	0.081863	23.333333	3.533333	4.866667		0.625	25	16	15	16	0	20	73.333333	6.6666667	0	2.866667	5.866667	0	6.66667
3	01632/86	CEHENG	Cashel	1292	898	8	4	6.25	12.25	32	96	0	16	7.05	2.28	93.75	0.004399	0.043993		4	4.5		0.333333	0	16	10	15	0	20	40	40	0	3.2	5.7	0	0	0
3	01633/86	SHANGSI	Cashel	1292	898	8	6	6.25	2.25	48	80	0	16	9.28125	3.926667	100	0.007301	0.116822	22	6.125	4.5625	0.375	0	16	16	15	0	6.25	93.75	0	0	2.9375		6.133333	0	0	0
3	01636/86	JINGDUNG(2)	Cashel	1292	898	7	5	2.25	6.25	35	77	0	16	13.525	6.775	100	0.014784	0.236538	23.8	10	7	0.5625	6.25	16	16	12	0	0		35.714286	64.285714	0	3.642857	5.923077	0	0	0
3	01637/86	JINGHONG	Cashel	1292	898	9	4	12.25	12.25	36	108	0	16	15.36	7.992857	93.75	0.018676	0.280143	22.166667	10.64286	5.714286		0.066667	0	16	15	14	0	0	71.428571	28.571429	0	3.285714	6.066667	0	0	0
3	01638/86	LANCANG	Cashel	1292	898	7	8	2.25	0.25	56	56	0	16	13.41875	7.075	100	0.015415	0.246645	23.03125	10.375	5.6875	0.625	6.25	16	16	16	0	6.25	43.75	50	0	3.4375	6.5625	0	0	0
3	01680/86	DAKHA	Cashel	1292	898	8	9	6.25	2.25	72	56	0	16	13.6	8.041667	93.75	0.015335	0.230032	22.071429	9.2	5.266667		0.733333	20	16	15	12	0	6.666667	60	33.333333	0	3.266667	6.583333	0	0	0
3	01772/88	ZOKHUA	Cashel	1292	898	7	7	2.25	0.25	49	63	0	16	11.575	5.65625	100	0.012382	0.198118	19.7	9.9375	6	0.3125	0	16	16	16	0	0	75	25	0	3.25	6.0625	0	0	0
3	01773/88	AUNGBAN	Cashel	1292	898	9	9	12.25	2.25	81	63	0	16	11.16	6.038462	93.75	0.011862	0.177924	21.461538	9.666667	5.6		0.466667	0	16	15	13	0	6.666667	60	33.333333	0	3.266667	5.769231	0	0	0
3	01777/88	JOHNMEIKLECSO	Cashel	1292	898	9	10	12.25	6.25	90	54	0	16	12.61875	7.358333	100	0.012861	0.205783	20.571429	11.21429	4.428571		0.0625	6.25	16	16	12	0	15.38462	46.153846	38.461538	0	3.230769	7.083333	0	0	0
3	01778/88	ZAMBIACSO	Cashel	1292	898	8	7	6.25	0.25	56	72	0	16	13.575	7.675	100	0.01467	0.234726	20.964286	13.71429	4.928571		0.125	0	16	16	12	0	7.142857	71.428571	21.428571	0	3.142857	7.916667	0	0	0
3	01786/88	MORARANO	Cashel	1292	898	9	7	12.25	0.25	63	81	0	16	12.9375	7.806667	100	0.013492	0.21588	21.666667	10.5625	5.625	0.1875	37.5	16	16	15	0	0	62.5	31.25	6.25	3.4375		7.266667	0	0	0
4	01447/84	MT.PROVINCE	Cashel	1292	898	11	4	30.25	12.25	44	132	0	16	14.0428571	8.092857	87.5	0.015699	0.219786	21.392857	12.42857	5.357143		0.785714	21.428571	16	14	14	0	0	92.857143	7.1428571	0	3.071429	6.285714	0	0	0
4	01448/84	BENQUET	Cashel	1292	898	10	7	20.25	0.25	70	90	0	16	14.1333333	8.315385	100	0.016496	0.247445	22.166667	11.6	4.066667		0.866667	13.333333	16	15	13	0	0	73.333333	26.666667	0	3.266667	6.428571	0	0	0
4	01515/85	DATHIEN	Cashel	1292	898	11	6	30.25	2.25	66	110	0	16	15.01875	8.292857	100	0.019191	0.307049	23.90625	9.933333	5		0.5	12.5	16	16	14	0	6.666667	73.333333	20	0	3.133333	7.071429	0	0	0
4	01517/85	HOTHIEN	Cashel	1292	898	10	10	20.25	6.25	100	60	0	16	12.9625	7.464286	100	0.01372	0.219527	21.6875	9.6875	4	0.3125	31.25	16	16	14	12.5	0	50	37.5	0	3.125	7	0	0	0
4	01519/85	LANGHANH	Cashel	1292	898	12	8	42.25	0.25	96	96	0	16	14.275	8.366667	100	0.016234	0.259737	22.833333	10.6875	4.1875	0.4375	18.75	16	16	15	0	6.25	68.75	25	0	3.1875		6.466667	0	6.66667	0
4	01521/85	NONGKRATING	Cashel	1292	898	10	9	20.25	2.25	90	70	0	16	12.63125	5.9	100	0.013269	0.212308	23.433333	10.125	5.25	0.3125	0	16	16	15	0	6.25	93.75	0	0	2.9375		5.933333	0	0	0
4	01523/85	DOIINTHANON	Cashel	1292	898	11	8	30.25	0.25	88	88	0	16	13	6.563636	100	0.01405	0.224794	23.545455	10.8	6.666667		0.375	6.25	16	16	11	0	7.692308	69.230769	23.076923	0	3.153846	6.636364	0	0	0
4	01525/85	NAMNOW	Cashel	1292	898	12	6	42.25	2.25	72	120	0	16	14.3461538	6.384615	81.25	0.016406	0.21328	23.333333	11.84615	5.538462		0.153846	0	16	13	13	0	0	84.615385	15.384615	0	3.153846	6.769231	0	0	0

Plot Data Set - Cashel

STEM 3	STEM 4	STEM 5	STEM 6	STEM 7	STEM 8	STEM 9	BRARATIO	FO_INDEX	INTNODE	CROWNPCT	FO_POS	TOP	VTREE	MFF	FCL	CCC	OCC	LOTH	APHID	DG	HG	PILOCORR	VHA
0	6.6666667	6.666667	40	33.333333	13.333333	0		50.8085	0.75391		0.496364	6.25	0.070958	1	1	1	1	1.9375	0	0	13.543079		19.997729	97.336343
6.25	6.25	43.75	25	12.5	0			21.714124	0.75111		0.4525	12.5	0.070412	1	1	1	1	1.5625	0	0	14.213814		21.024563	96.587381
0	14.285714	7.142857	7.1428571	14.285714	35.714286	14.285714		38.193802	0.94806		0.465	0	0.089106	1	1	1	1	1.1875	0	0	14.617883		24.718636	122.23105
6.666667	20	20	13.333333	6.6666667	13.333333	13.333333		27.772487	1.06066		0.31	0	0.092817	1	1	1	1	1.2	0	0	15.698132		25.063356	119.36377
0	0	6.666667	6.6666667	13.333333	53.333333	20		19.739248	0.93248		0.401667	18.75	0.070126	1	1	1	1	1.1875	0	0	14.531711		23.158492	96.194159
7.692308	23.076923	7.692308	30.769231	23.076923	7.6923077	0		70.436869	0.70768		0.2625	6.25	0.07609	1	1	1	1	1.33333	12.5	0	14.797567		23.206778	97.852947
20	6.6666667	26.66667	20	6.6666667	13.333333	0		33.039216	0.67438		0.32	6.25	0.077416	1	1	1	1	1.53333	0	0	15.701656		24.568042	99.558
0	6.6666667	6.666667	46.666667	13.333333	13.333333	13.333333		19.213934	0.49711		0.39	0	0.062854	1	1	1	1	1.0625	6.25	0	14.730559		24.038927	86.219661
0	8.3333333	0	33.333333	25	33.333333	0		65.25335	0.74037		0.446667	0	0.082498	1	1	1	1	1.13333	18.75	6.25	13.949146		24.100919	106.0929
6.25	31.25	6.25	50	0	0			369.5671	1.22156		0.07	0	0.043607	1	1	1	1	2.375	0	0	12.044241		25.294699	59.817465
0	25	43.75	31.25	0	0			123.65859	0.73115		0.104444	0	0.019533	1	1	1	1	1.875	0	0	8.8635584		27.779396	25.119498
0	21.428571	21.42857	42.857143	14.285714	0	0		251.06939	0.87721		0.218571	0	0.068916	1	1	1	1	2.28571	6.25	87.5	14.718574		27.782244	82.718233
0	33.333333	60	6.6666667	0	0	0		36.706536	0.79782		0.271667	0	0.060504	1	1	1	1	1.6875	6.25	0	14.010621		24.060355	82.996266
0	14.285714	21.42857	42.857143	14.285714	7.1428571	0		55.989072	0.8281		0.268	6.25	0.067058	1	1	1	1	1.86667	12.5	6.25	14.422942		25.50637	91.985633
18.75	56.25	12.5	6.25	0				46.081303	0.85206		0.366	6.25	0.078752	1	1	1	1	2	0	0	15.634677		24.428339	108.0274
16.66667	8.3333333	8.333333	16.666667	33.333333	0	0		63.074004	0.87991		0.24	0	0.07096	1	1	1	1	2.33333	25	0	14.263831		33.373674	97.338176
6.25	25	31.25	18.75	12.5	0			957.38942	0.70004		0.15375	0	0.072636	1	1	1	1	2.8125	0	100	15.199794		24.645992	99.638237
0	50	7.142857	21.428571	14.285714	7.1428571	0		36.881119	0.76747		0.155	0	0.061453	1	1	1	1	2	0	93.75	14.62801		23.600081	79.028623
0	23.076923	15.38462	15.384615	23.076923	7.6923077	15.384615		19.6448	0.7491		0.532	0	0.076213	1	1	1	1	1.0625	12.5	87.5	14.30911		23.115324	104.54512
0	7.1428571	7.142857	14.285714	14.285714	42.857143	14.285714		27.233083	0.64962		0.298	0	0.074584	1	1	1	1	1.375	12.5	0	14.715829		21.717232	102.30981
0	20	26.66667	26.666667	20	6.6666667	0		21.340619	0.60251		0.394444	0	0.064592	1	1	1	1	1.6	0	0	14.46566		24.368715	83.065401
0	0	28.57143	42.857143	7.1428571	21.428571	0		48.929203	0.65564		0.32375	0	0.055641	1	1	1	1	1.5625	12.5	6.25	12.900194		22.6171	76.324889
0	0	0	35.714286	42.857143	7.1428571	14.285714		28.446014	0.73685		0.388	12.5	0.068823	1	1	1	1	1.4375	6.25	6.25	13.790441		21.126438	94.406737
0	0	0	30.769231	46.153846	15.384615	7.6923077		22.038198	0.68476		0.437143	12.5	0.060382	1	1	1	1	1.14286	6.25	0	13.983204		23.759812	72.474689
0	0	7.692308	38.461538	38.461538	15.384615	0		50.818285	0.75211		0.351429	12.5	0.063703	1	1	1	1	1.25	12.5	0	14.051824		23.264781	87.384344
0	0	8.333333	0	33.333333	58.333333	0		28.31142	0.59136		0.368	18.75	0.063291	1	1	1	1	1.13333	18.75	6.25	13.379287		23.525178	81.392547
0	0	21.42857	42.857143	28.571429	7.1428571	0		29.893506	0.48012		0.298	6.25	0.042406	1	1	1	1	1.3125	6.25	0	12.632621		23.734939	58.169818
0	10	60	20	0	10	0		37.905704	0.52358		0.305	0	0.072152	1	1	1	1	1.06667	31.25	0	14.876648		21.698322	92.787469
0	12.5	25	43.75	6.25	6.25			17.624463	0.51226		0.37	0	0.049152	1	1	1	1	1.1875	0	0	13.039363		22.796432	67.424215
12.5	12.5	0	12.5	50	12.5	0		21.481213	0.59967		0.467778	6.25	0.065631	1	1	1	1	1.14286	31.25	0	13.034569		20.163468	78.77507
0	6.6666667	13.33333	60	6.6666667	6.6666667	0		103.9091	0.70411		0.112	0	0.021539	1	1	1	1	1.1875	6.25	0	8.8530362		23.321728	29.545766
0	0	23.07692	46.153846	15.384615	0	0		83.036275	0.49369		0.122	0	0.01449	1	1	1	1	1.25	0	6.25	7.4733372		22.958181	16.150162
0	0	0	57.142857	42.857143	0	0		34.444452	0.6277		0.236	0	0.028976	1	1	1	1	3	6.25	6.25	10.969959		22.418196	37.263189
0	0	26.66667	66.666667	6.6666667	0	0		162.77965	0.6052		0.327778	0	0.046344	1	1	1	1	2.25	6.25	0	12.843724		24.346768	63.571556
0	0	28.57143	35.714286	21.428571	14.285714	0		22.152764	0.66426		0.391429	6.25	0.053333	1	1	1	1	1.6875	0	0	13.333042		23.153057	73.159486
	7.1428571	14.28571	21.428571	35.714286	21.428571	0		170.2328	0.62072		0.257143	6.25	0.05311	1	1	1	1	1.8125	12.5	0	13.476461		22.624524	72.852928
0	0	20	50	20	10	0		19.419032	0.78926		0.458	0	0.074437	1	1	1	1	1.125	31.25	0	14.307843		21.194983	102.10816
0	18.75	68.75	12.5	0	0			134.47998	0.49822		0.105	0	0.035329	1	1	1	1	2.1875	0	18.75	11.561412		21.288537	48.462203
6.666667	20	13.33333	40	6.6666667	0	0		243.44619	0.56939		0.142	0	0.043517	1	1	1	1	1.53333	0	6.25	12.866986		22.055522	55.963818
0	11.111111	0	11.111111	55.555556	22.222222	0		15.169704	0.66719		0.4725	37.5	0.047209	1	1	1	1	1.4	12.5	0	13.078583		21.164037	60.711094
0	0	0	14.285714	14.285714	50	21.428571		20.168651	0.57632		0.326667	12.5	0.049904	1	1	1	1	1.4	6.25	0	12.600899		22.44116	64.177399
0	0	0	15.384615	61.538462	23.076923	0		31.276126	0.56378		0.368571	6.25	0.048442	1	1	1	1	1.25	18.75	0	12.42661		23.619283	66.449359
0	0	14.28571	35.714286	42.857143	7.1428571	0		16.996972	0.60377		0.478889	12.5	0.050409	1	1	1	1	1.0625	6.25	6.25	12.203995		21.130049	69.14788
0	0	0	21.428571	42.857143	28.571429	7.1428571		114.60235	0.77049		0.366667	0	0.054773	1	1	1	1	1.33333	6.25	0	12.720928		20.310998	70.437955
0	0	0	7.1428571	21.428571	64.285714	7.1428571		28.714329	0.86166		0.536667	12.5	0.095608	1	1	1	1	1.6	0	0	15.510921		22.019957	122.95223
0	0	6.25	18.75	25	50	0		23.665115	0.73874		0.426	6.25	0.074101	1	1	1	1	1.5	0	43.75	14.875777		21.420866	101.64791
0	0	0	0	38.461538	53.846154	7.6923077		24.073579	0.55336		0.422	6.25	0.05874	1	1	1	1	1.75	12.5	0	13.806769		21.529102	80.575823
6.25	18.75	31.25	31.25	12.5	0			19.329769	0.51238		0.3975	0	0.049338	1	1	1	1	1.25	6.25	12.5	13.662997		22.208278	67.678406
0	6.6666667	26.66667	40	20	6.6666667	0		17.934849	0.55749		0.428333	6.25	0.058346	1	1	1	1	1.125	0	0	14.011736		22.201922	80.036076
0	0	0	60	33.333333	6.6666667	0		69.423077	0.52298		0.17	0	0.055248	1	1	1	1	1.375	6.25	18.75	14.264269		20.860016	75.785507
0	7.1428571	14.28571	28.571429	28.571429	21.428571	0		26.750681	0.61094		0.435	0	0.070056	1	1	1	1	1.375	12.5	43.75	14.192053		18.487321	96.099103
0	0	20	40	33.333333	0	0		139.5969	0.88833		0.151111	0	0.018493	1	1	1	1	1	0	6.25	8.3342266		23.102202	23.781841
10	0	20	50	20	0	0		131.66667	0.61933		0.04	0	0.013978	1	1	1	1	1.2	0	6.25	7.4827134			11.983526
0	6.6666667	13.33333	53.333333	13.333333	13.333333	0		16.828025	0.57033		0.261667	0	0.021771	1	1	1	1	2.6875	6.25	0	9.6398587		21.928678	29.863825
7.692308	7.6923077	0	53.846154	30.769231	0	0		21.296244	0.76231		0.35	6.25	0.052643	1	1	1	1	2.625	25	37.5	13.71696		23.806862	72.212905
0	6.6666667	13.33333	46.666667	33.333333	0	0		16.170213	0.74531		0.47	6.25	0.075082	1	1	1	1	1.86667	0	0	15.417436		22.284165	96.555738
0	0	50	43.75	6.25	0			39.857047	0.64593		0.31125	0	0.058636	1	1	1	1	2	0	0	14.006963		23.005566	80.433999
8.333333	0	0	25	50	16.666667	0		122.01056	0.87407		0.334286	0	0.066489	1	1	1	1	1.46667	18.75	0	13.970636		22.082294	85.505407
12.5	6.25	50	25	6.25	0	213.625			0.5292		0.0875	0	0.041659	1	1	1	1	2.25	0	62.5	12.553635		19.555903	57.144952
0	0	46.15385	30.769231	23.076923	0	0		236.38145	0.58		0.141667	0	0.045112	1	1	1	1	1.86667	12.5	6.25	12.286822		21.292061	58.013884
0	0	8.333333	16.666667	33.333333	41.666667	0		13.918919	0.59695		0.74	12.5	0.052364	1	1	1	1	1	6.25	0	12.794164		20.492731	71.830447
0	0	0	8.3333333	0	83.333333	8.3333333		53.793103	0.52797		0.29	12.5	0.062222	1	1	1	1	1.5625	12.5	12.5	13.664324		20.971463	85.352204
0	0	0	13.333333	46.666667	40	0		15.464986	0.73061		0.5	0	0.056041	1	1	1	1	1.1875	6.25	12.5	13.104293		21.630748	76.874319
0	7.1428571	14.28571	35.714286	28.571429	14.285714	0		43.769533	0.65884		0.364286	0	0.065619	1	1	1	1	1.35714	0	37.5	14.135265		21.413206	78.760444
0	7.1428571	7.142857	28.571429	50	7.1428571	0		87.264639	0.68338		0.466667	0	0.066994	1	1	1	1	1.33333	6.25	18.75	14.489767		22.193689	86.155345
0	7.1428571	0	14.285714	35.714286	42.857143	0		163.80846	0.82664		0.45	6.25	0.081434	1	1	1	1	1.3125	6.25	6.25	15.6283		23.998579	111.70612
0	0		28.571429	42.857143	28.571429	0		29.128841	0.77845		0.3225	0	0.056116	1	1	1	1	1.125	12.5	25	13.214528		21.628164	76.976889
0	0	13.33333	13.333333	46.666667	20	0		23.127674	0.93282		0.378571	0	0.071478	1	1	1	1	1.75	6.25	6.25	14.373891		22.881008	98.048789
0	0	13.33333	80	6.6666667	0	0		32.532967	0.54784		0.336	0	0.045214	1	1	1	1	1.1875	6.25	6.25	12.995432		23.370862	62.021915
0	0	18.18182	27.272727	27.272727	27.272727	0		40.682696	0.57773		0.252	0	0.05114	1	1	1	1	1.75	31.25	18.75	13.372102		23.518387	70.151448
0	7.6923077	0	30.769231	30.769231	30.769231	0		15.169414	0.50306		0.475	0	0.056053	1	1	1	1	1.15385	0	12.5	14.4501		23.408298	62.473316

Plot Data Set - Cashel

BLOCK	SEED LOT	PROVNAME	SITE	DATE EST	DATE ASS	PLOT X	PLOT Y	PLOT TX	PLOT TY	PLOT TX_Y	_TYPE_	_FREQ_	DBH 1	HEIGHT	SURV	GMEAN	GSUM	MEAN PILO	WHORL S	BRANCH	DIABRA	FORK	FOXTAIL	NO_TOTAL	NO_G	NOHEIGHT	KRAFT 1	KRAFT 2	KRAFT 3	KRAFT 4	KRAFT 5	KRAFT	STEM	STEM 1	STEM 2
4	01572/85	COTOMINES	Cashel	1292	898	11	9	30.25	2.25	99	77	0	16	12.35	8.314286	100	0.013683	0.218924	21.035714	13.13333	3.2		0.625	6.25	16	16	14	6.25	12.5	81.25	0	0	2.75	6.333333	0	0
4	01624/86	BAOSHAN(1)	Cashel	1292	898	11	7	30.25	0.25	77	99	0	16	8.45625	3.81875	100	0.008323	0.133163	24.333333	3.9375	6.0625	1.1875	0	16	16	16	0	0	87.5	12.5	0	3.125	6.25	0	0	0
4	01632/86	CEHENG	Cashel	1292	898	10	4	20.25	12.25	40	120	0	16	7.72666667	3.19375	100	0.00607	0.091047	24.166667	4.125	6.5625	0.6875	0	16	15	16	0	0	100	0	0	3		6.333333	0	0
4	01633/86	SHANGSI	Cashel	1292	898	11	5	30.25	6.25	55	121	0	16	9.13125	3.873333	100	0.007732	0.12371	25.5	5.625	6.6875	0.5	0	16	16	15	0	12.5	75	12.5	0	3		6.066667	0	0
4	01636/86	JINGDUNG(2)	Cashel	1292	898	12	9	42.25	2.25	108	84	0	16	13.0375	6.4625	100	0.014044	0.224706	23.25	8.625	5.9375	0.375	6.25	16	16	16	0	12.5	50	31.25	6.25	3.3125	6.1875	0	0	0
4	01637/86	JINGHONG	Cashel	1292	898	10	8	20.25	0.25	80	80	0	16	13.1866667	7.064286	93.75	0.014331	0.214966	24.133333	10	6.428571		0.266667	0	16	15	14	0	7.142857	21.428571	71.428571	0	3.642857	6.357143	0	0
4	01638/86	LANCANG	Cashel	1292	898	10	5	20.25	6.25	50	110	0	16	15.1625	8.169231	100	0.018404	0.294464	23.2	10.4	5.866667		0.3125	6.25	16	16	13	0	0	66.666667	33.333333	0	3.333333	5.857143	0	7.14286
4	01680/86	DAKHA	Cashel	1292	898	12	10	42.25	6.25	120	72	0	16	12.86875	8.4	100	0.013533	0.21653	22.3	9.8	4.733333		0.3125	18.75	16	16	13	0	0	46.666667	53.333333	0	3.533333	6.5	0	0
4	01772/88	ZOKHUA	Cashel	1292	898	11	10	30.25	6.25	110	66	0	16	10.9933333	6.025	93.75	0.011265	0.16898	21	9.615385	5.923077		0.4	0	16	15	12	0	16.66667	50	33.333333	0	3.166667	7.25	0	0
4	01773/88	AUNGBAN	Cashel	1292	898	12	7	42.25	0.25	84	108	0	16	13.0933333	5.823077	93.75	0.01606	0.240907	23.785714	9	6.466667		0.533333	0	16	15	13	0	0	33.333333	66.666667	0	3.666667	5.615385	0	0
4	01777/88	JOHNMEIKLECSO	Cashel	1292	898	12	5	42.25	6.25	60	132	0	16	13.1533333	7.4375	100	0.015049	0.225731	22.1	11.72727	4.454545		0.6875	0	16	15	8	0	27.27273	63.636364	9.0909091	0	2.818182	7	0	0
4	01778/88	ZAMBIACSO	Cashel	1292	898	12	4	42.25	12.25	48	144	0	16	15.7571429	8.941667	87.5	0.019663	0.275278	22.428571	11.91667	4.916667		0.428571	14.285714	16	14	12	0	8.333333	75	16.666667	0	3.083333	7.571429	0	0
4	01786/88	MORARANO	Cashel	1292	898	10	6	20.25	2.25	60	100	0	16	14.16875	7.835714	100	0.016475	0.263593	22.892857	8.533333	4.666667		0.75	18.75	16	16	14	0	13.33333	66.666667	20	0	3.066667	6.857143	0	0
5	01447/84	MT.PROVINCE	Cashel	1292	898	13	4	56.25	12.25	52	156	0	16	13.53125	7.966667	100	0.0151	0.241595	21.3	12.13333	4.466667		0.4375	6.25	16	16	12	0	0	93.333333	6.6666667	0	3.066667	5.769231	0	0
5	01448/84	BENQUET	Cashel	1292	898	13	8	56.25	0.25	104	104	0	16	12.7875	7.976923	100	0.013027	0.208428	20.40625	11	4.125	0.4375	31.25	16	16	13	0	0	87.5	12.5	0	3.125		6.666667	0	0
5	01515/85	DATHIEN	Cashel	1292	898	15	9	90.25	2.25	135	105	0	16	13.925	8.430769	100	0.015726	0.251619	21.733333	9.25	3.8125	0.125	37.5	16	16	13	0		6.666667	66.666667	20	6.666667	3.266667	7.230769	0	0
5	01517/85	HOTHIEN	Cashel	1292	898	14	10	72.25	6.25	140	84	0	16	12.7625	7.628571	100	0.013044	0.208709	20.866667	10.0625	4.9375	0.25	25	16	16	14	0	18.75	62.5	18.75	0	3		7.142857	0	0
5	01519/85	LANGHANH	Cashel	1292	898	15	5	90.25	6.25	75	165	0	16	13.7266667	7.772727	100	0.015042	0.22563	20.923077	11.58333	4.75		0.375	12.5	16	15	11	0	0	92.307692	7.6923077	0	3.076923	7.076923	0	0
5	01521/85	NONGKRATING	Cashel	1292	898	14	4	72.25	12.25	56	168	0	16	14.9357143	6.683333	93.75	0.0177	0.247803	22.857143	11.30769	5.230769		0.466667	0	16	14	12	0	15.38462	61.538462	23.076923	0	3.076923	6.363636	0	0
5	01523/85	DOIINTHANON	Cashel	1292	898	13	5	56.25	6.25	65	143	0	16	14.4125	7.377778	100	0.01695	0.271202	22.071429	11.15385	5.384615		0.125	0	16	16	9	0	9.090909	90.909091	0	0	2.909091	5.666667	0	0
5	01525/85	NAMNOW	Cashel	1292	898	15	6	90.25	2.25	90	150	0	16	13.84375	6.653333	100	0.015181	0.242898	21.96875	12.6	5.4		0.1875	6.25	16	16	15	0	0	73.333333	26.666667	0	3.266667	6.533333	0	0
5	01572/85	COTOMINES	Cashel	1292	898	15	10	90.25	6.25	150	90	0	16	12.58125	8.792857	100	0.012735	0.203755	18.8	12.4375	4.125	0.75	6.25	16	16	14	0	18.75	81.25	0	0	2.8125	6.5	0	0	0
5	01624/86	BAOSHAN(1)	Cashel	1292	898	14	8	72.25	0.25	112	112	0	16	8.04666667	3.4125	100	0.006589	0.09883		3.8	5		0.857143	0	16	15	16	0	13.33333	86.666667	0	0	0	2.866667	6.4	0	0
5	01632/86	CEHENG	Cashel	1292	898	14	7	72.25	0.25	98	126	0	16	6.75833333	2.626667	93.75	0.005209	0.062504		3	5.538462		0.8	0	16	12	15	0	15.38462	84.615385	0	0	0	2.846154	6.307692	0	0
5	01633/86	SHANGSI	Cashel	1292	898	15	7	90.25	0.25	105	135	0	16	7.80714286	3.10625	100	0.00574	0.080359	24.5	4.6	4.733333		0.5	0	16	14	16	0	0	92.857143	7.1428571	0	0	3.071429	6.785714	0	0
5	01636/86	JINGDUNG(2)	Cashel	1292	898	13	7	56.25	0.25	91	117	0	16	13.45	6.845455	87.5	0.014862	0.208073	24.166667	9	4.461538		0.285714	21.428571	16	14	11	0	0	64.285714	21.428571	14.28571	3.5	6.272727	0	0
5	01637/86	JINGHONG	Cashel	1292	898	14	9	72.25	2.25	126	98	0	16	14.6533333	8.02	93.75	0.017006	0.255093	21.966667	10.13333	5.6		0.466667	13.333333	16	15	15	0	6.666667	53.333333	40	0	3.333333	6.866667	0	0
5	01638/86	LANCANG	Cashel	1292	898	15	8	90.25	0.25	120	120	0	16	14.18125	6.8	100	0.016651	0.266422	23.035714	9.8	6		0.375	0	16	16	15	0	0	33.333333	66.666667	0	3.666667	6.357143	0	0
5	01680/86	DAKHA	Cashel	1292	898	14	5	72.25	6.25	70	154	0	16	13.93125	8.458333	100	0.015623	0.249968	22.357143	8.928571	5.142857		0.5625	43.75	16	16	12	7.142857	7.142857	85.714286	0	0	2.785714	6.571429	0	7.14286
5	01772/88	ZOKHUA	Cashel	1292	898	15	4	90.25	12.25	60	180	0	16	13.3142857	6.207143	87.5	0.015967	0.223532	20.846154	9.357143	6		0.285714	0	16	14	14	0	7.142857	85.714286	7.1428571	0	3	6.384615	0	0
5	01773/88	AUNGBAN	Cashel	1292	898	13	10	56.25	6.25	130	78	0	16	11.95	5.690909	87.5	0.012261	0.171654	20.307692	8.357143	5		0.428571	0	16	14	11	0	0	64.285714	35.714286	0	3.357143	5.818182	0	0
5	01777/88	JOHNMEIKLECSO	Cashel	1292	898	13	9	56.25	2.25	117	91	0	16	12.85625	7.6	100	0.013381	0.214089	20.653846	11.66667	5		0.375	12.5	16	16	11	0	16.66667	66.666667	16.666667	0	3	7.538462	0	0
5	01778/88	ZAMBIACSO	Cashel	1292	898	13	6	56.25	2.25	78	130	0	16	14.70625	8.281818	100	0.017196	0.275134	21.625	14.375	5.125	0.125	0	16	16	11	0	0	87.5	12.5	0	3.125		7.666667	0	0
5	01786/88	MORARANO	Cashel	1292	898	14	6	72.25	2.25	84	140	0	16	14.3142857	8.1	93.75	0.01631	0.228338	22.785714	11.07143	5.214286		0.357143	13.333333	16	14	10	0	0	71.428571	28.571429	0	3.285714	7.5	0	0
6	01447/84	MT.PROVINCE	Cashel	1292	898	2	7	12.25	0.25	14	18	0	16	12.9625	8.384615	100	0.013414	0.21463	21.3	12.14286	4.714286		0.3125	31.25	16	16	13	0	0	75	25	0	3.25	6.714286	0	0
6	01448/84	BENQUET	Cashel	1292	898	3	13	6.25	30.25	39	9	0	16	13.2875	8.05	100	0.014715	0.235438	20.321429	10.375	3.8125	0.5625	25	16	16	14	6.25	12.5	68.75	12.5	0	2.875		6.933333	0	0
6	01515/85	DATHIEN	Cashel	1292	898	3	8	6.25	0.25	24	24	0	16	14.7266667	7.978571	93.75	0.01712	0.256802	22.3	10.2	4.533333		0.333333	33.333333	16	15	14	0	7.142857	50	42.857143	0	3.357143	7.214286	0	0
6	01517/85	HOTHIEN	Cashel	1292	898	1	9	20.25	2.25	9	7	0	16	13.98	7.638462	100	0.015578	0.233672	22.357143	10	4.076923		0.375	31.25	16	15	13	0	0	53.333333	46.666667	0	3.466667	6.8	0	6.66667
6	01519/85	LANGHANH	Cashel	1292	898	2	11	12.25	12.25	22	10	0	16	15.10625	7.976923	100	0.018573	0.297162	22.875	11.06667	3.866667		0.4375	31.25	16	16	13	0	6.666667	53.333333	33.333333	6.666667	3.4	7.071429	0	0
6	01521/85	NONGKRATING	Cashel	1292	898	3	9	6.25	2.25	27	21	0	16	13.7375	6.08	100	0.015315	0.245041	22.714286	10.53333	6.066667		0.125	0	16	16	15	6.666667	13.33333	73.333333	6.6666667	0	2.8	6.666667	0	0
6	01523/85	DOIINTHANON	Cashel	1292	898	3	10	6.25	6.25	30	18	0	16	14.6066667	7.007692	100	0.017076	0.256145	23.2	12.57143	5.571429		0.3125	0	16	15	13	0	0	85.714286	14.285714	0	3.142857	6.538462	0	0
6	01525/85	NAMNOW	Cashel	1292	898	1	11	20.25	12.25	11	5	0	16	13.2875	6.306667	100	0.014825	0.237194	23.833333	10.33333	5.2		0.4375	0	16	16	15	0	12.5	56.25	31.25	0	3.1875	6.466667	0	0
6	01572/85	COTOMINES	Cashel	1292	898	1	8	20.25	0.25	8	8	0	16	13.95625	8.871429	100	0.016112	0.257798	19.84375	12.13333	4		0.5625	12.5	16	16	14	0	7.142857	92.857143	0	0	2.928571	6.714286	0	0
6	01624/86	BAOSHAN(1)	Cashel	1292	898	3	12	6.25	20.25	36	12	0	16	7.85	3.566667	100	0.005608	0.089734		4	5.133333		0.875	0	16	16	15	0	25	62.5	12.5	0	2.875	6.4	0	0
6	01632/86	CEHENG	Cashel	1292	898	2	10	12.25	6.25	20	12	0	16	9	3.325	100	0.007567	0.105941	23	3.928571	5.214286		0.6875	0	16	14	16	0	0	50	50	0	3.5	6.071429	0	0
6	01633/86	SHANGSI	Cashel	1292	898	1	13	20.25	30.25	13	3	0	16	8.18571429	3.35	87.5	0.005649	0.079081	26.5	4.428571	4.571429		0.285714	0	16	14	14	0	7.142857	85.714286	7.1428571	0	3	6.642857	0	0
6	01636/86	JINGDUNG(2)	Cashel	1292	898	3	7	6.25	0.25	21	27	0	16	12.54	6.48	93.75	0.014055	0.210821	23.633333	8.933333	5.933333		0.533333	0	16	15	15	0	6.666667	73.333333	20	0	3.133333	6.266667	0	0
6	01637/86	JINGHONG	Cashel	1292	898	3	11	6.25	12.25	33	15	0	16	14.2625	7.569231	100	0.017799	0.284778	22.366667	11.28571	5.928571		0.4	0	16	16	13	7.142857	7.142857	64.285714	14.285714	7.142857	3.071429	6.357143	0	0
6	01638/86	LANCANG	Cashel	1292	898	2	9	12.25	2.25	18	14	0	16	13.8066667	6.815385	93.75	0.016673	0.250091	24.269231	10	5.533333		0.8	13.333333	16	15	13	0	0	66.666667	33.333333	0	3.333333	6.076923	0	0
6	01680/86	DAKHA	Cashel	1292	898	2	12	12.25	20.25	24	8	0	16	14.15	8.007143	100	0.016769	0.268296	22.714286	10.06667	5		0.4375	6.25	16	16	14	0	6.666667	66.666667	26.666667	0	3.2	6.571429	0	0
6	01772/88	ZOKHUA	Cashel	1292	898	2	8	12.25	0.25	16	16	0	16	12.41875	5.70625	100	0.012869	0.2059	21.066667	10.625	5.4375	0.375	0	16	16	16	0	6.25	87.5	6.25	0	3	6.375	0	6.25	0
6	01773/88	AUNGBAN	Cashel	1292	898	1	12	20.25	20.25	12	4	0	16	12.25	5.891667	87.5	0.015159	0.212229	22.958333	9.666667	5.083333		0.642857	0	16	14	12	0	0	69.230769	30.769231	0	3.307692	6.083333	0	0
6	01777/88	JOHNMEIKLECSO	Cashel	1292	898	1	7	20.25	0.25	7	9	0	16	14.3933333	7.972727	100	0.016442	0.246627	21.5	13.64286	4.5		0.375	6.25	16	15	11	0	0	86.666667	13.333333	0	3.133333	6.727273	0	0
6	01778/88	ZAMBIACSO	Cashel	1292	898	1	10	20.25	6.25	10	6	0	16	15.2066667	8.635714	93.75	0.018291	0.274358	22.233333	14.4	6.133333		0.2	0	16	15	14	0	0	93.333333	6.6666667	0	3.066667	8.142857	0	0
6	01786/88	MORARANO	Cashel	1292	898	2	13	12.25	30.25	26	6	0	16	14.16	8	93.75	0.016577	0.248653	23.1	9.466667	4.933333		0.333333	26.666667	16	15	15	0	6.666667	60	33.333333	0	3.266667	6.933333	0	0

Plot Data Set - Cashel

STEM3	STEM4	STEM5	STEM6	STEM7	STEM8	STEM9	BRAARATIO	FO_INDEX	INTNODE	CROWNPCT	FO_POS	TOP	VTREE	MF	FF	CL	CC	OC	LOTH	APID	DG	HG	PILOCORR	VHA
0	6.6666667	6.666667	46.666667	26.666667	13.333333	0		164.93274	0.60998		0.26375	6.25	0.062615	1	1	1	1	1.3125	6.25	31.25	13.196377		20.969134	85.891104
0	6.25	62.5	31.25	0	0			367.72982	0.89979		0.102727	0	0.027611	1	1	1	1	1.375	0	6.25	10.291987		24.188253	37.875701
0	0	6.666667	53.333333	40	0	0		173.38701	0.68495		0.098889	0	0.019987	1	1	1	1	1.5625	0	37.5	8.7893117		23.849484	25.704044
0	0	6.666667	80	13.333333	0	0		26.15022	0.64681		0.275714	0	0.02417	1	1	1	1	2.75	6.25	6.25	9.9199609		25.413926	33.155401
18.75	0	37.5	31.25	12.5	0			29.742257	0.70283		0.321667	0	0.050664	1	1	1	1	2.9375	0	6.25	13.369508		23.196195	69.49787
0	7.1428571	14.28571	35.714286	21.428571	21.428571	0		52.795301	0.67649		0.4275	6.25	0.05346	1	1	1	1	1.86667	0	0	13.505382		24.090531	68.750548
0	0	21.42857	35.714286	35.714286	0	0		23.51941	0.75221		0.4775	6.25	0.080646	1	1	1	1	1.75	6.25	12.5	15.304656		23.305206	110.62542
0	0	14.28571	42.857143	21.428571	21.428571	0		27.854497	0.84169		0.33	6.25	0.063007	1	1	1	1	1.5625	12.5	0	13.124		22.261131	86.429513
0	0	0	25	25	50	0		264.24923	0.58569		0.2	6.25	0.042901	1	1	1	1	2.4	12.5	31.25	11.974027		20.835062	55.17148
0	0	38.46154	61.538462	0	0	0		444.96429	0.59397		0.1	0	0.053262	1	1	1	1	2.2	12.5	12.5	14.297063		23.734939	68.495804
0	0	12.5	25	25	25	12.5		150.48488	0.64901		0.374286	25	0.069435	1	1	1	1	1	18.75	0	13.839412		22.213811	89.293582
0	0	0	21.428571	28.571429	21.428571	28.571429		16.033668	0.73991		0.575	12.5	0.086158	1	1	1	1	1.78571	0	0	15.819382		22.575363	103.41353
0	0	14.28571	21.428571	28.571429	35.714286	0		43.475582	0.93724		0.437778	6.25	0.066858	1	1	1	1	1.4375	6.25	25	14.480202		22.968525	91.711865
0	23.076923	7.692308	38.461538	30.769231	0	0		33.28611	0.62642		0.365	6.25	0.064126	1	1	1	1	1.625	12.5	18.75	13.86281		21.302436	87.964249
0	0	0	50	41.666667	0	8.3333333		35.021936	0.68797		0.34	0	0.05615	1	1	1	1	1.375	18.75	25	12.876141		20.334006	77.023014
0	7.6923077	0	7.6923077	30.769231	53.846154	0		20.460789	1.01857		0.49	0	0.065882	1	1	1	1	1.75	18.75	18.75	14.147482		21.777074	90.373777
0	0	7.142857	14.285714	42.857143	28.571429	7.1428571		13.444539	0.79447		0.6075	0	0.055683	1	1	1	1	1.375	12.5	12.5	12.884826		20.806654	76.382665
0	0	0	15.384615	61.538462	23.076923	0		26.101729	0.74813		0.364	25	0.065995	1	1	1	1	1.1875	6.25	0	13.836305		20.961257	84.869637
0	0	9.090909	54.545455	27.272727	9.0909091	0		18.818627	0.56655		0.338333	12.5	0.062992	1	1	1	1	1.4	12.5	6.25	15.009164		22.943347	75.607101
0	11.111111	33.33333	33.333333	22.222222	0	0		18.375	0.61989		0.43	18.75	0.067112	1	1	1	1	1	18.75	6.25	14.687707		22.187136	92.060117
6.666667	0	6.666667	13.333333	66.666667	6.6666667	0		37.125	0.50433		0.17	6.25	0.05381	1	1	1	1	1.66667	0	43.75	13.900157		21.974413	73.813446
0	0	50	50	0	0			18.710315	0.73649		0.512727	0	0.061677	1	1	1	1	1.1875	12.5	50	12.730991		18.724252	84.605392
0	0	0	60	40	0	0		139.5528	0.82956		0.117778	0	0.020711	1	1	1	1	1.85714	0	6.25	9.1572922			26.63486
0	0	0	69.230769	30.769231	0	0		275.98695	0.78269		0.044444	0	0.018945	1	1	1	1	1.26667	0	18.75	8.1419695			19.490334
0	0	0	21.428571	78.571429	0	0		327.98611	0.59767		0.1	0	0.01944	1	1	1	1	2.8	0	0	8.5471382		24.590946	23.333053
0	18.181818	9.090909	18.181818	36.363636	18.181818	0		42.916667	0.74078		0.293333	0	0.05687	1	1	1	1	2.92857	18.75	6.25	13.753441		24.162342	68.26017
0	0	0	26.666667	60	13.333333	0		19.826391	0.77918		0.447143	0	0.070173	1	1	1	1	1.93333	0	6.25	14.711991		22.032043	90.243662
0	0	0	71.428571	21.428571	7.1428571	0		27.824704	0.6493		0.328333	0	0.061859	1	1	1	1	2.5625	12.5	18.75	14.557708		23.102425	84.855202
0	0	7.142857	14.285714	50	21.428571	0		27.240334	1.01471		0.392857	6.25	0.074896	1	1	1	1	1.4	12.5	6.25	14.100997		22.413317	102.73766
0	0	23.07692	30.769231	30.769231	15.384615	0		240	0.62053		0.033333	0	0.055199	1	1	1	1	2.71429	6.25	12.5	14.2552		20.844618	66.254049
0	18.181818	0	63.636364	18.181818	0	0		60.538462	0.62097		0.146	0	0.040102	1	1	1	1	2.85714	18.75	0	12.491969		20.187576	48.133649
0	0	0	7.6923077	46.153846	30.769231	15.384615		25.360536	0.67428		0.412	12.5	0.058806	1	1	1	1	1.4375	12.5	0	13.049832		20.650041	80.667034
0	0	0	0	50	33.333333	16.666667		39.512195	0.55943		0.41	0	0.076647	1	1	1	1	1.6875	37.5	0	14.793812		21.694279	105.13963
0	0	0	0	70	10	20		17.285485	0.75153		0.482	6.25	0.072181	1	1	1	1	1.69231	25	0	14.407637		22.826083	86.636686
0	0	7.142857	28.571429	50	14.285714	0		25.701557	0.68086		0.416	6.25	0.058684	1	1	1	1	1.125	12.5	31.25	13.066321		21.248347	80.499155
0	0	6.666667	40	20	20	13.333333		23.599573	0.9232		0.4375	6.25	0.062996	1	1	1	1	1.35714	6.25	25	13.685028		20.365485	86.414246
0	0	0	7.1428571	64.285714	28.571429	0		37.849877	0.85138		0.41	0	0.070576	1	1	1	1	1.93333	6.25	12.5	14.761188		22.370785	90.760863
0	0	6.666667	6.6666667	60	6.6666667	13.333333		23.415584	0.77841		0.4	18.75	0.061577	1	1	1	1	1.33333	0	6.25	14.080743		22.398565	79.188281
0	0	0	28.571429	42.857143	21.428571	7.1428571		50.115688	0.7375		0.32	6.25	0.075961	1	1	1	1	1.3125	6.25	6.25	15.374614		22.973783	104.19903
0	0	6.666667	40	40	6.6666667	6.6666667		13.333333	0.53285		0.325	6.25	0.053063	1	1	1	1	1.26667	0	31.25	13.961331		22.763084	72.788168
0	0	0	53.846154	38.461538	7.6923077	0		14.076034	0.52182		0.311667	6.25	0.063127	1	1	1	1	1.5	6.25	25	14.742297		23.261934	81.182163
0	0	20	26.666667	40	13.333333	0		107.1147	0.59722		0.278333	0	0.052453	1	1	1	1	1.25	6.25	12.5	13.73597		23.822493	71.951656
0	0	7.142857	28.571429	50	14.285714	0		58.522603	0.71646		0.401429	6.25	0.073494	1	1	1	1	1	6.25	37.5	14.320156		19.857711	100.81549
0	6.6666667	0	40	53.333333	0	0		64.849492	0.78789		0.110909	0	0.018514	1	1	1	1	1.375	6.25	25	8.4486315			25.396009
0	7.1428571	14.28571	50	21.428571	7.1428571	0		56.291667	0.82571		0.09375	0	0.022259	1	1	1	1	1.8125	0	12.5	9.8137658		22.825416	26.717442
0	0	7.142857	42.857143	28.571429	21.428571	0		17.851552	0.6498		0.15	0	0.017008	1	1	1	1	2.14286	0	12.5	8.4788813		26.373359	20.414539
0	0	26.66667	26.666667	40	6.6666667	0		112.11024	0.68456		0.227143	0	0.051137	1	1	1	1	2.28571	0	12.5	13.374553		23.542834	65.762824
0	7.1428571	0	57.142857	28.571429	0	7.1428571		60.468922	0.64601		0.18	6.25	0.075181	1	1	1	1	1.9375	6.25	18.75	15.050851		22.430568	103.12867
0	7.6923077	7.692308	61.538462	15.384615	7.6923077	0		129.22413	0.66682		0.24875	6.25	0.061384	1	1	1	1	2.4	12.5	18.75	14.56704		24.325	78.94024
0	0	0	50	42.857143	7.1428571	0		192.06982	0.77257		0.236	6.25	0.072308	1	1	1	1	1.4375	6.25	6.25	14.608816		22.784685	99.187549
0	0	37.5	50	6.25	0			64.388206	0.50619		0.261667	6.25	0.042108	1	1	1	1	3.1875	0	50	12.797827		20.988952	57.761626
0	0	8.333333	75	16.666667	0	0		64.254915	0.56438		0.18125	0	0.053854	1	1	1	1	3	12.5	12.5	13.890104		22.866728	64.639437
0	9.0909091	9.090909	9.0909091	54.545455	9.0909091	9.0909091		30.228046	0.57654		0.414	6.25	0.069048	1	1	1	1	1.25	25	0	14.465799		21.546199	88.796025
0	0	0	7.1428571	14.285714	35.714286	42.857143		22.370318	0.57937		0.453333	0	0.081008	1	1	1	1	2.53333	6.25	0	15.257413		22.339523	104.17737
0	6.6666667	0	20	46.666667	20	6.6666667		48.47671	0.88398		0.3625	0	0.068609	1	1	1	1	1.53333	0	0	14.525105		23.128989	88.231454